

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 August 2003 (28.08.2003)

PCT


(10) International Publication Number
WO 03/070910 A2

(51) International Patent Classification ⁷ :	C12N	60/440,129	15 January 2003 (15.01.2003)	US
(21) International Application Number:	PCT/US03/05022	(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:		
(22) International Filing Date:	20 February 2003 (20.02.2003)	US	60/393,796 (CON)	
(25) Filing Language:	English	Filed on	3 July 2002 (03.07.2002)	
(26) Publication Language:	English	US	60/399,348 (CON)	
		Filed on	29 July 2002 (29.07.2002)	
		US	10/306,747 (CON)	
		Filed on	27 November 2002 (27.11.2002)	
(30) Priority Data:		US	10/287,949 (CON)	
60/358,580	20 February 2002 (20.02.2002)	US	4 November 2002 (04.11.2002)	
60/363,124	11 March 2002 (11.03.2002)	US	PCT/US02/17674 (CON)	
PCT/US02/17674	29 May 2002 (29.05.2002)	US	29 May 2002 (29.05.2002)	
60/386,782	6 June 2002 (06.06.2002)	US	60/358,580 (CON)	
60/393,796	3 July 2002 (03.07.2002)	US	20 February 2002 (20.02.2002)	
60/399,348	29 July 2002 (29.07.2002)	US	60/363,124 (CON)	
60/406,784	29 August 2002 (29.08.2002)	US	11 March 2002 (11.03.2002)	
60/408,378	5 September 2002 (05.09.2002)	US	60/386,782 (CON)	
60/409,293	9 September 2002 (09.09.2002)	US	6 June 2002 (06.06.2002)	
10/287,949	4 November 2002 (04.11.2002)	US	60/406,784 (CON)	
10/306,747	27 November 2002 (27.11.2002)	US	29 August 2002 (29.08.2002)	

[Continued on next page]

(54) Title: RNA INTERFERENCE MEDIATED INHIBITION OF VASCULAR ENDOTHELIAL GROWTH FACTOR AND VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR GENE EXPRESSION USING SHORT INTERFERING NUCLEIC ACID (siNA)

A375 24h 36B4 VEGFR1 mRNA Expression



(57) Abstract: The present invention concerns methods and reagents useful in modulating vascular endothelial growth factor (VEGF, VEGF-B, VEGF-C, VEGF-D) and/or vascular endothelial growth factor receptor (e.g., VEGFr1, VEGFr2, and/or VEGFr3) gene expression in a variety of applications, including use in therapeutic, diagnostic, target validation, and genomic discovery applications. Specifically, the invention relates to small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating RNA interference (RNAi) against VEGF and/or VEGFr gene expression and/or activity. The small nucleic acid molecules are useful in the diagnosis and treatment of cancer, proliferative diseases, and any other disease or condition that responds to modulation of VEGF and/or VEGFr expression or activity.

WO 03/070910 A2



US 60/408,378 (CON)
Filed on 5 September 2002 (05.09.2002)
US 60/409,293 (CON)
Filed on 9 September 2002 (09.09.2002)
US 60/440,129 (CON)
Filed on 15 January 2003 (15.01.2003)

(71) **Applicant** (*for all designated States except US*): **RI-BOZYME PHARMACEUTICALS, INCORPORATED** [US/US]; 2950 Wilderness Place, Boulder, CO 80301 (US).

(72) **Inventors; and**

(75) **Inventors/Applicants** (*for US only*): **McSWIGGEN, James** [US/US]; 4866 Franklin Drive, Boulder, CO 80301 (US). **BEIGELMAN, Leonid** [US/US]; 5530 Colt Drive, Longmont, CO 80503 (US). **PAVCO, Pamela** [US/US]; 705 Barberry Circle, Lafayette, CO 80026 (US).

(74) **Agent: TERPSTRA, Anita, J.**; McDonnell Boehnen Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).

(81) **Designated States (national)**: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional)**: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

RNA INTERFERENCE MEDIATED INHIBITION OF VASCULAR ENDOTHELIAL GROWTH FACTOR AND VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR GENE EXPRESSION USING SHORT INTERFERING NUCLEIC ACID (siNA)

5 This invention claims the benefit of McSwiggen, USSN 60/393,796 filed July 3, 2002,
of McSwiggen, USSN 60/399,348 filed July 29, 2002, of Pavco, USSN 10/306,747, filed
November 27, 2002, which claims the benefit of Pavco USSN 60/334461, filed November
30, 2001, of Pavco, USSN 10/287,949 filed November 4, 2002, of Pavco, PCT/US02/17674
10 filed May 29, 2002, of Beigelman USSN 60/358,580 filed February 20, 2002, of Beigelman
USSN 60/363,124 filed March 11, 2002, of Beigelman USSN 60/386,782 filed June 6, 2002,
of Beigelman USSN 60/406,784 filed August 29, 2002, of Beigelman USSN 60/408,378
filed September 5, 2002, of Beigelman USSN 60/409,293 filed September 9, 2002, and of
Beigelman USSN 60/440,129 filed January 15, 2003. These applications are hereby
incorporated by reference herein in their entireties, including the drawings.

15 **Field Of The Invention**

The present invention concerns compounds, compositions, and methods for the study, diagnosis, and treatment of conditions and diseases that respond to the modulation of vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor receptor (e.g., VEGFr1, VEGFr2 and/or VEGFr3) gene expression and/or activity. The present invention also concerns compounds, compositions, and methods relating to conditions and diseases that respond to the modulation of expression and/or activity of genes involved in VEGF and VEGF receptor pathways. Specifically, the invention relates to small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating RNA interference (RNAi) against VEGF and VEGF receptor gene expression.

Background Of The Invention

The following is a discussion of relevant art pertaining to RNAi. The discussion is provided only for understanding of the invention that follows. The summary is not an admission that any of the work described below is prior art to the claimed invention.

5 RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire *et al.*, 1998, *Nature*, 391, 806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-
10 conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that
15 specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2',5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.

20 The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein *et al.*, 2001, *Nature*, 409, 363). Short interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes (Elbashir *et al.*,
25 2001, *Genes Dev.*, 15, 188). Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having

sequence complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188).

RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, were the first to observe RNAi in *C. elegans*. Wianny and Goetz, 1999, *Nature Cell Biol.*, 2, 70, describe RNAi mediated by dsRNA in mouse embryos. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells. Recent work in *Drosophila* embryonic lysates (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877) has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity. These studies have shown that 21-nucleotide siRNA duplexes are most active when containing 3'-terminal dinucleotide overhangs. Furthermore, complete substitution of one or both siRNA strands with 2'-deoxy (2'-H) or 2'-O-methyl nucleotides abolishes RNAi activity, whereas substitution of the 3'-terminal siRNA overhang nucleotides with 2'-deoxy nucleotides (2'-H) was shown to be tolerated. Single mismatch sequences in the center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end of the guide sequence (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). Other studies have indicated that a 5'-phosphate on the target-complementary strand of a siRNA duplex is required for siRNA activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309).

Studies have shown that replacing the 3'-terminal nucleotide overhanging segments of a 21-mer siRNA duplex having two -nucleotide 3'-overhangs with deoxyribonucleotides does not have an adverse effect on RNAi activity. Replacing up to four nucleotides on each end of the siRNA with deoxyribonucleotides has been reported to be well tolerated, whereas complete substitution with deoxyribonucleotides results in no RNAi activity (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). In addition, Elbashir *et al.*, *supra*, also report that substitution of siRNA with 2'-O-methyl nucleotides completely abolishes RNAi activity. Li *et al.*,

International PCT Publication No. WO 00/44914, and Beach *et al.*, International PCT Publication No. WO 01/68836 preliminarily suggest that siRNA may include modifications to either the phosphate-sugar backbone or the nucleoside to include at least one of a nitrogen or sulfur heteroatom, however, neither application postulates to what extent such
5 modifications would be tolerated in siRNA molecules, nor provides any further guidance or examples of such modified siRNA. Kreutzer *et al.*, Canadian Patent Application No. 2,359,180, also describe certain chemical modifications for use in dsRNA constructs in order to counteract activation of double-stranded RNA-dependent protein kinase PKR, specifically 2'-amino or 2'-O-methyl nucleotides, and nucleotides containing a 2'-O or 4'-C
10 methylene bridge. However, Kreutzer *et al.* similarly fails to provide examples or guidance as to what extent these modifications would be tolerated in siRNA molecules.

Parrish *et al.*, 2000, *Molecular Cell*, 6, 1977-1087, tested certain chemical modifications targeting the unc-22 gene in *C. elegans* using long (>25 nt) siRNA transcripts. The authors describe the introduction of thiophosphate residues into these siRNA transcripts
15 by incorporating thiophosphate nucleotide analogs with T7 and T3 RNA polymerase and observed that RNAs with two phosphorothioate modified bases also had substantial decreases in effectiveness as RNAi. Further, Parrish *et al.* reported that phosphorothioate modification of more than two residues greatly destabilized the RNAs *in vitro* such that interference activities could not be assayed. *Id.* at 1081. The authors also tested certain
20 modifications at the 2'-position of the nucleotide sugar in the long siRNA transcripts and found that substituting deoxynucleotides for ribonucleotides produced a substantial decrease in interference activity, especially in the case of Uridine to Thymidine and/or Cytidine to deoxy-Cytidine substitutions. *Id.* In addition, the authors tested certain base modifications, including substituting, in sense and antisense strands of the siRNA, 4-thiouracil, 5-
25 bromouracil, 5-iodouracil, and 3-(aminoallyl)uracil for uracil, and inosine for guanosine. Whereas 4-thiouracil and 5-bromouracil substitution appeared to be tolerated, Parrish reported that inosine produced a substantial decrease in interference activity when incorporated in either strand. Parrish also reported that incorporation of 5-iodouracil and 3-(aminoallyl)uracil in the antisense strand resulted in a substantial decrease in RNAi activity
30 as well.

The use of longer dsRNA has been described. For example, Beach *et al.*, International PCT Publication No. WO 01/68836, describes specific methods for attenuating gene expression using endogenously-derived dsRNA. Tuschl *et al.*, International PCT Publication No. WO 01/75164, describe a *Drosophila in vitro* RNAi system and the use of
5 specific siRNA molecules for certain functional genomic and certain therapeutic applications; although Tuschl, 2001, *Chem. Biochem.*, 2, 239-245, doubts that RNAi can be used to cure genetic diseases or viral infection due to the danger of activating interferon response. Li *et al.*, International PCT Publication No. WO 00/44914, describe the use of specific dsRNAs for attenuating the expression of certain target genes. Zernicka-Goetz *et al.*,
10 International PCT Publication No. WO 01/36646, describe certain methods for inhibiting the expression of particular genes in mammalian cells using certain dsRNA molecules. Fire *et al.*, International PCT Publication No. WO 99/32619, describe particular methods for introducing certain dsRNA molecules into cells for use in inhibiting gene expression. Plaetinck *et al.*, International PCT Publication No. WO 00/01846, describe certain methods
15 for identifying specific genes responsible for conferring a particular phenotype in a cell using specific dsRNA molecules. Mello *et al.*, International PCT Publication No. WO 01/29058, describe the identification of specific genes involved in dsRNA-mediated RNAi. Deschamps Depaillette *et al.*, International PCT Publication No. WO 99/07409, describe specific compositions consisting of particular dsRNA molecules combined with certain anti-
20 viral agents. Waterhouse *et al.*, International PCT Publication No. 99/53050, describe certain methods for decreasing the phenotypic expression of a nucleic acid in plant cells using certain dsRNAs. Driscoll *et al.*, International PCT Publication No. WO 01/49844, describe specific DNA constructs for use in facilitating gene silencing in targeted organisms.

Others have reported on various RNAi and gene-silencing systems. For example,
25 Parrish *et al.*, 2000, *Molecular Cell*, 6, 1977-1087, describe specific chemically-modified siRNA constructs targeting the unc-22 gene of *C. elegans*. Grossniklaus, International PCT Publication No. WO 01/38551, describes certain methods for regulating polycomb gene expression in plants using certain dsRNAs. Churikov *et al.*, International PCT Publication No. WO 01/42443, describe certain methods for modifying genetic characteristics of an
30 organism using certain dsRNAs. Cogoni *et al.*, International PCT Publication No. WO

01/53475, describe certain methods for isolating a *Neurospora* silencing gene and uses thereof. Reed *et al.*, International PCT Publication No. WO 01/68836, describe certain methods for gene silencing in plants. Honer *et al.*, International PCT Publication No. WO 01/70944, describe certain methods of drug screening using transgenic nematodes as Parkinson's Disease models using certain dsRNAs. Deak *et al.*, International PCT Publication No. WO 01/72774, describe certain *Drosophila*-derived gene products that may be related to RNAi in *Drosophila*. Arndt *et al.*, International PCT Publication No. WO 01/92513 describe certain methods for mediating gene suppression by using factors that enhance RNAi. Tuschl *et al.*, International PCT Publication No. WO 02/44321, describe certain synthetic siRNA constructs. Pachuk *et al.*, International PCT Publication No. WO 00/63364, and Satishchandran *et al.*, International PCT Publication No. WO 01/04313, describe certain methods and compositions for inhibiting the function of certain polynucleotide sequences using certain dsRNAs. Echeverri *et al.*, International PCT Publication No. WO 02/38805, describe certain *C. elegans* genes identified via RNAi. Kreutzer *et al.*, International PCT Publications Nos. WO 02/055692, WO 02/055693, and EP 1144623 B1 describes certain methods for inhibiting gene expression using RNAi. Graham *et al.*, International PCT Publications Nos. WO 99/49029 and WO 01/70949, and AU 4037501 describe certain vector expressed siRNA molecules. Fire *et al.*, US 6,506,559, describe certain methods for inhibiting gene expression in vitro using certain long dsRNA (greater than 25 nucleotide) constructs that mediate RNAi.

SUMMARY OF THE INVENTION

This invention relates to compounds, compositions, and methods useful for modulating the expression of genes, such as those genes associated with angiogenesis and proliferation using short interfering nucleic acid (siNA) molecules. This invention also relates to compounds, compositions, and methods useful for modulating the expression and activity of vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor receptor (e.g., VEGFr1, VEGFr2, VEGFr3) genes, or genes involved in VEGF and/or VEGFr pathways of gene expression and/or VEGF activity by RNA interference (RNAi) using small nucleic acid molecules, such as short interfering nucleic acid (siNA), short

interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules. In particular, the instant invention features small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules and methods used to modulate the expression of VEGF and/or VEGFr genes. A siNA of the invention can be unmodified or chemically-modified. A siNA of the instant invention can be chemically synthesized, expressed from a vector or enzymatically synthesized. The instant invention also features various chemically-modified synthetic short interfering nucleic acid (siNA) molecules capable of modulating VEGF and/or VEGFr gene expression or activity in cells by RNA interference (RNAi). The use of chemically-modified siNA improves various properties of native siNA molecules through increased resistance to nuclease degradation *in vivo* and/or through improved cellular uptake. Further, contrary to earlier published studies, siNA having multiple chemical modifications retains its RNAi activity. The siNA molecules of the instant invention provide useful reagents and methods for a variety of therapeutic, diagnostic, target validation, genomic discovery, genetic engineering, and pharmacogenomic applications.

In one embodiment, the invention features one or more siNA molecules and methods that independently or in combination modulate the expression of gene(s) encoding proteins, such as vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor receptors (e.g., VEGFr1, VEGFr2, VEGFr3), associated with the maintenance and/or development of cancer and other proliferative diseases, such as genes encoding sequences comprising those sequences referred to by GenBank Accession Nos. shown in Table I, referred to herein generally as VEGF and/or VEGFr. The description below of the various aspects and embodiments of the invention is provided with reference to the exemplary VEGF and VEGFr (e.g., VEGFr1, VEGFr2, VEGFr3) genes referred to herein as VEGF and VEGFr respectively. However, the various aspects and embodiments are also directed to other VEGF and/or VEGFr genes, such as mutant VEGF and/or VEGFr genes, splice variants of VEGF and/or VEGFr genes, other VEGF and/or VEGFr ligands and receptors. The various aspects and embodiments are also directed to other genes that are involved in VEGF and/or VEGFr mediated pathways of signal transduction or gene expression that are

involved in the progression, development, and/or maintenance of disease (e.g., cancer). Those additional genes can be analyzed for target sites using the methods described for VEGF and/or VEGFr genes herein. Thus, the inhibition and the effects of such inhibition of the other genes can be performed as described herein.

5 In one embodiment, the invention features a siNA molecule that down-regulates expression of a VEGF gene, for example, wherein the VEGF gene comprises VEGF encoding sequence.

 In one embodiment, the invention features a siNA molecule that down-regulates expression of a VEGFr gene, for example, wherein the VEGFr gene comprises VEGFr
10 encoding sequence.

 In one embodiment, the invention features a siNA molecule having RNAi activity against VEGF and/or VEGFr RNA, wherein the siNA molecule comprises a sequence complementary to any RNA having VEGF and/or VEGFr or other VEGF and/or VEGFr
15 encoding sequence, such as those sequences having GenBank Accession Nos. shown in **Table I**. Chemical modifications as shown in **Tables III and IV** or otherwise described herein can be applied to any siNA construct of the invention.

 In one embodiment, the invention features a siNA molecule having RNAi activity against VEGF and/or VEGFr RNA, wherein the siNA molecule comprises a sequence complementary to any RNA having VEGF and/or VEGFr encoding sequence, such as those
20 sequences having VEGF and/or VEGFr GenBank Accession Nos. shown in **Table I**. Chemical modifications as shown in **Tables III and IV** or otherwise described herein can be applied to any siNA construct of the invention.

 In another embodiment, the invention features a siNA molecule having RNAi activity against a VEGF and/or VEGFr gene, wherein the siNA molecule comprises nucleotide
25 sequence complementary to nucleotide sequence of a VEGF and/or VEGFr gene, such as those VEGF and/or VEGFr sequences having GenBank Accession Nos. shown in **Table I**. In another embodiment, a siNA molecule of the invention includes nucleotide sequence that can interact with nucleotide sequence of a VEGF and/or VEGFr gene and thereby mediate

silencing of VEGF and/or VEGFr gene expression, for example, wherein the siNA mediates regulation of VEGF and/or VEGFr gene expression by cellular processes that modulate the chromatin structure of the VEGF and/or VEGFr gene and prevent transcription of the VEGF and/or VEGFr gene.

5 In another embodiment, the invention features a siNA molecule comprising nucleotide sequence, for example, nucleotide sequence in the antisense region of the siNA molecule that is complementary to a nucleotide sequence or portion of sequence of a VEGF and/or VEGFr gene. In another embodiment, the invention features a siNA molecule comprising a region, for example, the antisense region of the siNA construct, complementary to a
10 sequence or portion of sequence comprising a VEGF and/or VEGFr gene sequence.

In one embodiment, the antisense region of VEGFr1 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 1-427 or 1997-2000. In one embodiment, the antisense region can also comprise sequence having any of SEQ ID NOs. 428-854, 2024-2027, 2032-2035, 2040-2043, 2104-2107, 2109, 2117, 2120-2122,
15 2125-2132, 2137-2140, 2142, 2150, 2152, 2154, 2158-2160, 2164-2166, 2188-2190, 2197, 2199, 2203-2204, 2229, 2231, 2233, 2235, 2237, or 2238. In another embodiment, the sense region of VEGFr1 constructs can comprise sequence having any of SEQ ID NOs. 1-427, 1997-2000, 2009-2016, 2020-2023, 2028-2031, 2036-2039, 2092-2103, 2108, 2114, 2116, 2123-2124, 2133-2136, 2141, 2149, 2151, 2153, 2155-2157, 2161-2163, 2185-2187, 2198,
20 2200-2202, 2228, 2230, 2232, 2234, or 2236. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222.
25 The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, the antisense region of VEGFr2 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 855-1178 or 2001-2004. In one embodiment, the antisense region can also comprise sequence having any of SEQ ID NOs. 1179-1502, 2048-2051, 2056-2059, 2064-2067, 2208-2210, 2214-2216, or 2048-2051.

5 In another embodiment, the sense region of VEGFr2 constructs can comprise sequence having any of SEQ ID NOs. 855-1178, 2001-2004, 2044-2047, 2052-2055, 2060-2063, 2017-2019, 2205-2207, 2211-2213, or 2044-2047. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the

10 antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence

15 of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, the antisense region of VEGFr3 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 1503-1749 or 2005-2008. In one embodiment, the antisense region can also comprise sequence having any of SEQ ID

20 NOs. 1750-1996, 2072-2075, 2080-2083, or 2088-2091. In another embodiment, the sense region of VEGFr3 constructs can comprise sequence having any of SEQ ID NOs. 1503-1749, 2005-2008, 2068-2071, 2076-2079, or 2034-2087. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the

25 antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence

of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, a siNA molecule of the invention comprises any of SEQ ID NOs. 1-2238. The sequences shown in SEQ ID NOs: 1-2238 are not limiting. A siNA molecule
5 of the invention can comprise any contiguous VEGF and/or VEGFr sequence (e.g., about 19 to about 25, or about 19, 20, 21, 22, 23, 24 or 25 contiguous VEGF and/or VEGFr nucleotides).

In yet another embodiment, the invention features a siNA molecule comprising a sequence, for example, the antisense sequence of the siNA construct, complementary to a
10 sequence or portion of sequence comprising sequence represented by GenBank Accession Nos. shown in **Table I**. Chemical modifications in **Tables III and IV** and described herein can be applied to any siRNA construct of the invention.

In one embodiment of the invention a siNA molecule comprises an antisense strand having about 19 to about 29 nucleotides, wherein the antisense strand is complementary to a
15 RNA sequence encoding a VEGF and/or VEGFr protein, and wherein said siNA further comprises a sense strand having about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or 29) nucleotides, and wherein said sense strand and said antisense strand are distinct nucleotide sequences with at least about 19 complementary nucleotides.

In another embodiment of the invention a siNA molecule of the invention comprises
20 an antisense region having about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or 29) nucleotides, wherein the antisense region is complementary to a RNA sequence encoding a VEGF and/or VEGFr protein, and wherein said siNA further comprises a sense region having about 19 to about 29 nucleotides, wherein said sense region and said antisense region comprise a linear molecule with at least about 19 complementary
25 nucleotides.

In one embodiment of the invention a siNA molecule comprises an antisense strand comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof encoding a VEGF and/or VEGFr protein. The siNA further comprises a

sense strand, wherein said sense strand comprises a nucleotide sequence of a VEGF and/or VEGFr gene or a portion thereof.

In another embodiment, a siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof
5 encoding a VEGF and/or VEGFr protein. The siNA molecule further comprises a sense region, wherein said sense region comprises a nucleotide sequence of a VEGF and/or VEGFr gene or a portion thereof.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGF gene. Because VEGF genes can share
10 some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGF genes (and associated receptor or ligand genes) or alternately specific VEGF genes by selecting sequences that are either shared amongst different VEGF targets or alternatively that are unique for a specific VEGF target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGF RNA sequence
15 having homology between several VEGF genes so as to target several VEGF genes (e.g., different VEGF isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGF RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

20 In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGFr gene. Because VEGFr genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGFr genes (and associated receptor or ligand genes) or alternately specific VEGFr genes by selecting sequences that are either shared amongst different
25 VEGFr targets or alternatively that are unique for a specific VEGFr target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGFr RNA sequence having homology between several VEGFr genes so as to target several VEGFr genes (e.g., different VEGFr isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a

sequence that is unique to a specific VEGFr RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGFr gene. Because VEGFr genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGFr genes or alternately specific VEGFr genes by selecting sequences that are either shared amongst different VEGFr targets or alternatively that are unique for a specific VEGFr target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGFr RNA sequence having homology between several VEGFr genes so as to target several VEGFr genes (e.g., VEGFr1, VEGFr2 and/or VEGFr3, different VEGFr isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGFr RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGF gene. Because VEGF genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGF genes or alternately specific VEGF genes by selecting sequences that are either shared amongst different VEGF targets or alternatively that are unique for a specific VEGF target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGF RNA sequence having homology between several VEGF genes so as to target several VEGF genes (e.g., VEGF-A, VEGF-B, VEGF-C and/or VEGF-D, different VEGF isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGF RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, nucleic acid molecules of the invention that act as mediators of the RNA interference gene silencing response are double-stranded nucleic acid molecules. In another embodiment, the siNA molecules of the invention consist of duplexes containing

about 19 base pairs between oligonucleotides comprising about 19 to about 25 (*e.g.*, about 19, 20, 21, 22, 23, 24 or 25) nucleotides. In yet another embodiment, siNA molecules of the invention comprise duplexes with overhanging ends of about 1 to about 3 (*e.g.*, about 1, 2, or 3) nucleotides, for example, about 21-nucleotide duplexes with about 19 base pairs and 3'-terminal mononucleotide, dinucleotide, or trinucleotide overhangs.

In one embodiment, the invention features one or more chemically-modified siNA constructs having specificity for VEGF and/or VEGFr expressing nucleic acid molecules, such as RNA encoding a VEGF and/or VEGFr protein. Non-limiting examples of such chemical modifications include without limitation phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, and terminal glyceryl and/or inverted deoxy abasic residue incorporation. These chemical modifications, when used in various siNA constructs, are shown to preserve RNAi activity in cells while at the same time, dramatically increasing the serum stability of these compounds. Furthermore, contrary to the data published by Parrish *et al.*, *supra*, applicant demonstrates that multiple (greater than one) phosphorothioate substitutions are well-tolerated and confer substantial increases in serum stability for modified siNA constructs.

In one embodiment, a siNA molecule of the invention comprises modified nucleotides while maintaining the ability to mediate RNAi. The modified nucleotides can be used to improve *in vitro* or *in vivo* characteristics such as stability, activity, and/or bioavailability. For example, a siNA molecule of the invention can comprise modified nucleotides as a percentage of the total number of nucleotides present in the siNA molecule. As such, a siNA molecule of the invention can generally comprise about 5% to about 100% modified nucleotides (*e.g.*, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% modified nucleotides). The actual percentage of modified nucleotides present in a given siNA molecule will depend on the total number of nucleotides present in the siNA. If the siNA molecule is single stranded, the percent modification can be based upon the total number of nucleotides present in the single stranded siNA molecules. Likewise, if the siNA molecule is double stranded, the percent

modification can be based upon the total number of nucleotides present in the sense strand, antisense strand, or both the sense and antisense strands.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 21 nucleotides long.

In one embodiment, a siNA molecule of the invention comprises no ribonucleotides. In another embodiment, a siNA molecule of the invention comprises ribonucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene, and wherein the second strand of the double-stranded siNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein each strand of the siNA molecule comprises about 19 to about 23 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene, and wherein the siNA further comprises a sense region, wherein the sense region comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the antisense region and the sense region each comprise about 19 to about 23 nucleotides, and wherein the antisense region comprises at least about 19 nucleotides that
5 are complementary to nucleotides of the sense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide
10 sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments
15 wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule. The sense region can be connected to the antisense region via a linker molecule, such as a polynucleotide linker or a non-nucleotide linker.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene,
20 wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region, and wherein pyrimidine nucleotides in the sense region are 2'-O-methyl pyrimidine
25 nucleotides, 2'-deoxy purine nucleotides, or 2'-deoxy-2'-fluoro pyrimidine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments

wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule, and wherein the fragment comprising the sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the fragment comprising the sense region. In another embodiment, the terminal cap moiety is an inverted deoxy abasic moiety or glyceryl moiety. In another embodiment, each of the two fragments of the siNA molecule comprise about 21 nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region, and wherein the purine nucleotides present in the antisense region comprise 2'-deoxy- purine nucleotides. In another embodiment, the antisense region comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense region. In another embodiment, the antisense region comprises a glyceryl modification at the 3' end of the antisense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule, and wherein about 19 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule and wherein at least two 3' terminal nucleotides of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule. In another embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines, such as 2'-deoxy-thymidine. In another embodiment, all 21 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule. In another embodiment, about 19 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by the VEGF and/or VEGFr

gene. In another embodiment, 21 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by the VEGF and/or VEGFr gene. In another embodiment, the 5'-end of the fragment comprising said antisense region optionally includes a phosphate group.

5 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGF and/or VEGFr RNA sequence (e.g., wherein said target RNA sequence is encoded by a VEGF and/or VEGFr gene), wherein the siNA molecule comprises no ribonucleotides and wherein each strand of the double-stranded siNA molecule is about 21 nucleotides long.

10 In one embodiment, the invention features a medicament comprising a siNA molecule of the invention.

 In one embodiment, the invention features an active ingredient comprising a siNA molecule of the invention.

15 In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 21 nucleotides long.

 In one embodiment, a VEGFr gene contemplated by the invention is a VEGFr1, VEGFr2, or VEGFr3 gene.

20 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide
25 sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule

comprises a sugar modification. In one embodiment, the VEGFr gene is VEGFr2. In one embodiment, the VEGFr gene is VEGFr1.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one
5 of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein
10 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the nucleotide sequence of the antisense strand of the double-stranded siNA molecule is complementary to the nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof which encodes an protein or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic
15 acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein
20 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein each strand of the siNA molecule comprises about 19 to about 29 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.

In one embodiment, the invention features a double-stranded short interfering nucleic
25 acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein

a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the siNA molecule is assembled from two oligonucleotide fragments wherein one fragment comprises the nucleotide sequence of the antisense strand of the siNA molecule and a second fragment comprises nucleotide sequence of the sense region of the siNA molecule.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the sense strand is connected to the antisense strand via a linker molecule, such as a polynucleotide linker or a non-nucleotide linker.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein pyrimidine nucleotides present in the sense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein purine nucleotides present in the sense region are 2'-deoxy purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide

sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the sense strand comprises a 3'-end and a 5'-end, and wherein a terminal cap moiety (e.g., an inverted deoxy abasic moiety) is present at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the sense strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides and one or more 2'-O-methyl purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the pyrimidine nucleotides present in the antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein any purine nucleotides present in the antisense strand are 2'-O-methyl purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide

sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense strand.

5 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide
10 sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises a glyceryl modification at the 3' end.

In one embodiment, the invention features a double-stranded short interfering nucleic
15 acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide
20 sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein each of the two strands of the siNA molecule comprises 21 nucleotides. In another embodiment, about 19 nucleotides of each strand of the siNA molecule are base-paired to the complementary nucleotides of the other strand of the siNA molecule and wherein at least two 3' terminal nucleotides of each strand of the
25 siNA molecule are not base-paired to the nucleotides of the other strand of the siNA molecule. In another embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines, such as 2'-deoxy-thymidine. In another embodiment, each strand of the siNA molecule are base-paired to the complementary nucleotides of the other strand of the siNA molecule. In another embodiment, about 19
30 nucleotides of the antisense strand are base-paired to the nucleotide sequence of the VEGF

and/or VEGFr RNA or a portion thereof. In another embodiment, 21 nucleotides of the antisense strand are base-paired to the nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof.

5 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein
10 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the 5'-end of the antisense strand optionally includes a phosphate group.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one
15 of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule
20 comprises a sugar modification, and wherein the nucleotide sequence or a portion thereof of the antisense strand is complementary to a nucleotide sequence of the 5'-untranslated region or a portion thereof of the VEGF and/or VEGFr RNA.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one
25 of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule

comprises a sugar modification, and wherein the nucleotide sequence or a portion thereof of the antisense strand is complementary to a nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof that is present in the VEGF and/or VEGFr RNA.

5 In one embodiment, the invention features a pharmaceutical composition comprising a siNA molecule of the invention in an acceptable carrier or diluent.

In one embodiment, the invention features a medicament comprising an siNA molecule of the invention.

In one embodiment, the invention features an active ingredient comprising an siNA molecule of the invention.

10 In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which
15 comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification.


In a non-limiting example, the introduction of chemically-modified nucleotides into nucleic acid molecules provides a powerful tool in overcoming potential limitations of *in vivo* stability and bioavailability inherent to native RNA molecules that are delivered
20 exogenously. For example, the use of chemically-modified nucleic acid molecules can enable a lower dose of a particular nucleic acid molecule for a given therapeutic effect since chemically-modified nucleic acid molecules tend to have a longer half-life in serum. Furthermore, certain chemical modifications can improve the bioavailability of nucleic acid
25 molecules by targeting particular cells or tissues and/or improving cellular uptake of the nucleic acid molecule. Therefore, even if the activity of a chemically-modified nucleic acid molecule is reduced as compared to a native nucleic acid molecule, for example, when compared to an all-RNA nucleic acid molecule, the overall activity of the modified nucleic

acid molecule can be greater than that of the native molecule due to improved stability and/or delivery of the molecule. Unlike native unmodified siNA, chemically-modified siNA can also minimize the possibility of activating interferon activity in humans.

5 The antisense region of a siNA molecule of the invention can comprise a phosphorothioate internucleotide linkage at the 3'-end of said antisense region. The antisense region can comprise about one to about five phosphorothioate internucleotide linkages at the 5'-end of said antisense region. The 3'-terminal nucleotide overhangs of a siNA molecule of the invention can comprise ribonucleotides or deoxyribonucleotides that are chemically-modified at a nucleic acid sugar, base, or backbone. The 3'-terminal
10 nucleotide overhangs can comprise one or more universal base ribonucleotides. The 3'-terminal nucleotide overhangs can comprise one or more acyclic nucleotides.

One embodiment of the invention provides an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the invention in a manner that allows expression of the nucleic acid molecule. Another embodiment of the invention provides a
15 mammalian cell comprising such an expression vector. The mammalian cell can be a human cell. The siNA molecule of the expression vector can comprise a sense region and an antisense region. The antisense region can comprise sequence complementary to a RNA or DNA sequence encoding VEGF and/or VEGFr and the sense region can comprise sequence complementary to the antisense region. The siNA molecule can comprise two distinct
20 strands having complementary sense and antisense regions. The siNA molecule can comprise a single strand having complementary sense and antisense regions.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical
25 modification comprises one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides comprising a backbone modified internucleotide linkage having Formula I:




wherein each R1 and R2 is independently any nucleotide, non-nucleotide, or polynucleotide which can be naturally-occurring or chemically-modified, each X and Y is independently O, S, N, alkyl, or substituted alkyl, each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl, and wherein W, X, Y, and Z are optionally not all O.

The chemically-modified internucleotide linkages having Formula I, for example, wherein any Z, W, X, and/or Y independently comprises a sulphur atom, can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) chemically-modified internucleotide linkages having Formula I at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified internucleotide linkages having Formula I at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In another embodiment, a siNA molecule of the invention having internucleotide linkage(s) of Formula I also comprises a chemically-modified nucleotide or non-nucleotide having any of Formulae I-VII.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a

VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula II:




- 5 wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2, and B is a nucleosidic base
- 10 such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally occurring base that can be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring universal base that can be complementary or non-complementary to target RNA.
- 15

The chemically-modified nucleotide or non-nucleotide of Formula II can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more chemically-modified nucleotide or non-nucleotide of Formula II at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 5'-end of the sense strand, the antisense strand, or both

20

strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 3'-end of the sense strand, the antisense strand, or both strands.

- 5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula III:




10

- wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF₃, OCF₃, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO₂, NO₂, N₃, NH₂, aminoalkyl, aminoacid, aminoacyl, ONH₂, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH₂, S=O, CHF, or CF₂, and B is a nucleosidic base such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally occurring base that can be employed to be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring universal base that can be complementary or non-complementary to target RNA.
- 15
- 20

The chemically-modified nucleotide or non-nucleotide of Formula III can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more chemically-modified nucleotide or non-nucleotide of Formula III at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide(s) or non-nucleotide(s) of Formula III at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide or non-nucleotide of Formula III at the 3'-end of the sense strand, the antisense strand, or both strands.

In another embodiment, a siNA molecule of the invention comprises a nucleotide having Formula II or III, wherein the nucleotide having Formula II or III is in an inverted configuration. For example, the nucleotide having Formula II or III is connected to the siNA construct in a 3'-3', 3'-2', 2'-3', or 5'-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises a 5'-terminal phosphate group having Formula IV:



wherein each X and Y is independently O, S, N, alkyl, substituted alkyl, or alkylhalo; wherein each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, aralkyl, or alkylhalo; and wherein W, X, Y and Z are not all O.

In one embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand, for example, a strand complementary to a target RNA, wherein the siNA molecule comprises an all RNA siNA molecule. In another embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand wherein the siNA molecule also comprises about 1 to about 3 (e.g., about 1, 2, or 3) nucleotide 3'-terminal nucleotide overhangs having about 1 to about 4 (e.g., about 1, 2, 3, or 4) deoxyribonucleotides on the 3'-end of one or both strands. In another embodiment, a 5'-terminal phosphate group having Formula IV is present on the target-complementary strand of a siNA molecule of the invention, for example a siNA molecule having chemical modifications having any of Formulae I-VII.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more phosphorothioate internucleotide linkages. For example, in a non-limiting example, the invention features a chemically-modified short interfering nucleic acid (siNA) having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate internucleotide linkages in one siNA strand. In yet another embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) individually having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate internucleotide linkages in both siNA strands. The phosphorothioate internucleotide linkages can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more phosphorothioate internucleotide linkages at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) consecutive phosphorothioate internucleotide linkages at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine phosphorothioate internucleotide linkages in the sense

strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine phosphorothioate internucleotide linkages in the sense strand, the antisense strand, or both strands.

5 In one embodiment, the invention features a siNA molecule, wherein the sense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally
10 a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base
15 modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example about 1, 2, 3, 4, 5, 6,
20 7, 8, 9, 10, or more, phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the sense strand comprises about 1 to about 5, specifically about 1, 2, 3, 4, or 5 phosphorothioate
25 internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5, or more
30 phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7,

8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5 or more, for example about 1, 2, 3, 4, 5, or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

10 In one embodiment, the invention features a siNA molecule, wherein the antisense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally
15 a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base
20 modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9,
25 10 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3' and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7,
30 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about

1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages, and/or one or more
5 (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense
10 siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5, for example about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

In one embodiment, the invention features a chemically-modified short interfering
15 nucleic acid (siNA) molecule having about 1 to about 5, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages in each strand of the siNA molecule.

In another embodiment, the invention features a siNA molecule comprising 2'-5' internucleotide linkages. The 2'-5' internucleotide linkage(s) can be at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both siNA sequence strands. In addition, the 2'-5' internucleotide linkage(s) can be present at various other positions within one or both siNA
20 sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the siNA molecule can comprise a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of
25 the siNA molecule can comprise a 2'-5' internucleotide linkage.

In another embodiment, a chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified, wherein each strand is about 18 to about 27 (e.g., about 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27) nucleotides in length, wherein the duplex has about 18 to about 23 (e.g., about 18, 19,


20, 21, 22, or 23) base pairs, and wherein the chemical modification comprises a structure having any of Formulae I-VII. For example, an exemplary chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein each strand consists of about 21 nucleotides, each having a 2-nucleotide 3'-terminal nucleotide overhang, and wherein the duplex has about 19 base pairs. In another embodiment, a siNA molecule of the invention comprises a single stranded hairpin structure, wherein the siNA is about 36 to about 70 (*e.g.*, about 36, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (*e.g.*, about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a linear oligonucleotide having about 42 to about 50 (*e.g.*, about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein the linear oligonucleotide forms a hairpin structure having about 19 base pairs and a 2-nucleotide 3'-terminal nucleotide overhang. In another embodiment, a linear hairpin siNA molecule of the invention contains a stem loop motif, wherein the loop portion of the siNA molecule is biodegradable. For example, a linear hairpin siNA molecule of the invention is designed such that degradation of the loop portion of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.

In another embodiment, a siNA molecule of the invention comprises a circular nucleic acid molecule, wherein the siNA is about 38 to about 70 (*e.g.*, about 38, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (*e.g.*, about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification, which comprises a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a circular oligonucleotide having about 42 to about 50 (*e.g.*, about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any

of Formulae I-VII or any combination thereof, wherein the circular oligonucleotide forms a dumbbell shaped structure having about 19 base pairs and 2 loops.


In another embodiment, a circular siNA molecule of the invention contains two loop motifs, wherein one or both loop portions of the siNA molecule is biodegradable. For example, a circular siNA molecule of the invention is designed such that degradation of the loop portions of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.

In one embodiment, a siNA molecule of the invention comprises at least one (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) abasic moiety, for example a compound having Formula V:




wherein each R3, R4, R5, R6, R7, R8, R10, R11, R12, and R13 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2.

In one embodiment, a siNA molecule of the invention comprises at least one (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) inverted abasic moiety, for example a compound having Formula VI:



wherein each R3, R4, R5, R6, R7, R8, R10, R11, R12, and R13 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2, and either R2, R3, R8 or R13 serve as points of attachment to the siNA molecule of the invention.

10 In another embodiment, a siNA molecule of the invention comprises at least one (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) substituted polyalkyl moieties, for example a compound having Formula VII:



wherein each n is independently an integer from 1 to 12, each R1, R2 and R3 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or a group having Formula I, and R1, R2 or R3 serves as points of attachment to the siNA molecule of the invention.

In another embodiment, the invention features a compound having Formula VII, wherein R1 and R2 are hydroxyl (OH) groups, $n = 1$, and R3 comprises O and is the point of attachment to the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both strands of a double-stranded siNA molecule of the invention or to a single-stranded siNA molecule of the invention. This modification is referred to herein as "glyceryl" (for example modification 6 in Figure 10).

In another embodiment, a moiety having any of Formula V, VI or VII of the invention is at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of a siNA molecule of the invention. For example, a moiety having Formula V, VI or VII can be present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense strand, the sense strand, or both antisense and sense strands of the siNA molecule. In addition, a moiety having Formula VII can be present at the 3'-end or the 5'-end of a hairpin siNA molecule as described herein.

In another embodiment, a siNA molecule of the invention comprises an abasic residue having Formula V or VI, wherein the abasic residue having Formula VI or VI is connected to the siNA construct in a 3'-3', 3'-2', 2'-3', or 5'-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) locked nucleic acid (LNA) nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In another embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) acyclic nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a

plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all
10 pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in
15 said sense region are 2'-deoxy nucleotides.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein
20 all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine
25 nucleotides).

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein

all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in said antisense region are 2'-deoxy nucleotides.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (*e.g.*, about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid

molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 4 and 5** and **Tables III and IV** herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are purine ribonucleotides (e.g., wherein all purine nucleotides are purine ribonucleotides or alternately a plurality of purine nucleotides are purine ribonucleotides), and inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the siNA comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of

pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in **Figures 4 and 5** and **Tables III and IV** herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and for example where one or more purine nucleotides present in the sense region are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and wherein inverted deoxy abasic modifications are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid

molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and
5 wherein one or more purine nucleotides present in the antisense region are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the
10 group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense
15 sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages.

In another embodiment, any modified nucleotides present in the siNA molecules of the
20 invention, preferably in the antisense strand of the siNA molecules of the invention, but also optionally in the sense and/or both antisense and sense strands, comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for
25 example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the siNA molecules of the invention, preferably in the antisense strand of the siNA molecules of the invention, but also optionally in the sense and/or both antisense and sense strands, are resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi. Non-limiting examples
30 of nucleotides having a northern configuration include locked nucleic acid (LNA)

nucleotides (e.g., 2'-O, 4'-C-methylene-(D-ribofuranosyl) nucleotides); 2'-methoxyethoxy (MOE) nucleotides; 2'-methyl-thio-ethyl, 2'-deoxy-2'-fluoro nucleotides, 2'-deoxy-2'-chloro nucleotides, 2'-azido nucleotides, and 2'-O-methyl nucleotides.

In one embodiment, the invention features a chemically-modified short interfering
5 nucleic acid molecule (siNA) capable of mediating RNA interference (RNAi) against a
VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical
modification comprises a conjugate covalently attached to the chemically-modified siNA
molecule. In another embodiment, the conjugate is covalently attached to the chemically-
modified siNA molecule via a biodegradable linker. In one embodiment, the conjugate
10 molecule is attached at the 3'-end of either the sense strand, the antisense strand, or both
strands of the chemically-modified siNA molecule. In another embodiment, the conjugate
molecule is attached at the 5'-end of either the sense strand, the antisense strand, or both
strands of the chemically-modified siNA molecule. In yet another embodiment, the
conjugate molecule is attached both the 3'-end and 5'-end of either the sense strand, the
15 antisense strand, or both strands of the chemically-modified siNA molecule, or any
combination thereof. In one embodiment, a conjugate molecule of the invention comprises a
molecule that facilitates delivery of a chemically-modified siNA molecule into a biological
system, such as a cell. In another embodiment, the conjugate molecule attached to the
chemically-modified siNA molecule is a poly ethylene glycol, human serum albumin, or a
20 ligand for a cellular receptor that can mediate cellular uptake. Examples of specific
conjugate molecules contemplated by the instant invention that can be attached to
chemically-modified siNA molecules are described in Vargeese *et al.*, U.S. Serial No.
10/201,394, incorporated by reference herein. The type of conjugates used and the extent of
conjugation of siNA molecules of the invention can be evaluated for improved
25 pharmacokinetic profiles, bioavailability, and/or stability of siNA constructs while at the
same time maintaining the ability of the siNA to mediate RNAi activity. As such, one
skilled in the art can screen siNA constructs that are modified with various conjugates to
determine whether the siNA conjugate complex possesses improved properties while
maintaining the ability to mediate RNAi, for example in animal models as are generally
30 known in the art.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule of the invention, wherein the siNA further comprises a nucleotide, non-nucleotide, or mixed nucleotide/non-nucleotide linker that joins the sense region of the siNA to the antisense region of the siNA. In one embodiment, a nucleotide linker of the invention can be a linker of ≥ 2 nucleotides in length, for example 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides in length. In another embodiment, the nucleotide linker can be a nucleic acid aptamer. By “aptamer” or “nucleic acid aptamer” as used herein is meant a nucleic acid molecule that binds specifically to a target molecule wherein the nucleic acid molecule has sequence that comprises a sequence recognized by the target molecule in its natural setting. Alternately, an aptamer can be a nucleic acid molecule that binds to a target molecule where the target molecule does not naturally bind to a nucleic acid. The target molecule can be any molecule of interest. For example, the aptamer can be used to bind to a ligand-binding domain of a protein, thereby preventing interaction of the naturally occurring ligand with the protein. This is a non-limiting example and those in the art will recognize that other embodiments can be readily generated using techniques generally known in the art. (See, for example, Gold *et al.*, 1995, *Annu. Rev. Biochem.*, 64, 763; Brody and Gold, 2000, *J. Biotechnol.*, 74, 5; Sun, 2000, *Curr. Opin. Mol. Ther.*, 2, 100; Kusser, 2000, *J. Biotechnol.*, 74, 27; Hermann and Patel, 2000, *Science*, 287, 820; and Jayasena, 1999, *Clinical Chemistry*, 45, 1628.)

In yet another embodiment, a non-nucleotide linker of the invention comprises abasic nucleotide, polyether, polyamine, polyamide, peptide, carbohydrate, lipid, polyhydrocarbon, or other polymeric compounds (e.g. polyethylene glycols such as those having between 2 and 100 ethylene glycol units). Specific examples include those described by Seela and Kaiser, *Nucleic Acids Res.* 1990, 18:6353 and *Nucleic Acids Res.* 1987, 15:3113; Cload and Schepartz, *J. Am. Chem. Soc.* 1991, 113:6324; Richardson and Schepartz, *J. Am. Chem. Soc.* 1991, 113:5109; Ma *et al.*, *Nucleic Acids Res.* 1993, 21:2585 and *Biochemistry* 1993, 32:1751; Durand *et al.*, *Nucleic Acids Res.* 1990, 18:6353; McCurdy *et al.*, *Nucleosides & Nucleotides* 1991, 10:287; Jsche *et al.*, *Tetrahedron Lett.* 1993, 34:301; Ono *et al.*, *Biochemistry* 1991, 30:9914; Arnold *et al.*, International Publication No. WO 89/02439; Usman *et al.*, International Publication No. WO 95/06731; Dudycz *et al.*, International Publication No. WO 95/11910 and Ferentz and Verdine, *J. Am. Chem. Soc.* 1991, 113:4000,

all hereby incorporated by reference herein. A "non-nucleotide" further means any group or compound that can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound can be abasic
5 in that it does not contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine, for example at the C1 position of the sugar.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted in vitro system, wherein one or both strands of the siNA molecule that are assembled from two
10 separate oligonucleotides do not comprise any ribonucleotides. For example, a siNA molecule can be assembled from a single oligonucleotide where the sense and antisense regions of the siNA comprise separate oligonucleotides not having any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotides. In another example, a siNA molecule can be assembled from a single oligonucleotide where the sense and
15 antisense regions of the siNA are linked or circularized by a nucleotide or non-nucleotide linker as described herein, wherein the oligonucleotide does not have any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotide. Applicant has surprisingly found that the presence of ribonucleotides (e.g., nucleotides having a 2'-hydroxyl group) within the siNA molecule is not required or essential to support RNAi
20 activity. As such, in one embodiment, all positions within the siNA can include chemically modified nucleotides and/or non-nucleotides such as nucleotides and or non-nucleotides having Formula I, II, III, IV, V, VI, or VII or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

In one embodiment, a siNA molecule of the invention is a single stranded siNA
25 molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group and a 3'-
30 terminal phosphate group (e.g., a 2',3'-cyclic phosphate). In another embodiment, the single

stranded siNA molecule of the invention comprises about 19 to about 29 nucleotides. In yet another embodiment, the single stranded siNA molecule of the invention comprises one or more chemically modified nucleotides or non-nucleotides described herein. For example, all the positions within the siNA molecule can include chemically-modified nucleotides such as nucleotides having any of Formulae I-VII, or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine

nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and a terminal cap modification, such as any
5 modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the
10 siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a
15 target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are locked nucleic acid (LNA) nucleotides (e.g.,
20 wherein all purine nucleotides are LNA nucleotides or alternately a plurality of purine nucleotides are LNA nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the
25 siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the
30 siNA molecule comprises a single stranded polynucleotide having complementarity to a

target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine
5 nucleotides present in the antisense region are 2'-methoxyethyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-methoxyethyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-methoxyethyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in **Figure 10**, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense
10 sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

15 In another embodiment, any modified nucleotides present in the single stranded siNA molecules of the invention comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid*
20 *Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the single stranded siNA molecules of the invention are preferably resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi.

In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of
25 the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and
5 wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

In another embodiment, the invention features a method for modulating the expression of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing
10 siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr genes; and (b) introducing the siNA molecules into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In another embodiment, the invention features a method for modulating the expression
15 of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecules into a cell under
20 conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In one embodiment, siNA molecules of the invention are used as reagents in ex vivo applications. For example, siNA reagents are introduced into tissue or cells that are transplanted into a subject for therapeutic effect. The cells and/or tissue can be derived from an organism or subject that later receives the explant, or can be derived from another
25 organism or subject prior to transplantation. The siNA molecules can be used to modulate the expression of one or more genes in the cells or tissue, such that the cells or tissue obtain a desired phenotype or are able to perform a function when transplanted in vivo. In one embodiment, certain target cells from a patient are extracted. These extracted cells are contacted with siNAs targeting a specific nucleotide sequence within the cells under

conditions suitable for uptake of the siNAs by these cells (e.g. using delivery reagents such as cationic lipids, liposomes and the like or using techniques such as electroporation to facilitate the delivery of siNAs into cells). The cells are then reintroduced back into the same patient or other patients. In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in that organism.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr genes; and (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF

and/or VEGFr genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in that organism.

5 In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into the organism under conditions suitable to modulate the
10 expression of the VEGF and/or VEGFr gene in the organism.

 In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr
15 genes; and (b) introducing the siNA molecules into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

 In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single
20 stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

 In another embodiment, the invention features a method for modulating the expression of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing
25 siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) contacting the siNA molecule with a cell in vitro or in vivo under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and
5 (b) contacting the siNA molecule with a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr
10 gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or
15 VEGFr gene; and (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression
20 of the VEGF and/or VEGFr genes in that organism.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and
25 (b) introducing the siNA molecule into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA

comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecules into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

5 In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising contacting the organism with a siNA molecule of the invention under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the organism.

10 In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising contacting the organism with one or more siNA molecules of the invention under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

The siNA molecules of the invention can be designed to inhibit target (VEGF and/or VEGFr) gene expression through RNAi targeting of a variety of RNA molecules. In one embodiment, the siNA molecules of the invention are used to target various RNAs
15 corresponding to a target gene. Non-limiting examples of such RNAs include messenger RNA (mRNA), alternate RNA splice variants of target gene(s), post-transcriptionally modified RNA of target gene(s), pre-mRNA of target gene(s), and/or RNA templates. If alternate splicing produces a family of transcripts that are distinguished by usage of appropriate exons, the instant invention can be used to inhibit gene expression through the
20 appropriate exons to specifically inhibit or to distinguish among the functions of gene family members. For example, a protein that contains an alternatively spliced transmembrane domain can be expressed in both membrane bound and secreted forms. Use of the invention to target the exon containing the transmembrane domain can be used to determine the functional consequences of pharmaceutical targeting of membrane bound as opposed to the
25 secreted form of the protein. Non-limiting examples of applications of the invention relating to targeting these RNA molecules include therapeutic pharmaceutical applications, pharmaceutical discovery applications, molecular diagnostic and gene function applications, and gene mapping, for example using single nucleotide polymorphism mapping with siNA

molecules of the invention. Such applications can be implemented using known gene sequences or from partial sequences available from an expressed sequence tag (EST).

In another embodiment, the siNA molecules of the invention are used to target conserved sequences corresponding to a gene family or gene families such as VEGF and/or VEGFr family genes. As such, siNA molecules targeting multiple VEGF and/or VEGFr targets can provide increased therapeutic effect. In addition, siNA can be used to characterize pathways of gene function in a variety of applications. For example, the present invention can be used to inhibit the activity of target gene(s) in a pathway to determine the function of uncharacterized gene(s) in gene function analysis, mRNA function analysis, or translational analysis. The invention can be used to determine potential target gene pathways involved in various diseases and conditions toward pharmaceutical development. The invention can be used to understand pathways of gene expression involved in, for example, the progression and/or maintenance of cancer.

In one embodiment, siNA molecule(s) and/or methods of the invention are used to inhibit the expression of gene(s) that encode RNA referred to by Genbank Accession, for example VEGF and/or VEGFr genes encoding RNA sequence(s) referred to herein by Genbank Accession number, for example, Genbank Accession Nos. shown in Table I.

In one embodiment, the invention features a method comprising: (a) generating a library of siNA constructs having a predetermined complexity; and (b) assaying the siNA constructs of (a) above, under conditions suitable to determine RNAi target sites within the target RNA sequence. In another embodiment, the siNA molecules of (a) have strands of a fixed length, for example, about 23 nucleotides in length. In yet another embodiment, the siNA molecules of (a) are of differing length, for example having strands of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. In another embodiment, fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA

sequence. The target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In one embodiment, the invention features a method comprising: (a) generating a
5 randomized library of siNA constructs having a predetermined complexity, such as of 4^N ,
where N represents the number of base paired nucleotides in each of the siNA construct
strands (eg. for a siNA construct having 21 nucleotide sense and antisense strands with 19
base pairs, the complexity would be 4^{19}); and (b) assaying the siNA constructs of (a) above,
under conditions suitable to determine RNAi target sites within the target VEGF and/or
10 VEGFr RNA sequence. In another embodiment, the siNA molecules of (a) have strands of a
fixed length, for example about 23 nucleotides in length. In yet another embodiment, the
siNA molecules of (a) are of differing length, for example having strands of about 19 to
about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one
embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described in
15 Example 7 herein. In another embodiment, the assay can comprise a cell culture system in
which target RNA is expressed. In another embodiment, fragments of VEGF and/or VEGFr
RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis,
northern blot analysis, or RNase protection assays, to determine the most suitable target
site(s) within the target VEGF and/or VEGFr RNA sequence. The target VEGF and/or
20 VEGFr RNA sequence can be obtained as is known in the art, for example, by cloning
and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In another embodiment, the invention features a method comprising: (a) analyzing the
sequence of a RNA target encoded by a target gene; (b) synthesizing one or more sets of
siNA molecules having sequence complementary to one or more regions of the RNA of (a);
25 and (c) assaying the siNA molecules of (b) under conditions suitable to determine RNAi
targets within the target RNA sequence. In one embodiment, the siNA molecules of (b)
have strands of a fixed length, for example about 23 nucleotides in length. In another
embodiment, the siNA molecules of (b) are of differing length, for example having strands
of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In
30 one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described

herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. Fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA sequence. The target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by expression in *in vivo* systems.

By "target site" is meant a sequence within a target RNA that is "targeted" for cleavage mediated by a siNA construct which contains sequences within its antisense region that are complementary to the target sequence.

By "detectable level of cleavage" is meant cleavage of target RNA (and formation of cleaved product RNAs) to an extent sufficient to discern cleavage products above the background of RNAs produced by random degradation of the target RNA. Production of cleavage products from 1-5% of the target RNA is sufficient to detect above the background for most methods of detection.

In one embodiment, the invention features a composition comprising a siNA molecule of the invention, which can be chemically-modified, in a pharmaceutically acceptable carrier or diluent. In another embodiment, the invention features a pharmaceutical composition comprising siNA molecules of the invention, which can be chemically-modified, targeting one or more genes in a pharmaceutically acceptable carrier or diluent. In another embodiment, the invention features a method for treating or preventing a disease or condition in a subject, comprising administering to the subject a composition of the invention under conditions suitable for the treatment or prevention of the disease or condition in the subject, alone or in conjunction with one or more other therapeutic compounds. In yet another embodiment, the invention features a method for reducing or preventing tissue rejection in a subject comprising administering to the subject a composition of the invention under conditions suitable for the reduction or prevention of tissue rejection in the subject.

In another embodiment, the invention features a method for validating a VEGF and/or VEGFr gene target, comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a VEGF and/or VEGFr target gene; (b) introducing the siNA molecule into a cell, tissue, or organism under conditions suitable for modulating expression of the VEGF and/or VEGFr target gene in the cell, tissue, or organism; and (c) determining the function of the gene by assaying for any phenotypic change in the cell, tissue, or organism.

In another embodiment, the invention features a method for validating a VEGF and/or VEGFr target comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a VEGF and/or VEGFr target gene; (b) introducing the siNA molecule into a biological system under conditions suitable for modulating expression of the VEGF and/or VEGFr target gene in the biological system; and (c) determining the function of the gene by assaying for any phenotypic change in the biological system.

By "biological system" is meant, material, in a purified or unpurified form, from biological sources, including but not limited to human, animal, plant, insect, bacterial, viral or other sources, wherein the system comprises the components required for RNAi activity. The term "biological system" includes, for example, a cell, tissue, or organism, or extract thereof. The term biological system also includes reconstituted RNAi systems that can be used in an *in vitro* setting.

By "phenotypic change" is meant any detectable change to a cell that occurs in response to contact or treatment with a nucleic acid molecule of the invention (e.g., siNA). Such detectable changes include, but are not limited to, changes in shape, size, proliferation, motility, protein expression or RNA expression or other physical or chemical changes as can be assayed by methods known in the art. The detectable change can also include expression of reporter genes/molecules such as Green Florescent Protein (GFP) or various tags that are used to identify an expressed protein or any other cellular component that can be assayed.

In one embodiment, the invention features a kit containing a siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of a VEGF and/or VEGFr target gene in a cell, tissue, or organism. In another embodiment, the invention features a kit containing more than one siNA molecule of the invention, which
5 can be chemically-modified, that can be used to modulate the expression of more than one VEGF and/or VEGFr target gene in a cell, tissue, or organism.

In one embodiment, the invention features a cell containing one or more siNA molecules of the invention, which can be chemically-modified. In another embodiment, the cell containing a siNA molecule of the invention is a mammalian cell. In yet another
10 embodiment, the cell containing a siNA molecule of the invention is a human cell.

In one embodiment, the synthesis of a siNA molecule of the invention, which can be chemically-modified, comprises: (a) synthesis of two complementary strands of the siNA molecule; (b) annealing the two complementary strands together under conditions suitable to obtain a double-stranded siNA molecule. In another embodiment, synthesis of the two
15 complementary strands of the siNA molecule is by solid phase oligonucleotide synthesis. In yet another embodiment, synthesis of the two complementary strands of the siNA molecule is by solid phase tandem oligonucleotide synthesis.

In one embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing a first oligonucleotide sequence strand of the siNA
20 molecule, wherein the first oligonucleotide sequence strand comprises a cleavable linker molecule that can be used as a scaffold for the synthesis of the second oligonucleotide sequence strand of the siNA; (b) synthesizing the second oligonucleotide sequence strand of siNA on the scaffold of the first oligonucleotide sequence strand, wherein the second oligonucleotide sequence strand further comprises a chemical moiety than can be used to
25 purify the siNA duplex; (c) cleaving the linker molecule of (a) under conditions suitable for the two siNA oligonucleotide strands to hybridize and form a stable duplex; and (d) purifying the siNA duplex utilizing the chemical moiety of the second oligonucleotide sequence strand. In one embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example under hydrolysis conditions

using an alkylamine base such as methylamine. In one embodiment, the method of synthesis comprises solid phase synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized linker and the cleavable linker of (a) takes place concomitantly. In another embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group, which can be employed in a trityl-on synthesis strategy as described herein. In yet another embodiment, the chemical moiety, such as a dimethoxytrityl group, is removed during purification, for example, using acidic conditions.

In a further embodiment, the method for siNA synthesis is a solution phase synthesis or hybrid phase synthesis wherein both strands of the siNA duplex are synthesized in tandem using a cleavable linker attached to the first sequence which acts a scaffold for synthesis of the second sequence. Cleavage of the linker under conditions suitable for hybridization of the separate siNA sequence strands results in formation of the double-stranded siNA molecule.

In another embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing one oligonucleotide sequence strand of the siNA molecule, wherein the sequence comprises a cleavable linker molecule that can be used as a scaffold for the synthesis of another oligonucleotide sequence; (b) synthesizing a second oligonucleotide sequence having complementarity to the first sequence strand on the scaffold of (a), wherein the second sequence comprises the other strand of the double-stranded siNA molecule and wherein the second sequence further comprises a chemical moiety than can be used to isolate the attached oligonucleotide sequence; (c) purifying the product of (b) utilizing the chemical moiety of the second oligonucleotide sequence strand under conditions suitable for isolating the full-length sequence comprising both siNA oligonucleotide strands connected by the cleavable linker and under conditions suitable for the two siNA oligonucleotide strands to hybridize and form a stable duplex. In one

embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example under hydrolysis conditions. In another embodiment, cleavage of the linker molecule in (c) above takes place after deprotection of the oligonucleotide. In another embodiment, the method of synthesis comprises solid phase
5 synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity or differing reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized linker and the
10 cleavable linker of (a) takes place either concomitantly or sequentially. In one embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group.

In another embodiment, the invention features a method for making a double-stranded siNA molecule in a single synthetic process comprising: (a) synthesizing an oligonucleotide
15 having a first and a second sequence, wherein the first sequence is complementary to the second sequence, and the first oligonucleotide sequence is linked to the second sequence via a cleavable linker, and wherein a terminal 5'-protecting group, for example, a 5'-O-dimethoxytrityl group (5'-O-DMT) remains on the oligonucleotide having the second sequence; (b) deprotecting the oligonucleotide whereby the deprotection results in the
20 cleavage of the linker joining the two oligonucleotide sequences; and (c) purifying the product of (b) under conditions suitable for isolating the double-stranded siNA molecule, for example using a trityl-on synthesis strategy as described herein.

In another embodiment, the method of synthesis of siNA molecules of the invention comprises the teachings of Scaringe *et al.*, US Patent Nos. 5,889,136; 6,008,400; and
25 6,111,086, incorporated by reference herein in their entirety.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications, for example, one or more chemical modifications having any of Formulae I-VII or any combination thereof that increases the nuclease resistance of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased nuclease resistance comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased nuclease resistance.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the sense and antisense strands of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the sense and antisense strands of the siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the sense and antisense strands of the siNA molecule.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target RNA sequence within a cell.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target DNA sequence within a cell.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having

increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence.

5 In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target DNA sequence comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the antisense strand of the siNA molecule and a complementary target DNA sequence.

10 In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulate the polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA construct.

15 In another embodiment, the invention features a method for generating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to a chemically-modified siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA molecule.

20 In one embodiment, the invention features chemically-modified siNA constructs that mediate RNAi against a VEGF and/or VEGFr in a cell, wherein the chemical modifications do not significantly effect the interaction of siNA with a target RNA molecule, DNA molecule and/or proteins or other factors that are essential for RNAi in a manner that would decrease the efficacy of RNAi mediated by such siNA constructs.

In another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against VEGF and/or VEGFr comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a VEGF and/or VEGFr target RNA comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the target RNA.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a VEGF and/or VEGFr target DNA comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the target DNA.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the cellular uptake of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules against VEGF and/or VEGFr with improved cellular uptake comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved cellular uptake.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that increases the bioavailability of the siNA construct, for

example, by attaching polymeric conjugates such as polyethyleneglycol or equivalent conjugates that improve the pharmacokinetics of the siNA construct, or by attaching conjugates that target specific tissue types or cell types *in vivo*. Non-limiting examples of such conjugates are described in Vargeese *et al.*, U.S. Serial No. 10/201,394 incorporated by
5 reference herein.

In one embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability, comprising (a) introducing a conjugate into the structure of a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability. Such
10 conjugates can include ligands for cellular receptors, such as peptides derived from naturally occurring protein ligands; protein localization sequences, including cellular ZIP code sequences; antibodies; nucleic acid aptamers; vitamins and other co-factors, such as folate and N-acetylgalactosamine; polymers, such as polyethyleneglycol (PEG); phospholipids; polyamines, such as spermine or spermidine; and others.

15 In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing an excipient formulation to a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability. Such excipients include polymers such as cyclodextrins, lipids, cationic lipids, polyamines,
20 phospholipids, and others.

In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing nucleotides having any of Formulae I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA
25 molecules having improved bioavailability.

In another embodiment, polyethylene glycol (PEG) can be covalently attached to siNA compounds of the present invention. The attached PEG can be any molecular weight, preferably from about 2,000 to about 50,000 daltons (Da).

The present invention can be used alone or as a component of a kit having at least one of the reagents necessary to carry out the *in vitro* or *in vivo* introduction of RNA to test samples and/or subjects. For example, preferred components of the kit include a siNA molecule of the invention and a vehicle that promotes introduction of the siNA into cells of interest as described herein (e.g., using lipids and other methods of transfection known in the art, see for example Beigelman *et al.*, US 6,395,713). The kit can be used for target validation, such as in determining gene function and/or activity, or in drug optimization, and in drug discovery (see for example Usman *et al.*, USSN 60/402,996). Such a kit can also include instructions to allow a user of the kit to practice the invention.

The term "short interfering nucleic acid", "siNA", "short interfering RNA", "siRNA", "short interfering nucleic acid molecule", "short interfering oligonucleotide molecule", or "chemically-modified short interfering nucleic acid molecule" as used herein refers to any nucleic acid molecule capable of inhibiting or down regulating gene expression, for example by mediating RNA interference "RNAi" or gene silencing in a sequence-specific manner; see for example Bass, 2001, *Nature*, 411, 428-429; Elbashir *et al.*, 2001, *Nature*, 411, 494-498; and Kreutzer *et al.*, International PCT Publication No. WO 00/44895; Zernicka-Goetz *et al.*, International PCT Publication No. WO 01/36646; Fire, International PCT Publication No. WO 99/32619; Plaetinck *et al.*, International PCT Publication No. WO 00/01846; Mello and Fire, International PCT Publication No. WO 01/29058; Deschamps-Depaillette, International PCT Publication No. WO 99/07409; and Li *et al.*, International PCT Publication No. WO 00/44914; Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237; Hutvagner and Zamore, 2002, *Science*, 297, 2056-60; McManus *et al.*, 2002, *RNA*, 8, 842-850; Reinhart *et al.*, 2002, *Gene & Dev.*, 16, 1616-1626; and Reinhart & Bartel, 2002, *Science*, 297, 1831). Non limiting examples of siNA molecules of the invention are shown in **Figures 4-6, and Tables II, III, and IV** herein. For example the siNA can be a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic

acid sequence or a portion thereof. The siNA can be assembled from two separate oligonucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary (i.e. each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double stranded structure, for example wherein the double stranded region is about 19 base pairs); the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof.

Alternatively, the siNA is assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the siNA are linked by means of a nucleic acid based or non-nucleic acid-based linker(s). The siNA can be a polynucleotide with a hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The siNA can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either *in vivo* or *in vitro* to generate an active siNA molecule capable of mediating RNAi. The siNA can also comprise a single stranded polynucleotide having nucleotide sequence complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof (for example, where such siNA molecule does not require the presence within the siNA molecule of nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide can further comprise a terminal phosphate group, such as a 5'-phosphate (see for example Martinez *et al.*, 2002, *Cell.*, 110, 563-574 and Schwarz *et al.*, 2002, *Molecular Cell*, 10, 537-568), or 5',3'-diphosphate. In certain embodiment, the siNA molecule of the invention comprises separate sense and antisense

sequences or regions, wherein the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linkers molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. In certain embodiments, the siNA molecules of the invention comprise nucleotide sequence that is complementary to nucleotide sequence of a target gene. In another embodiment, the siNA molecule of the invention interacts with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene. As used herein, siNA molecules need not be limited to those molecules containing only RNA, but further encompasses chemically-modified nucleotides and non-nucleotides. In certain embodiments, the short interfering nucleic acid molecules of the invention lack 2'-hydroxy (2'-OH) containing nucleotides. Applicant describes in certain embodiments short interfering nucleic acids that do not require the presence of nucleotides having a 2'-hydroxy group for mediating RNAi and as such, short interfering nucleic acid molecules of the invention optionally do not include any ribonucleotides (e.g., nucleotides having a 2'-OH group). Such siNA molecules that do not require the presence of ribonucleotides within the siNA molecule to support RNAi can however have an attached linker or linkers or other attached or associated groups, moieties, or chains containing one or more nucleotides with 2'-OH groups. Optionally, siNA molecules can comprise ribonucleotides at about 5, 10, 20, 30, 40, or 50% of the nucleotide positions. The modified short interfering nucleic acid molecules of the invention can also be referred to as short interfering modified oligonucleotides "siMON." As used herein, the term siNA is meant to be equivalent to other terms used to describe nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically-modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term RNAi is meant to be equivalent to other terms used to describe sequence specific RNA interference, such as post transcriptional gene silencing, or epigenetics. For example, siNA molecules of the invention can be used to epigenetically silence genes at both the post-transcriptional level or the pre-transcriptional level. In a non-limiting example, epigenetic regulation of gene expression by

siNA molecules of the invention can result from siNA mediated modification of chromatin structure to alter gene expression (see, for example, Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237).

5 By "modulate" is meant that the expression of the gene, or level of RNA molecule or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits is up regulated or down regulated, such that expression, level, or activity is greater than or less than that observed in the absence of the modulator. For example, the term "modulate" can mean "inhibit," but the use of the word
10 "modulate" is not limited to this definition.

By "inhibit", "down-regulate", or "reduce", it is meant that the expression of the gene, or level of RNA molecules or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits, is reduced below that observed in the absence of the nucleic acid molecules (e.g., siNA) of the invention. In
15 one embodiment, inhibition, down-regulation or reduction with an siNA molecule is below that level observed in the presence of an inactive or attenuated molecule. In another embodiment, inhibition, down-regulation, or reduction with siNA molecules is below that level observed in the presence of, for example, an siNA molecule with scrambled sequence or with mismatches. In another embodiment, inhibition, down-regulation, or reduction of
20 gene expression with a nucleic acid molecule of the instant invention is greater in the presence of the nucleic acid molecule than in its absence.

By "gene" or "target gene" is meant, a nucleic acid that encodes an RNA, for example, nucleic acid sequences including, but not limited to, structural genes encoding a polypeptide. The target gene can be a gene derived from a cell, an endogenous gene, a transgene, or
25 exogenous genes such as genes of a pathogen, for example a virus, which is present in the cell after infection thereof. The cell containing the target gene can be derived from or contained in any organism, for example a plant, animal, protozoan, virus, bacterium, or fungus. Non-limiting examples of plants include monocots, dicots, or gymnosperms. Non-

limiting examples of animals include vertebrates or invertebrates. Non-limiting examples of fungi include molds or yeasts.

By "VEGF" as used herein is meant, any vascular endothelial growth factor (e.g., VEGF, VEGF-A, VEGF-B, VEGF-C, VEGF-D) protein, peptide, or polypeptide having
5 vascular endothelial growth factor activity, such as encoded by VEGF Genbank Accession Nos. shown in **Table I**. The term VEGF also refers to nucleic acid sequences encoding any vascular endothelial growth factor protein, peptide, or polypeptide having vascular endothelial growth factor activity.

By "VEGF-B" is meant, protein, peptide, or polypeptide receptor or a derivative
10 thereof, such as encoded by Genbank Accession No. NM_003377, having vascular endothelial growth factor type B activity. The term VEGF-B also refers to nucleic acid sequences encoding any VEGF-B protein, peptide, or polypeptide having VEGF-B activity.

By "VEGF-C" is meant, protein, peptide, or polypeptide receptor or a derivative
15 thereof, such as encoded by Genbank Accession No. NM_005429, having vascular endothelial growth factor type C activity. The term VEGF-C also refers to nucleic acid sequences encoding any VEGF-C protein, peptide, or polypeptide having VEGF-C activity.

By "VEGF-D" is meant, protein, peptide, or polypeptide receptor or a derivative
20 thereof, such as encoded by Genbank Accession No. NM_004469, having vascular endothelial growth factor type D activity. The term VEGF-D also refers to nucleic acid sequences encoding any VEGF-D protein, peptide, or polypeptide having VEGF-D activity.

By "VEGFr" as used herein is meant, any vascular endothelial growth factor receptor
protein, peptide, or polypeptide (e.g., VEGFr1, VEGFr2, or VEGFr3, including both
membrane bound and/or soluble forms thereof) having vascular endothelial growth factor
receptor activity, such as encoded by VEGFr Genbank Accession Nos. shown in **Table I**.
25 The term VEGFr also refers to nucleic acid sequences encoding any vascular endothelial
growth factor receptor protein, peptide, or polypeptide having vascular endothelial growth
factor receptor activity.

By "VEGFr1" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM_002019, having vascular endothelial growth factor receptor type 1 (*flt*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF1 also refers to nucleic acid sequences encoding any VEGFr1 protein, peptide, or polypeptide having VEGFr1 activity.

By "VEGFr2" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM_002253, having vascular endothelial growth factor receptor type 2 (*kdr*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF2 also refers to nucleic acid sequences encoding any VEGFr2 protein, peptide, or polypeptide having VEGFr2 activity.

By "VEGFr3" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM_002020 having vascular endothelial growth factor receptor type 3 (*kdr*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF3 also refers to nucleic acid sequences encoding any VEGFr3 protein, peptide, or polypeptide having VEGFr3 activity.

By "highly conserved sequence region" is meant, a nucleotide sequence of one or more regions in a target gene does not vary significantly from one generation to the other or from one biological system to the other.

By "sense region" is meant a nucleotide sequence of a siNA molecule having complementarity to an antisense region of the siNA molecule. In addition, the sense region of a siNA molecule can comprise a nucleic acid sequence having homology with a target nucleic acid sequence.

By "antisense region" is meant a nucleotide sequence of a siNA molecule having complementarity to a target nucleic acid sequence. In addition, the antisense region of a siNA molecule can optionally comprise a nucleic acid sequence having complementarity to a sense region of the siNA molecule.

By "target nucleic acid" is meant any nucleic acid sequence whose expression or activity is to be modulated. The target nucleic acid can be DNA or RNA.

By "complementarity" is meant that a nucleic acid can form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types. In reference to the nucleic molecules of the present invention, the binding free energy for a nucleic acid molecule with its complementary sequence is sufficient to allow the relevant function of the nucleic acid to proceed, e.g., RNAi activity. Determination of binding free energies for nucleic acid molecules is well known in the art (see, e.g., Turner *et al.*, 1987, *CSH Symp. Quant. Biol.* LII pp.123-133; Frier *et al.*, 1986, *Proc. Nat. Acad. Sci.* USA 83:9373-9377; Turner *et al.*, 1987, *J. Am. Chem. Soc.* 109:3783-3785). A percent complementarity indicates the percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). "Perfectly complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence.

The siRNA molecules of the invention represent a novel therapeutic approach to treat a variety of pathologic indications or other conditions, such as tumor angiogenesis and cancer, including but not limited to breast cancer, lung cancer (including non-small cell lung carcinoma), prostate cancer, colorectal cancer, brain cancer, esophageal cancer, bladder cancer, pancreatic cancer, cervical cancer, head and neck cancer, skin cancers, nasopharyngeal carcinoma, liposarcoma, epithelial carcinoma, renal cell carcinoma, gallbladder adeno carcinoma, parotid adenocarcinoma, ovarian cancer, melanoma, lymphoma, glioma, endometrial sarcoma, multidrug resistant cancers, diabetic retinopathy, macular degeneration, neovascular glaucoma, myopic degeneration, arthritis, psoriasis, endometriosis, female reproduction, verruca vulgaris, angiofibroma of tuberous sclerosis, pot-wine stains, Sturge Weber syndrome, Kippel-Trenaunay-Weber syndrome, Osler-Weber-Rendu syndrome, renal disease such as Autosomal dominant polycystic kidney disease (ADPKD), and any other diseases or conditions that are related to or will respond to the levels of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 in a cell or tissue, alone or in

combination with other therapies. The reduction of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 expression (specifically VEGF, VEGFr1, VEGFr2 and/or VEGFr3 gene RNA levels) and thus reduction in the level of the respective protein relieves, to some extent, the symptoms of the disease or condition.ue

5 In one embodiment of the present invention, each sequence of a siNA molecule of the invention is independently about 18 to about 24 nucleotides in length, in specific embodiments about 18, 19, 20, 21, 22, 23, or 24 nucleotides in length. In another embodiment, the siNA duplexes of the invention independently comprise about 17 to about 23 base pairs (e.g., about 17, 18, 19, 20, 21, 22 or 23). In yet another embodiment, siNA
10 molecules of the invention comprising hairpin or circular structures are about 35 to about 55 (e.g., about 35, 40, 45, 50 or 55) nucleotides in length, or about 38 to about 44 (e.g., 38, 39, 40, 41, 42, 43 or 44) nucleotides in length and comprising about 16 to about 22 (e.g., about 16, 17, 18, 19, 20, 21 or 22) base pairs. Exemplary siNA molecules of the invention are shown in Table II. Exemplary synthetic siNA molecules of the invention are shown in
15 Tables III and IV and/or Figures 4-5.

As used herein "cell" is used in its usual biological sense, and does not refer to an entire multicellular organism, e.g., specifically does not refer to a human. The cell can be present in an organism, e.g., birds, plants and mammals such as humans, cows, sheep, apes, monkeys, swine, dogs, and cats. The cell can be prokaryotic (e.g., bacterial cell) or
20 eukaryotic (e.g., mammalian or plant cell). The cell can be of somatic or germ line origin, totipotent or pluripotent, dividing or non-dividing. The cell can also be derived from or can comprise a gamete or embryo, a stem cell, or a fully differentiated cell.

The siNA molecules of the invention are added directly, or can be complexed with cationic lipids, packaged within liposomes, or otherwise delivered to target cells or tissues.
25 The nucleic acid or nucleic acid complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, infusion pump or stent, with or without their incorporation in biopolymers. In particular embodiments, the nucleic acid molecules of the invention comprise sequences shown in Tables II-III and/or Figures 4-5. Examples of such nucleic acid molecules consist essentially of sequences defined in these tables and figures.

Furthermore, the chemically modified constructs described in **Table IV** can be applied to any siNA sequence of the invention.

In another aspect, the invention provides mammalian cells containing one or more siNA molecules of this invention. The one or more siNA molecules can independently be
5 targeted to the same or different sites.

By "RNA" is meant a molecule comprising at least one ribonucleotide residue. By "ribonucleotide" is meant a nucleotide with a hydroxyl group at the 2' position of a β -D-ribo-furanose moiety. The terms include double-stranded RNA, single-stranded RNA, isolated RNA such as partially purified RNA, essentially pure RNA, synthetic RNA,
10 recombinantly produced RNA, as well as altered RNA that differs from naturally occurring RNA by the addition, deletion, substitution and/or alteration of one or more nucleotides. Such alterations can include addition of non-nucleotide material, such as to the end(s) of the siNA or internally, for example at one or more nucleotides of the RNA. Nucleotides in the RNA molecules of the instant invention can also comprise non-standard nucleotides, such as
15 non-naturally occurring nucleotides or chemically synthesized nucleotides or deoxynucleotides. These altered RNAs can be referred to as analogs or analogs of naturally-occurring RNA.

By "subject" is meant an organism, which is a donor or recipient of explanted cells or the cells themselves. "Subject" also refers to an organism to which the nucleic acid
20 molecules of the invention can be administered. In one embodiment, a subject is a mammal or mammalian cells. In another embodiment, a subject is a human or human cells.

The term "phosphorothioate" as used herein refers to an internucleotide linkage having Formula I, wherein Z and/or W comprise a sulfur atom. Hence, the term phosphorothioate refers to both phosphorothioate and phosphorodithioate internucleotide linkages.

25 The term "universal base" as used herein refers to nucleotide base analogs that form base pairs with each of the natural DNA/RNA bases with little discrimination between them. Non-limiting examples of universal bases include C-phenyl, C-naphthyl and other aromatic derivatives, inosine, azole carboxamides, and nitroazole derivatives such as 3-nitropyrrole,

4-nitroindole, 5-nitroindole, and 6-nitroindole as known in the art (see for example Loakes, 2001, *Nucleic Acids Research*, 29, 2437-2447).

5 The term "acyclic nucleotide" as used herein refers to any nucleotide having an acyclic ribose sugar, for example where any of the ribose carbons (C1, C2, C3, C4, or C5), are independently or in combination absent from the nucleotide.

10 The nucleic acid molecules of the instant invention, individually, or in combination or in conjunction with other drugs, can be used to treat diseases or conditions discussed herein (e.g., cancers and othe proliferative conditions). For example, to treat a particular disease or condition, the siNA molecules can be administered to a subject or can be administered to other appropriate cells evident to those skilled in the art, individually or in combination with one or more drugs under conditions suitable for the treatment.

15 In a further embodiment, the siNA molecules can be used in combination with other known treatments to treat conditions or diseases discussed above. For example, the described molecules could be used in combination with one or more known therapeutic agents to treat a disease or condition. Non-limiting examples of other therapeutic agents that can be readily combined with a siNA molecule of the invention are enzymatic nucleic acid molecules, allosteric nucleic acid molecules, antisense, decoy, or aptamer nucleic acid molecules, antibodies such as monoclonal antibodies, small molecules, and other organic and/or inorganic compounds including metals, salts and ions.

20 In one embodiment, the invention features an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the invention, in a manner which allows expression of the siNA molecule. For example, the vector can contain sequence(s) encoding both strands of a siNA molecule comprising a duplex. The vector can also contain sequence(s) encoding a single nucleic acid molecule that is self-complementary and thus
25 forms a siNA molecule. Non-limiting examples of such expression vectors are described in Paul *et al.*, 2002, *Nature Biotechnology*, 19, 505; Miyagishi and Taira, 2002, *Nature Biotechnology*, 19, 497; Lee *et al.*, 2002, *Nature Biotechnology*, 19, 500; and Novina *et al.*, 2002, *Nature Medicine*, advance online publication doi:10.1038/nm725.

In another embodiment, the invention features a mammalian cell, for example, a human cell, including an expression vector of the invention.

In yet another embodiment, the expression vector of the invention comprises a sequence for a siNA molecule having complementarity to a RNA molecule referred to by a
5 Genbank Accession numbers, for example Genbank Accession Nos. shown in Table I.

In one embodiment, an expression vector of the invention comprises a nucleic acid sequence encoding two or more siNA molecules, which can be the same or different.

In another aspect of the invention, siNA molecules that interact with target RNA molecules and down-regulate gene encoding target RNA molecules (for example target
10 RNA molecules referred to by Genbank Accession numbers herein) are expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant
15 vectors capable of expressing the siNA molecules can be delivered as described herein, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of siNA molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the siNA molecules bind and down-regulate gene function or expression via RNA interference (RNAi). Delivery of siNA expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted
20 from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell.

By "vectors" is meant any nucleic acid- and/or viral-based technique used to deliver a desired nucleic acid.

Other features and advantages of the invention will be apparent from the following
25 description of the preferred embodiments thereof, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a non-limiting example of a scheme for the synthesis of siNA molecules. The complementary siNA sequence strands, strand 1 and strand 2, are synthesized in tandem and are connected by a cleavable linkage, such as a nucleotide succinate or abasic succinate, which can be the same or different from the cleavable linker used for solid phase synthesis on a solid support. The synthesis can be either solid phase or solution phase, in the example shown, the synthesis is a solid phase synthesis. The synthesis is performed such that a protecting group, such as a dimethoxytrityl group, remains intact on the terminal nucleotide of the tandem oligonucleotide. Upon cleavage and deprotection of the oligonucleotide, the two siNA strands spontaneously hybridize to form a siNA duplex, which allows the purification of the duplex by utilizing the properties of the terminal protecting group, for example by applying a trityl on purification method wherein only duplexes/oligonucleotides with the terminal protecting group are isolated.

Figure 2 shows a MALDI-TOV mass spectrum of a purified siNA duplex synthesized by a method of the invention. The two peaks shown correspond to the predicted mass of the separate siNA sequence strands. This result demonstrates that the siNA duplex generated from tandem synthesis can be purified as a single entity using a simple trityl-on purification methodology.

Figure 3 shows a non-limiting proposed mechanistic representation of target RNA degradation involved in RNAi. Double-stranded RNA (dsRNA), which is generated by RNA-dependent RNA polymerase (RdRP) from foreign single-stranded RNA, for example viral, transposon, or other exogenous RNA, activates the DICER enzyme that in turn generates siNA duplexes. Alternately, synthetic or expressed siNA can be introduced directly into a cell by appropriate means. An active siNA complex forms which recognizes a target RNA, resulting in degradation of the target RNA by the RISC endonuclease complex or in the synthesis of additional RNA by RNA-dependent RNA polymerase (RdRP), which can activate DICER and result in additional siNA molecules, thereby amplifying the RNAi response.

Figure 4A-F shows non-limiting examples of chemically-modified siNA constructs of the present invention. In the figure, N stands for any nucleotide (adenosine, guanosine, cytosine, uridine, or optionally thymidine, for example thymidine can be substituted in the overhanging regions designated by parenthesis (N N). Various modifications are shown for the sense and antisense strands of the siNA constructs.

Figure 4A: The sense strand comprises 21 nucleotides having four phosphorothioate 5'- and 3'-terminal internucleotide linkages, wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and four 5'-terminal phosphorothioate internucleotide linkages and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4B: The sense strand comprises 21 nucleotides wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4C: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4D: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein and wherein and all purine nucleotides that may be present are 2'-deoxy nucleotides. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4E: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the

two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4F: The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-deoxy nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand of constructs A-F comprise sequence complementary to any target nucleic acid sequence of the invention.

Figure 5A-F shows non-limiting examples of specific chemically-modified siNA sequences of the invention. A-F applies the chemical modifications described in **Figure 4A-F** to a VEGFr1 siNA sequence. Such chemical modifications can be applied to any sequence herein, such as any VEGF, VEGFr1, VEGFr2, or VEGFr3 sequence.

Figure 6 shows non-limiting examples of different siNA constructs of the invention. The examples shown (constructs 1, 2, and 3) have 19 representative base pairs; however, different embodiments of the invention include any number of base pairs described herein. Bracketed regions represent nucleotide overhangs, for example comprising about 1, 2, 3, or 4 nucleotides in length, preferably about 2 nucleotides. Constructs 1 and 2 can be used independently for RNAi activity. Construct 2 can comprise a polynucleotide or non-nucleotide linker, which can optionally be designed as a biodegradable linker. In one

embodiment, the loop structure shown in construct 2 can comprise a biodegradable linker that results in the formation of construct 1 *in vivo* and/or *in vitro*. In another example, construct 3 can be used to generate construct 2 under the same principle wherein a linker is used to generate the active siNA construct 2 *in vivo* and/or *in vitro*, which can optionally
5 utilize another biodegradable linker to generate the active siNA construct 1 *in vivo* and/or *in vitro*. As such, the stability and/or activity of the siNA constructs can be modulated based on the design of the siNA construct for use *in vivo* or *in vitro* and/or *in vitro*.

Figure 7A-C is a diagrammatic representation of a scheme utilized in generating an expression cassette to generate siNA hairpin constructs.

10 **Figure 7A:** A DNA oligomer is synthesized with a 5'-restriction site (R1) sequence followed by a region having sequence identical (sense region of siNA) to a predetermined VEGF and/or VEGFr target sequence, wherein the sense region comprises, for example, about 19, 20, 21, or 22 nucleotides (N) in length, which is followed by a loop sequence of defined sequence (X), comprising, for example, about 3 to about 10 nucleotides.

15 **Figure 7B:** The synthetic construct is then extended by DNA polymerase to generate a hairpin structure having self-complementary sequence that will result in a siNA transcript having specificity for a VEGF and/or VEGFr target sequence and having self-complementary sense and antisense regions.

20 **Figure 7C:** The construct is heated (for example to about 95°C) to linearize the sequence, thus allowing extension of a complementary second DNA strand using a primer to the 3'-restriction sequence of the first strand. The double-stranded DNA is then inserted into an appropriate vector for expression in cells. The construct can be designed such that a 3'-terminal nucleotide overhang results from the transcription, for example by engineering restriction sites and/or utilizing a poly-U termination region as described in Paul *et al.*, 2002,
25 *Nature Biotechnology*, 29, 505-508.

Figure 8A-C is a diagrammatic representation of a scheme utilized in generating an expression cassette to generate double-stranded siNA constructs.

Figure 8A: A DNA oligomer is synthesized with a 5'-restriction (R1) site sequence followed by a region having sequence identical (sense region of siNA) to a predetermined VEGF and/or VEGFr target sequence, wherein the sense region comprises, for example, about 19, 20, 21, or 22 nucleotides (N) in length, and which is followed by a 3'-restriction site (R2) which is adjacent to a loop sequence of defined sequence (X).

Figure 8B: The synthetic construct is then extended by DNA polymerase to generate a hairpin structure having self-complementary sequence.

Figure 8C: The construct is processed by restriction enzymes specific to R1 and R2 to generate a double-stranded DNA which is then inserted into an appropriate vector for expression in cells. The transcription cassette is designed such that a U6 promoter region flanks each side of the dsDNA which generates the separate sense and antisense strands of the siNA. Poly T termination sequences can be added to the constructs to generate U overhangs in the resulting transcript.

Figure 9A-E is a diagrammatic representation of a method used to determine target sites for siNA mediated RNAi within a particular target nucleic acid sequence, such as messenger RNA.

Figure 9A: A pool of siNA oligonucleotides are synthesized wherein the antisense region of the siNA constructs has complementarity to target sites across the target nucleic acid sequence, and wherein the sense region comprises sequence complementary to the antisense region of the siNA.

Figure 9B&C: (**Figure 9B**) The sequences are pooled and are inserted into vectors such that (**Figure 9C**) transfection of a vector into cells results in the expression of the siNA.

Figure 9D: Cells are sorted based on phenotypic change that is associated with modulation of the target nucleic acid sequence.

Figure 9E: The siNA is isolated from the sorted cells and is sequenced to identify efficacious target sites within the target nucleic acid sequence.

Figure 10 shows non-limiting examples of different stabilization chemistries (1-10) that can be used, for example, to stabilize the 3'-end of siNA sequences of the invention, including (1) [3-3']-inverted deoxyribose; (2) deoxyribonucleotide; (3) [5'-3']-3'-deoxyribonucleotide; (4) [5'-3']-ribonucleotide; (5) [5'-3']-3'-O-methyl ribonucleotide; (6) 3'-glyceryl; (7) [3'-5']-3'-deoxyribonucleotide; (8) [3'-3']-deoxyribonucleotide; (9) [5'-2']-deoxyribonucleotide; and (10) [5-3']-dideoxyribonucleotide. In addition to modified and unmodified backbone chemistries indicated in the figure, these chemistries can be combined with different backbone modifications as described herein, for example, backbone modifications having Formula I. In addition, the 2'-deoxy nucleotide shown 5' to the terminal modifications shown can be another modified or unmodified nucleotide or non-nucleotide described herein, for example modifications having any of Formulae I-VII or any combination thereof.

Figure 11 shows a non-limiting example of a strategy used to identify chemically modified siNA constructs of the invention that are nuclease resistance while preserving the ability to mediate RNAi activity. Chemical modifications are introduced into the siNA construct based on educated design parameters (e.g. introducing 2'-modifications, base modifications, backbone modifications, terminal cap modifications etc). The modified construct is tested in an appropriate system (e.g. human serum for nuclease resistance, shown, or an animal model for PK/delivery parameters). In parallel, the siNA construct is tested for RNAi activity, for example in a cell culture system such as a luciferase reporter assay). Lead siNA constructs are then identified which possess a particular characteristic while maintaining RNAi activity, and can be further modified and assayed once again. This same approach can be used to identify siNA-conjugate molecules with improved pharmacokinetic profiles, delivery, and RNAi activity.

Figure 12 shows a non-limiting example of siNA mediated inhibition of VEGF-induced angiogenesis using the rat corneal model of angiogenesis. siNA targeting site 2340 of VEGFr1 RNA 29695/29699 (shown as RPI No. sense strand/antisense strand) was compared to an inverted control siNA 29983/29984 (shown as RPI No. sense strand/antisense strand) at three different concentrations (1ug, 3ug, and 10ug) and compared to a VEGF control in which no siNA was administered. As shown in the Figure, siNA

constructs targeting VEGFr1 RNA can provide significant inhibition of angiogenesis in the rat corneal model.

Figure 13 shows a non-limiting example of reduction of VEGFr1 mRNA in A375 cells mediated by chemically-modified siNAs that target VEGFr1 mRNA. A549 cells were
5 transfected with 0.25 ug/well of lipid complexed with 25 nM siNA. A screen of siNA constructs (Stabilization "Stab" chemistries are shown in Table IV, constructs are referred to by RPI number, see Table III) comprising Stab 4/5 chemistry (RPI 31190/31193), Stab 1/2 chemistry (RPI 31183/31186 and RPI 31184/31187), and unmodified RNA (RPI 30075/30076) were compared to untreated cells, matched chemistry inverted control siNA
10 constructs, (RPI 31208/31211, RPI 31201/31204, RPI 31202/31205, and RPI 30077/30078) scrambled siNA control constructs (Scram1 and Scram2), and cells transfected with lipid alone (transfection control). All of the siNA constructs show significant reduction of VEGFr1 RNA expression.

DETAILED DESCRIPTION OF THE INVENTION

15 Mechanism of action of Nucleic Acid Molecules of the Invention

The discussion that follows discusses the proposed mechanism of RNA interference mediated by short interfering RNA as is presently known, and is not meant to be limiting and is not an admission of prior art. Applicant demonstrates herein that chemically-modified short interfering nucleic acids possess similar or improved capacity to mediate
20 RNAi as do siRNA molecules and are expected to possess improved stability and activity *in vivo*; therefore, this discussion is not meant to be limiting only to siRNA and can be applied to siNA as a whole. By "improved capacity to mediate RNAi" or "improved RNAi activity" is meant to include RNAi activity measured *in vitro* and/or *in vivo* where the RNAi activity is a reflection of both the ability of the siNA to mediate RNAi and the stability of the siNAs
25 of the invention. In this invention, the product of these activities can be increased *in vitro* and/or *in vivo* compared to an all RNA siRNA or a siNA containing a plurality of ribonucleotides. In some cases, the activity or stability of the siNA molecule can be

decreased (i.e., less than ten-fold), but the overall activity of the siRNA molecule is enhanced *in vitro* and/or *in vivo*.

RNA interference refers to the process of sequence specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire *et al.*, 1998, *Nature*, 391, 806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes which is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2', 5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.

The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as Dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein *et al.*, 2001, *Nature*, 409, 363). Short interfering RNAs derived from Dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes. Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex containing a siRNA, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence homologous to the siRNA. Cleavage of the target RNA takes place in the middle of the region complementary to the guide sequence of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188). In addition, RNA interference can also involve small RNA

(e.g., micro-RNA or miRNA) mediated gene silencing, presumably through cellular mechanisms that regulate chromatin structure and thereby prevent transcription of target gene sequences (see for example Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 5 2002, *Science*, 297, 2232-2237). As such, siRNA molecules of the invention can be used to mediate gene silencing via interaction with RNA transcripts or alternately by interaction with particular gene sequences, wherein such interaction results in gene silencing either at the transcriptional level or post-transcriptional level.

RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, 10 were the first to observe RNAi in *C. elegans*. Wianny and Goetz, 1999, *Nature Cell Biol.*, 2, 70, describe RNAi mediated by dsRNA in mouse embryos. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa 15 cells. Recent work in *Drosophila* embryonic lysates has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity. These studies have shown that 21 nucleotide siRNA duplexes are most active when containing two 2-nucleotide 3'-terminal nucleotide overhangs. Furthermore, substitution of one or both siRNA strands with 2'-deoxy or 2'-O-methyl 20 nucleotides abolishes RNAi activity, whereas substitution of 3'-terminal siRNA nucleotides with deoxy nucleotides was shown to be tolerated. Mismatch sequences in the center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). 25 Other studies have indicated that a 5'-phosphate on the target-complementary strand of a siRNA duplex is required for siRNA activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309); however, siRNA molecules lacking a 5'-phosphate are active when introduced exogenously, suggesting that 5'-phosphorylation of siRNA constructs may occur *in vivo*.

Synthesis of Nucleic acid Molecules

Synthesis of nucleic acids greater than 100 nucleotides in length is difficult using automated methods, and the therapeutic cost of such molecules is prohibitive. In this invention, small nucleic acid motifs ("small" refers to nucleic acid motifs no more than 100
5 nucleotides in length, preferably no more than 80 nucleotides in length, and most preferably no more than 50 nucleotides in length; *e.g.*, individual siNA oligonucleotide sequences or siNA sequences synthesized in tandem) are preferably used for exogenous delivery. The simple structure of these molecules increases the ability of the nucleic acid to invade targeted regions of protein and/or RNA structure. Exemplary molecules of the instant
10 invention are chemically synthesized, and others can similarly be synthesized.

Oligonucleotides (*e.g.*, certain modified oligonucleotides or portions of oligonucleotides lacking ribonucleotides) are synthesized using protocols known in the art, for example as described in Caruthers *et al.*, 1992, *Methods in Enzymology* 211, 3-19, Thompson *et al.*, International PCT Publication No. WO 99/54459, Wincott *et al.*, 1995,
15 *Nucleic Acids Res.* 23, 2677-2684, Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, Brennan *et al.*, 1998, *Biotechnol Bioeng.*, 61, 33-45, and Brennan, U.S. Pat. No. 6,001,311. All of these references are incorporated herein by reference. The synthesis of oligonucleotides makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale
20 syntheses are conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 μ mol scale protocol with a 2.5 min coupling step for 2'-O-methylated nucleotides and a 45 sec coupling step for 2'-deoxy nucleotides or 2'-deoxy-2'-fluoro nucleotides. Table V outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2 μ mol scale can be performed on a 96-well plate synthesizer, such as the
25 instrument produced by Protogene (Palo Alto, CA) with minimal modification to the cycle. A 33-fold excess (60 μ L of 0.11 M = 6.6 μ mol) of 2'-O-methyl phosphoramidite and a 105-fold excess of S-ethyl tetrazole (60 μ L of 0.25 M = 15 μ mol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 22-fold excess (40 μ L of 0.11 M = 4.4 μ mol) of deoxy phosphoramidite and a 70-fold excess of S-ethyl
30 tetrazole (40 μ L of 0.25 M = 10 μ mol) can be used in each coupling cycle of deoxy residues

relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%. Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in
5 methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); and oxidation solution is 16.9 mM I₂, 49 mM pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American
10 International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide, 0.05 M in acetonitrile) is used.

Deprotection of the DNA-based oligonucleotides is performed as follows: the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aq. methylamine (1 mL) at 65 °C for 10 min. After cooling
15 to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H₂O/3:1:1, vortexed and the supernatant is then added to the first supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder.

The method of synthesis used for RNA including certain siNA molecules of the invention follows the procedure as described in Usman *et al.*, 1987, *J. Am. Chem. Soc.*, 109, 7845; Scaringe *et al.*, 1990, *Nucleic Acids Res.*, 18, 5433; and Wincott *et al.*, 1995, *Nucleic
20 Acids Res.* 23, 2677-2684 Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, and makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale syntheses are
25 conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 µmol scale protocol with a 7.5 min coupling step for alkylsilyl protected nucleotides and a 2.5 min coupling step for 2'-O-methylated nucleotides. Table V outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2 µmol scale can be
30 done on a 96-well plate synthesizer, such as the instrument produced by Protogene (Palo Alto, CA) with minimal modification to the cycle. A 33-fold excess (60 µL of 0.11 M = 6.6

μmol) of 2'-O-methyl phosphoramidite and a 75-fold excess of S-ethyl tetrazole (60 μL of 0.25 M = 15 μmol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 66-fold excess (120 μL of 0.11 M = 13.2 μmol) of alkylsilyl (ribo) protected phosphoramidite and a 150-fold excess of S-ethyl tetrazole (120 μL of 0.25 M = 30 μmol) can be used in each coupling cycle of ribo residues relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%. Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); oxidation solution is 16.9 mM I₂, 49 mM pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide 0.05 M in acetonitrile) is used.

Deprotection of the RNA is performed using either a two-pot or one-pot protocol. For the two-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aq. methylamine (1 mL) at 65 °C for 10 min. After cooling to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H₂O/3:1:1, vortexed and the supernatant is then added to the first supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder. The base deprotected oligoribonucleotide is resuspended in anhydrous TEA/HF/NMP solution (300 μL of a solution of 1.5 mL *N*-methylpyrrolidinone, 750 μL TEA and 1 mL TEA•3HF to provide a 1.4 M HF concentration) and heated to 65 °C. After 1.5 h, the oligomer is quenched with 1.5 M NH₄HCO₃.

Alternatively, for the one-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution

of 33% ethanolic methylamine/DMSO: 1/1 (0.8 mL) at 65 °C for 15 min. The vial is brought to rt. TEA•3HF (0.1 mL) is added and the vial is heated at 65 °C for 15 min. The sample is cooled at -20 °C and then quenched with 1.5 M NH₄HCO₃.

For purification of the trityl-on oligomers, the quenched NH₄HCO₃ solution is loaded
5 onto a C-18 containing cartridge that had been prewashed with acetonitrile followed by 50 mM TEAA. After washing the loaded cartridge with water, the RNA is detritylated with 0.5% TFA for 13 min. The cartridge is then washed again with water, salt exchanged with 1 M NaCl and washed with water again. The oligonucleotide is then eluted with 30% acetonitrile.

10 The average stepwise coupling yields are typically >98% (Wincott *et al.*, 1995 *Nucleic Acids Res.* 23, 2677-2684). Those of ordinary skill in the art will recognize that the scale of synthesis can be adapted to be larger or smaller than the example described above including but not limited to 96-well format.

Alternatively, the nucleic acid molecules of the present invention can be synthesized
15 separately and joined together post-synthetically, for example, by ligation (Moore *et al.*, 1992, *Science* 256, 9923; Draper *et al.*, International PCT publication No. WO 93/23569; Shabarova *et al.*, 1991, *Nucleic Acids Research* 19, 4247; Bellon *et al.*, 1997, *Nucleosides & Nucleotides*, 16, 951; Bellon *et al.*, 1997, *Bioconjugate Chem.* 8, 204), or by hybridization following synthesis and/or deprotection.

20 The siNA molecules of the invention can also be synthesized via a tandem synthesis methodology as described in Example 1 herein, wherein both siNA strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a cleavable linker which is subsequently cleaved to provide separate siNA fragments or strands that hybridize and permit purification of the siNA duplex. The linker can be a polynucleotide linker or a
25 non-nucleotide linker. The tandem synthesis of siNA as described herein can be readily adapted to both multiwell/multiplate synthesis platforms such as 96 well or similarly larger multi-well platforms. The tandem synthesis of siNA as described herein can also be readily

adapted to large scale synthesis platforms employing batch reactors, synthesis columns and the like.

A siNA molecule can also be assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes
5 the antisense region of the RNA molecule.

The nucleic acid molecules of the present invention can be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H (for a review see Usman and Cedergren, 1992, *TIBS* 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163). siNA constructs can be purified
10 by gel electrophoresis using general methods or can be purified by high pressure liquid chromatography (HPLC; see Wincott *et al.*, *supra*, the totality of which is hereby incorporated herein by reference) and re-suspended in water.

In another aspect of the invention, siNA molecules of the invention are expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can be DNA
15 plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant vectors capable of expressing the siNA molecules can be delivered as described herein, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of siNA molecules.

20 Optimizing Activity of the nucleic acid molecule of the invention.

Chemically synthesizing nucleic acid molecules with modifications (base, sugar and/or phosphate) can prevent their degradation by serum ribonucleases, which can increase their potency (see *e.g.*, Eckstein *et al.*, International Publication No. WO 92/07065; Perrault
25 *et al.*, 1990 *Nature* 344, 565; Pieken *et al.*, 1991, *Science* 253, 314; Usman and Cedergren, 1992, *Trends in Biochem. Sci.* 17, 334; Usman *et al.*, International Publication No. WO 93/15187; and Rossi *et al.*, International Publication No. WO 91/03162; Sproat, U.S. Pat. No. 5,334,711; Gold *et al.*, U.S. Pat. No. 6,300,074; and Burgin *et al.*, *supra*; all of which are incorporated by reference herein). All of the above references describe various chemical

modifications that can be made to the base, phosphate and/or sugar moieties of the nucleic acid molecules described herein. Modifications that enhance their efficacy in cells, and removal of bases from nucleic acid molecules to shorten oligonucleotide synthesis times and reduce chemical requirements are desired.

5 There are several examples in the art describing sugar, base and phosphate modifications that can be introduced into nucleic acid molecules with significant enhancement in their nuclease stability and efficacy. For example, oligonucleotides are modified to enhance stability and/or enhance biological activity by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-O-allyl, 2'-H, nucleotide base modifications (for a review see Usman and Cedergren, 1992, *TIBS*, 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163; Burgin *et al.*, 1996, *Biochemistry*, 35, 14090). Sugar modification of nucleic acid molecules have been extensively described in the art (see Eckstein *et al.*, *International Publication* PCT No. WO 92/07065; Perrault *et al.* *Nature*, 1990, 344, 565-568; Pieken *et al.* *Science*, 1991, 253, 314-317; Usman and Cedergren, *Trends in Biochem. Sci.*, 1992, 17, 334-339; Usman *et al.* *International Publication* PCT No. WO 93/15187; Sproat, *U.S. Pat.* No. 5,334,711 and Beigelman *et al.*, 1995, *J. Biol. Chem.*, 270, 25702; Beigelman *et al.*, *International PCT* publication No. WO 97/26270; Beigelman *et al.*, *U.S. Pat.* No. 5,716,824; Usman *et al.*, *U.S. Pat.* No. 5,627,053; Woolf *et al.*, *International PCT Publication* No. WO 98/13526; Thompson *et al.*, *USSN* 60/082,404 which was filed on April 20, 1998; Karpeisky *et al.*, 1998, *Tetrahedron Lett.*, 39, 1131; Earnshaw and Gait, 1998, *Biopolymers (Nucleic Acid Sciences)*, 48, 39-55; Verma and Eckstein, 1998, *Annu. Rev. Biochem.*, 67, 99-134; and Burlina *et al.*, 1997, *Bioorg. Med. Chem.*, 5, 1999-2010; all of the references are hereby incorporated in their totality by reference herein). Such publications describe general methods and strategies to determine the location of incorporation of sugar, base and/or phosphate modifications and the like into nucleic acid molecules without modulating catalysis, and are incorporated by reference herein. In view of such teachings, similar modifications can be used as described herein to modify the siNA nucleic acid molecules of the instant invention so long as the ability of siNA to promote RNAi in cells is not significantly inhibited.

While chemical modification of oligonucleotide internucleotide linkages with phosphorothioate, phosphorodithioate, and/or 5'-methylphosphonate linkages improves stability, excessive modifications can cause some toxicity or decreased activity. Therefore, when designing nucleic acid molecules, the amount of these internucleotide linkages should be minimized. The reduction in the concentration of these linkages should lower toxicity, resulting in increased efficacy and higher specificity of these molecules.

Short interfering nucleic acid (siNA) molecules having chemical modifications that maintain or enhance activity are provided. Such a nucleic acid is also generally more resistant to nucleases than an unmodified nucleic acid. Accordingly, the *in vitro* and/or *in vivo* activity should not be significantly lowered. In cases in which modulation is the goal, therapeutic nucleic acid molecules delivered exogenously should optimally be stable within cells until translation of the target RNA has been modulated long enough to reduce the levels of the undesirable protein. This period of time varies between hours to days depending upon the disease state. Improvements in the chemical synthesis of RNA and DNA (Wincott *et al.*, 1995, *Nucleic Acids Res.* 23, 2677; Caruthers *et al.*, 1992, *Methods in Enzymology* 211,3-19 (incorporated by reference herein)) have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability, as described above.

In one embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) G-clamp nucleotides. A G-clamp nucleotide is a modified cytosine analog wherein the modifications confer the ability to hydrogen bond both Watson-Crick and Hoogsteen faces of a complementary guanine within a duplex, see for example Lin and Matteucci, 1998, *J. Am. Chem. Soc.*, 120, 8531-8532. A single G-clamp analog substitution within an oligonucleotide can result in substantially enhanced helical thermal stability and mismatch discrimination when hybridized to complementary oligonucleotides. The inclusion of such nucleotides in nucleic acid molecules of the invention results in both enhanced affinity and specificity to nucleic acid targets, complementary sequences, or template strands. In another embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) LNA "locked nucleic acid" nucleotides such as a 2', 4'-C methylene bicyclo

nucleotide (see for example Wengel *et al.*, International PCT Publication No. WO 00/66604 and WO 99/14226).

In another embodiment, the invention features conjugates and/or complexes of siNA molecules of the invention. Such conjugates and/or complexes can be used to facilitate
5 delivery of siNA molecules into a biological system, such as a cell. The conjugates and complexes provided by the instant invention can impart therapeutic activity by transferring therapeutic compounds across cellular membranes, altering the pharmacokinetics, and/or modulating the localization of nucleic acid molecules of the invention. The present invention encompasses the design and synthesis of novel conjugates and complexes for the
10 delivery of molecules, including, but not limited to, small molecules, lipids, phospholipids, nucleosides, nucleotides, nucleic acids, antibodies, toxins, negatively charged polymers and other polymers, for example proteins, peptides, hormones, carbohydrates, polyethylene glycols, or polyamines, across cellular membranes. In general, the transporters described are designed to be used either individually or as part of a multi-component system, with or
15 without degradable linkers. These compounds are expected to improve delivery and/or localization of nucleic acid molecules of the invention into a number of cell types originating from different tissues, in the presence or absence of serum (see Sullenger and Cech, U.S. Pat. No. 5,854,038). Conjugates of the molecules described herein can be attached to biologically active molecules via linkers that are biodegradable, such as
20 biodegradable nucleic acid linker molecules.

The term "biodegradable linker" as used herein, refers to a nucleic acid or non-nucleic acid linker molecule that is designed as a biodegradable linker to connect one molecule to another molecule, for example, a biologically active molecule to a siNA molecule of the invention or the sense and antisense strands of a siNA molecule of the invention. The
25 biodegradable linker is designed such that its stability can be modulated for a particular purpose, such as delivery to a particular tissue or cell type. The stability of a nucleic acid-based biodegradable linker molecule can be modulated by using various chemistries, for example combinations of ribonucleotides, deoxyribonucleotides, and chemically-modified nucleotides, such as 2'-O-methyl, 2'-fluoro, 2'-amino, 2'-O-amino, 2'-C-allyl, 2'-O-allyl, and
30 other 2'-modified or base modified nucleotides. The biodegradable nucleic acid linker

molecule can be a dimer, trimer, tetramer or longer nucleic acid molecule, for example, an oligonucleotide of about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides in length, or can comprise a single nucleotide with a phosphorus-based linkage, for example, a phosphoramidate or phosphodiester linkage. The biodegradable nucleic acid linker molecule can also comprise nucleic acid backbone, nucleic acid sugar, or nucleic acid base modifications.

The term "biodegradable" as used herein, refers to degradation in a biological system, for example enzymatic degradation or chemical degradation.

The term "biologically active molecule" as used herein, refers to compounds or molecules that are capable of eliciting or modifying a biological response in a system. Non-limiting examples of biologically active siNA molecules either alone or in combination with other molecules contemplated by the instant invention include therapeutically active molecules such as antibodies, hormones, antivirals, peptides, proteins, chemotherapeutics, small molecules, vitamins, co-factors, nucleosides, nucleotides, oligonucleotides, enzymatic nucleic acids, antisense nucleic acids, triplex forming oligonucleotides, 2,5-A chimeras, siNA, dsRNA, allozymes, aptamers, decoys and analogs thereof. Biologically active molecules of the invention also include molecules capable of modulating the pharmacokinetics and/or pharmacodynamics of other biologically active molecules, for example, lipids and polymers such as polyamines, polyamides, polyethylene glycol and other polyethers.

The term "phospholipid" as used herein, refers to a hydrophobic molecule comprising at least one phosphorus group. For example, a phospholipid can comprise a phosphorus-containing group and saturated or unsaturated alkyl group, optionally substituted with OH, COOH, oxo, amine, or substituted or unsubstituted aryl groups.

Therapeutic nucleic acid molecules (e.g., siNA molecules) delivered exogenously optimally are stable within cells until reverse transcription of the RNA has been modulated long enough to reduce the levels of the RNA transcript. The nucleic acid molecules are resistant to nucleases in order to function as effective intracellular therapeutic agents.

Improvements in the chemical synthesis of nucleic acid molecules described in the instant invention and in the art have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability as described above.

5 In yet another embodiment, siNA molecules having chemical modifications that maintain or enhance enzymatic activity of proteins involved in RNAi are provided. Such nucleic acids are also generally more resistant to nucleases than unmodified nucleic acids. Thus, *in vitro* and/or *in vivo* the activity should not be significantly lowered.

10 Use of the nucleic acid-based molecules of the invention will lead to better treatment of the disease progression by affording the possibility of combination therapies (*e.g.*, multiple siNA molecules targeted to different genes; nucleic acid molecules coupled with known small molecule modulators; or intermittent treatment with combinations of molecules, including different motifs and/or other chemical or biological molecules). The treatment of subjects with siNA molecules can also include combinations of different types of nucleic acid molecules, such as enzymatic nucleic acid molecules (ribozymes),
15 allozymes, antisense, 2,5-A oligoadenylate, decoys, and aptamers.

In another aspect a siNA molecule of the invention comprises one or more 5' and/or a 3'- cap structure, for example on only the sense siNA strand, the antisense siNA strand, or both siNA strands.

20 By "cap structure" is meant chemical modifications, which have been incorporated at either terminus of the oligonucleotide (see, for example, Adamic *et al.*, U.S. Pat. No. 5,998,203, incorporated by reference herein). These terminal modifications protect the nucleic acid molecule from exonuclease degradation, and may help in delivery and/or localization within a cell. The cap may be present at the 5'-terminus (5'-cap) or at the 3'-terminal (3'-cap) or may be present on both termini. In non-limiting examples, the 5'-cap is
25 selected from the group consisting of glyceryl, inverted deoxy abasic residue (moiety); 4',5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide, 4'-thio nucleotide; carbocyclic nucleotide; 1,5-anhydrohexitol nucleotide; L-nucleotides; alpha-nucleotides; modified base nucleotide; phosphorodithioate linkage; *threo*-pentofuransyl nucleotide;

acyclic 3',4'-seco nucleotide; acyclic 3,4-dihydroxybutyl nucleotide; acyclic 3,5-dihydroxypentyl nucleotide, 3'-3'-inverted nucleotide moiety; 3'-3'-inverted abasic moiety; 3'-2'-inverted nucleotide moiety; 3'-2'-inverted abasic moiety; 1,4-butanediol phosphate; 3'-phosphoramidate; hexylphosphate; aminohexyl phosphate; 3'-phosphate; 3'-phosphorothioate; phosphorodithioate; or bridging or non-bridging methylphosphonate moiety.

In non-limiting examples, the 3'-cap is selected from the group consisting of glyceryl, inverted deoxy abasic residue (moiety), 4', 5'-methylene nucleotide; 1-(beta-D-erythrofuranosyl) nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate; 3-aminopropyl phosphate; 6-aminohexyl phosphate; 1,2-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; L-nucleotide; alpha-nucleotide; modified base nucleotide; phosphorodithioate; *threo*-pentofuranosyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted nucleotide moiety; 5'-5'-inverted abasic moiety; 5'-phosphoramidate; 5'-phosphorothioate; 1,4-butanediol phosphate; 5'-amino; bridging and/or non-bridging 5'-phosphoramidate, phosphorothioate and/or phosphorodithioate, bridging or non bridging methylphosphonate and 5'-mercapto moieties (for more details see Beaucage and Iyer, 1993, *Tetrahedron* 49, 1925; incorporated by reference herein).

By the term "non-nucleotide" is meant any group or compound which can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound is abasic in that it does not contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine and therefore lacks a base at the 1'-position.

An "alkyl" group refers to a saturated aliphatic hydrocarbon, including straight-chain, branched-chain, and cyclic alkyl groups. Preferably, the alkyl group has 1 to 12 carbons. More preferably, it is a lower alkyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkyl group can be substituted or unsubstituted. When substituted the substituted

group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO₂ or N(CH₃)₂, amino, or SH. The term also includes alkenyl groups that are unsaturated hydrocarbon groups containing at least one carbon-carbon double bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkenyl group has 1 to 12 carbons. More preferably, it is a lower alkenyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkenyl group may be substituted or unsubstituted. When substituted the substituted group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO₂, halogen, N(CH₃)₂, amino, or SH. The term "alkyl" also includes alkynyl groups that have an unsaturated hydrocarbon group containing at least one carbon-carbon triple bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkynyl group has 1 to 12 carbons. More preferably, it is a lower alkynyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkynyl group may be substituted or unsubstituted. When substituted the substituted group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO₂ or N(CH₃)₂, amino or SH.

Such alkyl groups can also include aryl, alkylaryl, carbocyclic aryl, heterocyclic aryl, amide and ester groups. An "aryl" group refers to an aromatic group that has at least one ring having a conjugated pi electron system and includes carbocyclic aryl, heterocyclic aryl and biaryl groups, all of which may be optionally substituted. The preferred substituent(s) of aryl groups are halogen, trihalomethyl, hydroxyl, SH, OH, cyano, alkoxy, alkyl, alkenyl, alkynyl, and amino groups. An "alkylaryl" group refers to an alkyl group (as described above) covalently joined to an aryl group (as described above). Carbocyclic aryl groups are groups wherein the ring atoms on the aromatic ring are all carbon atoms. The carbon atoms are optionally substituted. Heterocyclic aryl groups are groups having from 1 to 3 heteroatoms as ring atoms in the aromatic ring and the remainder of the ring atoms are carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen, and include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl and the like, all optionally substituted. An "amide" refers to an -C(O)-NH-R, where R is either alkyl, aryl, alkylaryl or hydrogen. An "ester" refers to an -C(O)-OR', where R is either alkyl, aryl, alkylaryl or hydrogen.

By "nucleotide" as used herein is as recognized in the art to include natural bases (standard), and modified bases well known in the art. Such bases are generally located at the 1' position of a nucleotide sugar moiety. Nucleotides generally comprise a base, sugar and a phosphate group. The nucleotides can be unmodified or modified at the sugar, phosphate and/or base moiety, (also referred to interchangeably as nucleotide analogs, modified nucleotides, non-natural nucleotides, non-standard nucleotides and other; see, for example, Usman and McSwiggen, *supra*; Eckstein *et al.*, International PCT Publication No. WO 92/07065; Usman *et al.*, International PCT Publication No. WO 93/15187; Uhlman & Peyman, *supra*, all are hereby incorporated by reference herein). There are several examples of modified nucleic acid bases known in the art as summarized by Limbach *et al.*, 1994, *Nucleic Acids Res.* 22, 2183. Some of the non-limiting examples of base modifications that can be introduced into nucleic acid molecules include, inosine, purine, pyridin-4-one, pyridin-2-one, phenyl, pseudouracil, 2, 4, 6-trimethoxy benzene, 3-methyluracil, dihydrouridine, naphthyl, aminophenyl, 5-alkylcytidines (*e.g.*, 5-methylcytidine), 5-alkyluridines (*e.g.*, ribothymidine), 5-halouridine (*e.g.*, 5-bromouridine) or 6-azapyrimidines or 6-alkylpyrimidines (*e.g.* 6-methyluridine), propyne, and others (Burgin *et al.*, 1996, *Biochemistry*, 35, 14090; Uhlman & Peyman, *supra*). By "modified bases" in this aspect is meant nucleotide bases other than adenine, guanine, cytosine and uracil at 1' position or their equivalents.

In one embodiment, the invention features modified siNA molecules, with phosphate backbone modifications comprising one or more phosphorothioate, phosphorodithioate, methylphosphonate, phosphotriester, morpholino, amidate carbamate, carboxymethyl, acetamidate, polyamide, sulfonate, sulfonamide, sulfamate, formacetal, thioformacetal, and/or alkylsilyl, substitutions. For a review of oligonucleotide backbone modifications, see Hunziker and Leumann, 1995, *Nucleic Acid Analogues: Synthesis and Properties*, in *Modern Synthetic Methods*, VCH, 331-417, and Mesmaeker *et al.*, 1994, *Novel Backbone Replacements for Oligonucleotides*, in *Carbohydrate Modifications in Antisense Research*, ACS, 24-39.

By "abasic" is meant sugar moieties lacking a base or having other chemical groups in place of a base at the 1' position, see for example Adamic *et al.*, U.S. Pat. No. 5,998,203.

By "unmodified nucleoside" is meant one of the bases adenine, cytosine, guanine, thymine, or uracil joined to the 1' carbon of β -D-ribo-furanose.

By "modified nucleoside" is meant any nucleotide base which contains a modification in the chemical structure of an unmodified nucleotide base, sugar and/or phosphate. Non-limiting examples of modified nucleotides are shown by Formulae I-VII and/or other
5 modifications described herein.

In connection with 2'-modified nucleotides as described for the present invention, by "amino" is meant 2'-NH₂ or 2'-O- NH₂, which can be modified or unmodified. Such modified groups are described, for example, in Eckstein *et al.*, U.S. Pat. No. 5,672,695 and
10 Matulic-Adamic *et al.*, U.S. Pat. No. 6,248,878, which are both incorporated by reference in their entireties.

Various modifications to nucleic acid siNA structure can be made to enhance the utility of these molecules. Such modifications will enhance shelf-life, half-life *in vitro*, stability, and ease of introduction of such oligonucleotides to the target site, *e.g.*, to enhance
15 penetration of cellular membranes, and confer the ability to recognize and bind to targeted cells.

Administration of Nucleic Acid Molecules

A siNA molecule of the invention can be adapted for use to treat, for example, tumor angiogenesis and cancer, including but not limited to breast cancer, lung cancer (including
20 non-small cell lung carcinoma), prostate cancer, colorectal cancer, brain cancer, esophageal cancer, bladder cancer, pancreatic cancer, cervical cancer, head and neck cancer, skin cancers, nasopharyngeal carcinoma, liposarcoma, epithelial carcinoma, renal cell carcinoma, gallbladder adeno carcinoma, parotid adenocarcinoma, ovarian cancer, melanoma, lymphoma, glioma, endometrial sarcoma, multidrug resistant cancers, diabetic retinopathy,
25 macular degeneration, neovascular glaucoma, myopic degeneration, arthritis, psoriasis, endometriosis, female reproduction, verruca vulgaris, angiofibroma of tuberous sclerosis, pot-wine stains, Sturge Weber syndrome, Kippel-Trenaunay-Weber syndrome, Osler-Weber-Rendu syndrome, renal disease such as Autosomal dominant polycystic kidney

disease (ADPKD), and any other diseases or conditions that are related to or will respond to the levels of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 in a cell or tissue, alone or in combination with other therapies. For example, a siNA molecule can comprise a delivery vehicle, including liposomes, for administration to a subject, carriers and diluents and their salts, and/or can be present in pharmaceutically acceptable formulations. Methods for the delivery of nucleic acid molecules are described in Akhtar *et al.*, 1992, *Trends Cell Bio.*, 2, 139; *Delivery Strategies for Antisense Oligonucleotide Therapeutics*, ed. Akhtar, 1995, Maurer *et al.*, 1999, *Mol. Membr. Biol.*, 16, 129-140; Hofland and Huang, 1999, *Handb. Exp. Pharmacol.*, 137, 165-192; and Lee *et al.*, 2000, *ACS Symp. Ser.*, 752, 184-192, all of which are incorporated herein by reference. Beigelman *et al.*, U.S. Pat. No. 6,395,713 and Sullivan *et al.*, PCT WO 94/02595 further describe the general methods for delivery of nucleic acid molecules. These protocols can be utilized for the delivery of virtually any nucleic acid molecule. Nucleic acid molecules can be administered to cells by a variety of methods known to those of skill in the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins (see for example Gonzalez *et al.*, 1999, *Bioconjugate Chem.*, 10, 1068-1074), biodegradable nanocapsules, and bioadhesive microspheres, or by proteinaceous vectors (O'Hare and Normand, International PCT Publication No. WO 00/53722). Alternatively, the nucleic acid/vehicle combination is locally delivered by direct injection or by use of an infusion pump. Direct injection of the nucleic acid molecules of the invention, whether subcutaneous, intramuscular, or intradermal, can take place using standard needle and syringe methodologies, or by needle-free technologies such as those described in Conry *et al.*, 1999, *Clin. Cancer Res.*, 5, 2330-2337 and Barry *et al.*, International PCT Publication No. WO 99/31262. The molecules of the instant invention can be used as pharmaceutical agents. Pharmaceutical agents prevent, modulate the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state in a subject.

Thus, the invention features a pharmaceutical composition comprising one or more nucleic acid(s) of the invention in an acceptable carrier, such as a stabilizer, buffer, and the like. The polynucleotides of the invention can be administered (*e.g.*, RNA, DNA or protein) and introduced into a subject by any standard means, with or without stabilizers, buffers, and

the like, to form a pharmaceutical composition. When it is desired to use a liposome delivery mechanism, standard protocols for formation of liposomes can be followed. The compositions of the present invention can also be formulated and used as tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions, suspensions for injectable administration, and the other compositions known in the art.

The present invention also includes pharmaceutically acceptable formulations of the compounds described. These formulations include salts of the above compounds, *e.g.*, acid addition salts, for example, salts of hydrochloric, hydrobromic, acetic acid, and benzene sulfonic acid.

A pharmacological composition or formulation refers to a composition or formulation in a form suitable for administration, *e.g.*, systemic administration, into a cell or subject, including for example a human. Suitable forms, in part, depend upon the use or the route of entry, for example oral, transdermal, or by injection. Such forms should not prevent the composition or formulation from reaching a target cell (*i.e.*, a cell to which the negatively charged nucleic acid is desirable for delivery). For example, pharmacological compositions injected into the blood stream should be soluble. Other factors are known in the art, and include considerations such as toxicity and forms that prevent the composition or formulation from exerting its effect.

By "systemic administration" is meant *in vivo* systemic absorption or accumulation of drugs in the blood stream followed by distribution throughout the entire body. Administration routes that lead to systemic absorption include, without limitation: intravenous, subcutaneous, intraperitoneal, inhalation, oral, intrapulmonary and intramuscular. Each of these administration routes exposes the siNA molecules of the invention to an accessible diseased tissue. The rate of entry of a drug into the circulation has been shown to be a function of molecular weight or size. The use of a liposome or other drug carrier comprising the compounds of the instant invention can potentially localize the drug, for example, in certain tissue types, such as the tissues of the reticular endothelial system (RES). A liposome formulation that can facilitate the association of drug with the surface of cells, such as, lymphocytes and macrophages is also useful. This approach can

provide enhanced delivery of the drug to target cells by taking advantage of the specificity of macrophage and lymphocyte immune recognition of abnormal cells, such as cells producing excess VEGF and/or VEGFr.

By "pharmaceutically acceptable formulation" is meant, a composition or formulation
5 that allows for the effective distribution of the nucleic acid molecules of the instant invention in the physical location most suitable for their desired activity. Non-limiting examples of agents suitable for formulation with the nucleic acid molecules of the instant invention include: P-glycoprotein inhibitors (such as Pluronic P85), which can enhance entry of drugs into the CNS (Jolliet-Riant and Tillement, 1999, *Fundam. Clin. Pharmacol.*,
10 13, 16-26); biodegradable polymers, such as poly (DL-lactide-coglycolide) microspheres for sustained release delivery after intracerebral implantation (Emerich, DF *et al.*, 1999, *Cell Transplant*, 8, 47-58) (Alkermes, Inc. Cambridge, MA); and loaded nanoparticles, such as those made of polybutylcyanoacrylate, which can deliver drugs across the blood brain barrier and can alter neuronal uptake mechanisms (*Prog Neuropsychopharmacol Biol*
15 *Psychiatry*, 23, 941-949, 1999). Other non-limiting examples of delivery strategies for the nucleic acid molecules of the instant invention include material described in Boado *et al.*, 1998, *J. Pharm. Sci.*, 87, 1308-1315; Tyler *et al.*, 1999, *FEBS Lett.*, 421, 280-284; Pardridge *et al.*, 1995, *PNAS USA.*, 92, 5592-5596; Boado, 1995, *Adv. Drug Delivery Rev.*, 15, 73-107; Aldrian-Herrada *et al.*, 1998, *Nucleic Acids Res.*, 26, 4910-4916; and Tyler *et al.*, 1999,
20 *PNAS USA.*, 96, 7053-7058.

The invention also features the use of the composition comprising surface-modified liposomes containing poly (ethylene glycol) lipids (PEG-modified, or long-circulating liposomes or stealth liposomes). These formulations offer a method for increasing the accumulation of drugs in target tissues. This class of drug carriers resists opsonization and
25 elimination by the mononuclear phagocytic system (MPS or RES), thereby enabling longer blood circulation times and enhanced tissue exposure for the encapsulated drug (Lasic *et al.* *Chem. Rev.* 1995, 95, 2601-2627; Ishiwata *et al.*, *Chem. Pharm. Bull.* 1995, 43, 1005-1011). Such liposomes have been shown to accumulate selectively in tumors, presumably by extravasation and capture in the neovascularized target tissues (Lasic *et al.*, *Science* 1995,
30 267, 1275-1276; Oku *et al.*, 1995, *Biochim. Biophys. Acta*, 1238, 86-90). The long-

circulating liposomes enhance the pharmacokinetics and pharmacodynamics of DNA and RNA, particularly compared to conventional cationic liposomes which are known to accumulate in tissues of the MPS (Liu *et al.*, *J. Biol. Chem.* 1995, 42, 24864-24870; Choi *et al.*, International PCT Publication No. WO 96/10391; Ansell *et al.*, International PCT Publication No. WO 96/10390; Holland *et al.*, International PCT Publication No. WO 96/10392). Long-circulating liposomes are also likely to protect drugs from nuclease degradation to a greater extent compared to cationic liposomes, based on their ability to avoid accumulation in metabolically aggressive MPS tissues such as the liver and spleen.

The present invention also includes compositions prepared for storage or administration that include a pharmaceutically effective amount of the desired compounds in a pharmaceutically acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co. (A.R. Gennaro edit. 1985), hereby incorporated by reference herein. For example, preservatives, stabilizers, dyes and flavoring agents can be provided. These include sodium benzoate, sorbic acid and esters of *p*-hydroxybenzoic acid. In addition, antioxidants and suspending agents can be used.

A pharmaceutically effective dose is that dose required to prevent, inhibit the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state. The pharmaceutically effective dose depends on the type of disease, the composition used, the route of administration, the type of mammal being treated, the physical characteristics of the specific mammal under consideration, concurrent medication, and other factors that those skilled in the medical arts will recognize. Generally, an amount between 0.1 mg/kg and 100 mg/kg body weight/day of active ingredients is administered dependent upon potency of the negatively charged polymer.

The nucleic acid molecules of the invention and formulations thereof can be administered orally, topically, parenterally, by inhalation or spray, or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and/or vehicles. The term parenteral as used herein includes percutaneous, subcutaneous, intravascular (*e.g.*, intravenous), intramuscular, or intrathecal injection or

infusion techniques and the like. In addition, there is provided a pharmaceutical formulation comprising a nucleic acid molecule of the invention and a pharmaceutically acceptable carrier. One or more nucleic acid molecules of the invention can be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or
5 adjuvants, and if desired other active ingredients. The pharmaceutical compositions containing nucleic acid molecules of the invention can be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use can be prepared according to any method known to
10 the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more such sweetening agents, flavoring agents, coloring agents or preservative agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can
15 be, for example, inert diluents; such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets can be uncoated or they can be coated by known techniques. In some cases such coatings can be
20 prepared by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate,
25 calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in a mixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropyl-methylcellulose,

sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents can be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions can also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions can contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents can be added to provide palatable oral preparations. These compositions can be preserved by the addition of an anti-oxidant such as ascorbic acid

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents or suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, can also be present.

Pharmaceutical compositions of the invention can also be in the form of oil-in-water emulsions. The oily phase can be a vegetable oil or a mineral oil or mixtures of these. Suitable emulsifying agents can be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for

example polyoxyethylene sorbitan monooleate. The emulsions can also contain sweetening and flavoring agents.

Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations can also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions can be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension can be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The nucleic acid molecules of the invention can also be administered in the form of suppositories, *e.g.*, for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Nucleic acid molecules of the invention can be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per subject per day). The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the host treated

and the particular mode of administration. Dosage unit forms generally contain between from about 1 mg to about 500 mg of an active ingredient.

It is understood that the specific dose level for any particular subject depends upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

For administration to non-human animals, the composition can also be added to the animal feed or drinking water. It can be convenient to formulate the animal feed and drinking water compositions so that the animal takes in a therapeutically appropriate quantity of the composition along with its diet. It can also be convenient to present the composition as a premix for addition to the feed or drinking water.

The nucleic acid molecules of the present invention can also be administered to a subject in combination with other therapeutic compounds to increase the overall therapeutic effect. The use of multiple compounds to treat an indication can increase the beneficial effects while reducing the presence of side effects.

In one embodiment, the invention comprises compositions suitable for administering nucleic acid molecules of the invention to specific cell types. For example, the asialoglycoprotein receptor (ASGPr) (Wu and Wu, 1987, *J. Biol. Chem.* 262, 4429-4432) is unique to hepatocytes and binds branched galactose-terminal glycoproteins, such as asialoorosomucoid (ASOR). In another example, the folate receptor is overexpressed in many cancer cells. Binding of such glycoproteins, synthetic glycoconjugates, or folates to the receptor takes place with an affinity that strongly depends on the degree of branching of the oligosaccharide chain, for example, triantennary structures are bound with greater affinity than biantennary or monoantennary chains (Baenziger and Fiete, 1980, *Cell*, 22, 611-620; Connolly *et al.*, 1982, *J. Biol. Chem.*, 257, 939-945). Lee and Lee, 1987, *Glycoconjugate J.*, 4, 317-328, obtained this high specificity through the use of N-acetyl-D-galactosamine as the carbohydrate moiety, which has higher affinity for the receptor, compared to galactose. This "clustering effect" has also been described for the binding and uptake of mannosyl-

terminating glycoproteins or glycoconjugates (Ponpipom *et al.*, 1981, *J. Med. Chem.*, 24, 1388-1395). The use of galactose, galactosamine, or folate based conjugates to transport exogenous compounds across cell membranes can provide a targeted delivery approach to, for example, the treatment of liver disease, cancers of the liver, or other cancers. The use of
5 bioconjugates can also provide a reduction in the required dose of therapeutic compounds required for treatment. Furthermore, therapeutic bioavailability, pharmacodynamics, and pharmacokinetic parameters can be modulated through the use of nucleic acid bioconjugates of the invention. Non-limiting examples of such bioconjugates are described in Vargeese *et al.*, USSN 10/201,394, filed August 13, 2001; and Matulic-Adamic *et al.*, USSN
10 60/362,016, filed March 6, 2002.

Alternatively, certain siNA molecules of the instant invention can be expressed within cells from eukaryotic promoters (*e.g.*, Izant and Weintraub, 1985, *Science*, 229, 345; McGarry and Lindquist, 1986, *Proc. Natl. Acad. Sci. USA* 83, 399; Scanlon *et al.*, 1991, *Proc. Natl. Acad. Sci. USA*, 88, 10591-5; Kashani-Sabet *et al.*, 1992, *Antisense Res. Dev.*, 2,
15 3-15; Dropulic *et al.*, 1992, *J. Virol.*, 66, 1432-41; Weerasinghe *et al.*, 1991, *J. Virol.*, 65, 5531-4; Ojwang *et al.*, 1992, *Proc. Natl. Acad. Sci. USA*, 89, 10802-6; Chen *et al.*, 1992, *Nucleic Acids Res.*, 20, 4581-9; Sarver *et al.*, 1990 *Science*, 247, 1222-1225; Thompson *et al.*, 1995, *Nucleic Acids Res.*, 23, 2259; Good *et al.*, 1997, *Gene Therapy*, 4, 45. Those skilled in the art realize that any nucleic acid can be expressed in eukaryotic cells from the
20 appropriate DNA/RNA vector. The activity of such nucleic acids can be augmented by their release from the primary transcript by a enzymatic nucleic acid (Draper *et al.*, PCT WO 93/23569, and Sullivan *et al.*, PCT WO 94/02595; Ohkawa *et al.*, 1992, *Nucleic Acids Symp. Ser.*, 27, 15-6; Taira *et al.*, 1991, *Nucleic Acids Res.*, 19, 5125-30; Ventura *et al.*, 1993, *Nucleic Acids Res.*, 21, 3249-55; Chowrira *et al.*, 1994, *J. Biol. Chem.*, 269, 25856.

25 In another aspect of the invention, RNA molecules of the present invention can be expressed from transcription units (see for example Couture *et al.*, 1996, *TIG.*, 12, 510) inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. In another embodiment, pol
30 III based constructs are used to express nucleic acid molecules of the invention (see for

example Thompson, U.S. Pats. Nos. 5,902,880 and 6,146,886). The recombinant vectors capable of expressing the siNA molecules can be delivered as described above, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of nucleic acid molecules. Such vectors can be repeatedly administered as necessary. Once
5 expressed, the siNA molecule interacts with the target mRNA and generates an RNAi response. Delivery of siNA molecule expressing vectors can be systemic, such as by intravenous or intra-muscular administration, by administration to target cells ex-planted from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell (for a review see Couture *et al.*, 1996,
10 *TIG.*, 12, 510).

In one aspect the invention features an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the instant invention. The expression vector can encode one or both strands of a siNA duplex, or a single self-complementary strand that self hybridizes into a siNA duplex. The nucleic acid sequences encoding the
15 siNA molecules of the instant invention can be operably linked in a manner that allows expression of the siNA molecule (see for example Paul *et al.*, 2002, *Nature Biotechnology*, 19, 505; Miyagishi and Taira, 2002, *Nature Biotechnology*, 19, 497; Lee *et al.*, 2002, *Nature Biotechnology*, 19, 500; and Novina *et al.*, 2002, *Nature Medicine*, advance online publication doi:10.1038/nm725).

20 In another aspect, the invention features an expression vector comprising: a) a transcription initiation region (*e.g.*, eukaryotic pol I, II or III initiation region); b) a transcription termination region (*e.g.*, eukaryotic pol I, II or III termination region); and c) a nucleic acid sequence encoding at least one of the siNA molecules of the instant invention, wherein said sequence is operably linked to said initiation region and said
25 termination region in a manner that allows expression and/or delivery of the siNA molecule. The vector can optionally include an open reading frame (ORF) for a protein operably linked on the 5' side or the 3'-side of the sequence encoding the siNA of the invention; and/or an intron (intervening sequences).

Transcription of the siNA molecule sequences can be driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters are expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type depends on the nature of the gene regulatory sequences (enhancers, silencers, etc.) present nearby. Prokaryotic RNA polymerase promoters are also used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein and Moss, 1990, *Proc. Natl. Acad. Sci. U S A*, 87, 6743-7; Gao and Huang 1993, *Nucleic Acids Res.*, 21, 2867-72; Lieber *et al.*, 1993, *Methods Enzymol.*, 217, 47-66; Zhou *et al.*, 1990, *Mol. Cell. Biol.*, 10, 4529-37). Several investigators have demonstrated that nucleic acid molecules expressed from such promoters can function in mammalian cells (e.g. Kashani-Sabet *et al.*, 1992, *Antisense Res. Dev.*, 2, 3-15; Ojwang *et al.*, 1992, *Proc. Natl. Acad. Sci. U S A*, 89, 10802-6; Chen *et al.*, 1992, *Nucleic Acids Res.*, 20, 4581-9; Yu *et al.*, 1993, *Proc. Natl. Acad. Sci. U S A*, 90, 6340-4; L'Huillier *et al.*, 1992, *EMBO J.*, 11, 4411-8; Lisiewicz *et al.*, 1993, *Proc. Natl. Acad. Sci. U. S. A.*, 90, 8000-4; Thompson *et al.*, 1995, *Nucleic Acids Res.*, 23, 2259; Sullenger & Cech, 1993, *Science*, 262, 1566). More specifically, transcription units such as the ones derived from genes encoding U6 small nuclear (snRNA), transfer RNA (tRNA) and adenovirus VA RNA are useful in generating high concentrations of desired RNA molecules such as siNA in cells (Thompson *et al.*, *supra*; Couture and Stinchcomb, 1996, *supra*; Noonberg *et al.*, 1994, *Nucleic Acid Res.*, 22, 2830; Noonberg *et al.*, U.S. Pat. No. 5,624,803; Good *et al.*, 1997, *Gene Ther.*, 4, 45; Beigelman *et al.*, International PCT Publication No. WO 96/18736. The above siNA transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated virus vectors), or viral RNA vectors (such as retroviral or alphavirus vectors) (for a review see Couture and Stinchcomb, 1996, *supra*).

In another aspect the invention features an expression vector comprising a nucleic acid sequence encoding at least one of the siNA molecules of the invention in a manner that allows expression of that siNA molecule. The expression vector comprises in one embodiment; a) a transcription initiation region; b) a transcription termination region; and c)

a nucleic acid sequence encoding at least one strand of the siNA molecule, wherein the sequence is operably linked to the initiation region and the termination region in a manner that allows expression and/or delivery of the siNA molecule.

5 In another embodiment the expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an open reading frame; and d) a nucleic acid sequence encoding at least one strand of a siNA molecule, wherein the sequence is operably linked to the 3'-end of the open reading frame and wherein the sequence is operably linked to the initiation region, the open reading frame and the termination region in a manner that allows expression and/or delivery of the siNA molecule. In yet another embodiment, the
10 expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an intron; and d) a nucleic acid sequence encoding at least one siNA molecule, wherein the sequence is operably linked to the initiation region, the intron and the termination region in a manner which allows expression and/or delivery of the nucleic acid molecule.

15 In another embodiment, the expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an intron; d) an open reading frame; and e) a nucleic acid sequence encoding at least one strand of a siNA molecule, wherein the sequence is operably linked to the 3'-end of the open reading frame and wherein the sequence is operably linked to the initiation region, the intron, the open reading frame and
20 the termination region in a manner which allows expression and/or delivery of the siNA molecule.

VEGF/VEGFr biology and biochemistry

The following discussion is adapted from R&D Systems, Cytokine Mini Reviews, Vascular Endothelial Growth Factor (VEGF), Copyright ©2002 R&D Systems.
25 Angiogenesis is a process of new blood vessel development from pre-existing vasculature. It plays an essential role in embryonic development, normal growth of tissues, wound healing, the female reproductive cycle (i.e., ovulation, menstruation and placental development), as well as a major role in many diseases. Particular interest has focused on cancer, since

tumors cannot grow beyond a few millimeters in size without developing a new blood supply. Angiogenesis is also necessary for the spread and growth of tumor cell metastases.

One of the most important growth and survival factors for endothelium is vascular endothelial growth factor (VEGF). VEGF induces angiogenesis and endothelial cell proliferation and plays an important role in regulating vasculogenesis. VEGF is a heparin-binding glycoprotein that is secreted as a homodimer of 45 kDa. Most types of cells, but usually not endothelial cells themselves, secrete VEGF. Since the initially discovered VEGF, VEGF-A, increases vascular permeability, it was known as vascular permeability factor. In addition, VEGF causes vasodilatation, partly through stimulation of nitric oxide synthase in endothelial cells. VEGF can also stimulate cell migration and inhibit apoptosis.

There are several splice variants of VEGF-A. The major ones include: 121, 165, 189 and 206 amino acids (aa), each one comprising a specific exon addition. VEGF165 is the most predominant protein, but transcripts of VEGF 121 may be more abundant. VEGF206 is rarely expressed and has been detected only in fetal liver. Recently, other splice variants of 145 and 183 aa have also been described. The 165, 189 and 206 aa splice variants have heparin-binding domains, which help anchor them in extracellular matrix and are involved in binding to heparin sulfate and presentation to VEGF receptors. Such presentation is a key factor for VEGF potency (i.e., the heparin-binding forms are more active). Several other members of the VEGF family have been cloned including VEGF-B, -C, and -D. Placenta growth factor (PlGF) is also closely related to VEGF-A. VEGF-A, -B, -C, -D, and PlGF are all distantly related to platelet-derived growth factors-A and -B. Less is known about the function and regulation of VEGF-B, -C, and -D, but they do not seem to be regulated by the major pathways that regulate VEGF-A.

VEGF-A transcription is potentiated in response to hypoxia and by activated oncogenes. The transcription factors, hypoxia inducible factor-1a (hif-1a) and -2a, are degraded by proteosomes in normoxia and stabilized in hypoxia. This pathway is dependent on the Von Hippel-Lindau gene product. Hif-1a and hif-2 a heterodimerize with the aryl hydrocarbon nuclear translocator in the nucleus and bind the VEGF promoter/enhancer. This is a key pathway expressed in most types of cells. Hypoxia inducibility, in particular,

characterizes VEGF-A versus other members of the VEGF family and other angiogenic factors. VEGF transcription in normoxia is activated by many oncogenes, including H-ras and several transmembrane tyrosine kinases, such as the epidermal growth factor receptor and erbB2. These pathways together account for a marked upregulation of VEGF-A in
5 tumors compared to normal tissues and are often of prognostic importance.

There are three receptors in the VEGF receptor family. They have the common properties of multiple IgG-like extracellular domains and tyrosine kinase activity. The enzyme domains of VEGF receptor 1 (VEGFr1, also known as Flt-1), VEGFr2 (also known as KDR or Flk-1), and VEGFr3 (also known as Flt-4) are divided by an inserted sequence.
10 Endothelial cells also express additional VEGF receptors, Neuropilin-1 and Neuropilin-2. VEGF-A binds to VEGFr1 and VEGFr2 and to Neuropilin-1 and Neuropilin-2. PlGF and VEGF-B bind VEGFr1 and Neuropilin-1. VEGF-C and -D bind VEGFr3 and VEGFr2.

The VEGF-C/VEGFr3 pathway is important for lymphatic proliferation. VEGFr3 is specifically expressed on lymphatic endothelium. A soluble form of Flt-1 can be detected in
15 peripheral blood and is a high affinity ligand for VEGF. Soluble Flt-1 can be used to antagonize VEGF function. VEGFr1 and VEGFr2 are upregulated in tumor and proliferating endothelium, partly by hypoxia and also in response to VEGF-A itself. VEGFr1 and VEGFr2 can interact with multiple downstream signaling pathways via proteins such as PLC-g, Ras, Shc, Nck, PKC and PI3-kinase. VEGFr1 is of higher affinity than VEGFr2 and
20 mediates motility and vascular permeability. VEGFr2 is necessary for proliferation.

VEGF can be detected in both plasma and serum samples of patients, with much higher levels in serum. Platelets release VEGF upon aggregation and may be a major source of VEGF delivery to tumors. Several studies have shown that association of high serum levels of VEGF with poor prognosis in cancer patients may be correlated with an elevated
25 platelet count. Many tumors release cytokines that can stimulate the production of megakaryocytes in the marrow and elevate the platelet count. This can result in an indirect increase of VEGF delivery to tumors.

VEGF is implicated in several other pathological conditions associated with enhanced angiogenesis. For example, VEGF plays a role in both psoriasis and rheumatoid arthritis. Diabetic retinopathy is associated with high intraocular levels of VEGF. Inhibition of VEGF function may result in infertility by blockade of corpus luteum function. Direct demonstration of the importance of VEGF in tumor growth has been achieved using dominant negative VEGF receptors to block in vivo proliferation, as well as blocking antibodies to VEGF39 or to VEGFr2.

The use of small interfering nucleic acid molecules targeting VEGF and corresponding receptors and ligands therefore provides a class of novel therapeutic agents that can be used in the diagnosis of and the treatment of cancer, proliferative diseases, or any other disease or condition that responds to modulation of VEGF and/or VEGFr genes.

Examples:

The following are non-limiting examples showing the selection, isolation, synthesis and activity of nucleic acids of the instant invention.

Example 1: Tandem synthesis of siNA constructs

Exemplary siNA molecules of the invention are synthesized in tandem using a cleavable linker, for example, a succinyl-based linker. Tandem synthesis as described herein is followed by a one-step purification process that provides RNAi molecules in high yield. This approach is highly amenable to siNA synthesis in support of high throughput RNAi screening, and can be readily adapted to multi-column or multi-well synthesis platforms.

After completing a tandem synthesis of a siNA oligo and its complement in which the 5'-terminal dimethoxytrityl (5'-O-DMT) group remains intact (trityl on synthesis), the oligonucleotides are deprotected as described above. Following deprotection, the siNA sequence strands are allowed to spontaneously hybridize. This hybridization yields a duplex in which one strand has retained the 5'-O-DMT group while the complementary strand comprises a terminal 5'-hydroxyl. The newly formed duplex behaves as a single molecule

during routine solid-phase extraction purification (Trityl-On purification) even though only one molecule has a dimethoxytrityl group. Because the strands form a stable duplex, this dimethoxytrityl group (or an equivalent group, such as other trityl groups or other hydrophobic moieties) is all that is required to purify the pair of oligos, for example, by using a C18 cartridge.

Standard phosphoramidite synthesis chemistry is used up to the point of introducing a tandem linker, such as an inverted deoxy abasic succinate or glyceryl succinate linker (see **Figure 1**) or an equivalent cleavable linker. A non-limiting example of linker coupling conditions that can be used includes a hindered base such as diisopropylethylamine (DIPA) and/or DMAP in the presence of an activator reagent such as Bromotripyrrolidinophosphoniumhexafluorophosphate (PyBrOP). After the linker is coupled, standard synthesis chemistry is utilized to complete synthesis of the second sequence leaving the terminal the 5'-O-DMT intact. Following synthesis, the resulting oligonucleotide is deprotected according to the procedures described herein and quenched with a suitable buffer, for example with 50mM NaOAc or 1.5M $\text{NH}_4\text{H}_2\text{CO}_3$.

Purification of the siNA duplex can be readily accomplished using solid phase extraction, for example using a Waters C18 SepPak 1g cartridge conditioned with 1 column volume (CV) of acetonitrile, 2 CV H_2O , and 2 CV 50mM NaOAc. The sample is loaded and then washed with 1 CV H_2O or 50mM NaOAc. Failure sequences are eluted with 1 CV 14% ACN (Aqueous with 50mM NaOAc and 50mM NaCl). The column is then washed, for example with 1 CV H_2O followed by on-column detritylation, for example by passing 1 CV of 1% aqueous trifluoroacetic acid (TFA) over the column, then adding a second CV of 1% aqueous TFA to the column and allowing to stand for approximately 10 minutes. The remaining TFA solution is removed and the column washed with H_2O followed by 1 CV 1M NaCl and additional H_2O . The siNA duplex product is then eluted, for example, using 1 CV 20% aqueous CAN.

Figure 2 provides an example of MALDI-TOV mass spectrometry analysis of a purified siNA construct in which each peak corresponds to the calculated mass of an individual siNA strand of the siNA duplex. The same purified siNA provides three peaks

when analyzed by capillary gel electrophoresis (CGE), one peak presumably corresponding to the duplex siNA, and two peaks presumably corresponding to the separate siNA sequence strands. Ion exchange HPLC analysis of the same siNA construct only shows a single peak. Testing of the purified siNA construct using a luciferase reporter assay described below demonstrated the same RNAi activity compared to siNA constructs generated from separately synthesized oligonucleotide sequence strands.

Example 2: Identification of potential siNA target sites in any RNA sequence

The sequence of an RNA target of interest, such as a viral or human mRNA transcript, is screened for target sites, for example by using a computer folding algorithm. In a non-limiting example, the sequence of a gene or RNA gene transcript derived from a database, such as Genbank, is used to generate siNA targets having complementarity to the target. Such sequences can be obtained from a database, or can be determined experimentally as known in the art. Target sites that are known, for example, those target sites determined to be effective target sites based on studies with other nucleic acid molecules, for example ribozymes or antisense, or those targets known to be associated with a disease or condition such as those sites containing mutations or deletions, can be used to design siNA molecules targeting those sites. Various parameters can be used to determine which sites are the most suitable target sites within the target RNA sequence. These parameters include but are not limited to secondary or tertiary RNA structure, the nucleotide base composition of the target sequence, the degree of homology between various regions of the target sequence, or the relative position of the target sequence within the RNA transcript. Based on these determinations, any number of target sites within the RNA transcript can be chosen to screen siNA molecules for efficacy, for example by using *in vitro* RNA cleavage assays, cell culture, or animal models. In a non-limiting example, anywhere from 1 to 1000 target sites are chosen within the transcript based on the size of the siNA construct to be used. High throughput screening assays can be developed for screening siNA molecules using methods known in the art, such as with multi-well or multi-plate assays to determine efficient reduction in target gene expression.

Example 3: Selection of siNA molecule target sites in a RNA

The following non-limiting steps can be used to carry out the selection of siNAs targeting a given gene sequence or transcript.

1. The target sequence is parsed *in silico* into a list of all fragments or subsequences of a particular length, for example 23 nucleotide fragments, contained within the target sequence. This step is typically carried out using a custom Perl script, but commercial sequence analysis programs such as Oligo, MacVector, or the GCG Wisconsin Package can be employed as well.
2. In some instances the siNAs correspond to more than one target sequence; such would be the case for example in targeting different transcripts of the same gene, targeting different transcripts of more than one gene, or for targeting both the human gene and an animal homolog. In this case, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find matching sequences in each list. The subsequences are then ranked according to the number of target sequences that contain the given subsequence; the goal is to find subsequences that are present in most or all of the target sequences. Alternately, the ranking can identify subsequences that are unique to a target sequence, such as a mutant target sequence. Such an approach would enable the use of siNA to target specifically the mutant sequence and not effect the expression of the normal sequence.
3. In some instances the siNA subsequences are absent in one or more sequences while present in the desired target sequence; such would be the case if the siNA targets a gene with a paralogous family member that is to remain untargeted. As in case 2 above, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find sequences that are present in the target gene but are absent in the untargeted paralog.
4. The ranked siNA subsequences can be further analyzed and ranked according to GC content. A preference can be given to sites containing 30-70% GC, with a further preference to sites containing 40-60% GC.

5. The ranked siNA subsequences can be further analyzed and ranked according to self-folding and internal hairpins. Weaker internal folds are preferred; strong hairpin structures are to be avoided.
6. The ranked siNA subsequences can be further analyzed and ranked according to whether they have runs of GGG or CCC in the sequence. GGG (or even more Gs) in either strand can make oligonucleotide synthesis problematic and can potentially interfere with RNAi activity, so it is avoided whenever better sequences are available. CCC is searched in the target strand because that will place GGG in the antisense strand.
7. The ranked siNA subsequences can be further analyzed and ranked according to whether they have the dinucleotide UU (uridine dinucleotide) on the 3'-end of the sequence, and/or AA on the 5'-end of the sequence (to yield 3' UU on the antisense sequence). These sequences allow one to design siNA molecules with terminal TT thymidine dinucleotides.
8. Four or five target sites are chosen from the ranked list of subsequences as described above. For example, in subsequences having 23 nucleotides, the right 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the upper (sense) strand of the siNA duplex, while the reverse complement of the left 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the lower (antisense) strand of the siNA duplex (see **Tables II and III**). If terminal TT residues are desired for the sequence (as described in paragraph 7), then the two 3' terminal nucleotides of both the sense and antisense strands are replaced by TT prior to synthesizing the oligos.
9. The siNA molecules are screened in an *in vitro*, cell culture or animal model system to identify the most active siNA molecule or the most preferred target site within the target RNA sequence.

In an alternate approach, a pool of siNA constructs specific to a VEGF and/or VEGFr target sequence is used to screen for target sites in cells expressing VEGF and/or VEGFr RNA, such as HUVEC, HMVEC, or A375 cells. The general strategy used in this approach

is shown in Figure 9. A non-limiting example of such is a pool comprising sequences having any of SEQ ID NOS 1-2238. Cells expressing VEGF and/or VEGFr (e.g., HUVEC, HMVEC, or A375 cells) are transfected with the pool of siNA constructs and cells that demonstrate a phenotype associated with VEGF and/or VEGFr inhibition are sorted. The pool of siNA constructs can be expressed from transcription cassettes inserted into appropriate vectors (see for example Figure 7 and Figure 8). The siNA from cells demonstrating a positive phenotypic change (e.g., decreased proliferation, decreased VEGF and/or VEGFr mRNA levels or decreased VEGF and/or VEGFr protein expression), are sequenced to determine the most suitable target site(s) within the target VEGF and/or VEGFr RNA sequence.

Example 4: VEGF and/or VEGFr targeted siNA design

siNA target sites were chosen by analyzing sequences of the VEGF and/or VEGFr RNA target and optionally prioritizing the target sites on the basis of folding (structure of any given sequence analyzed to determine siNA accessibility to the target), by using a library of siNA molecules as described in Example 3, or alternately by using an *in vitro* siNA system as described in Example 6 herein. siNA molecules were designed that could bind each target and are optionally individually analyzed by computer folding to assess whether the siNA molecule can interact with the target sequence. Varying the length of the siNA molecules can be chosen to optimize activity. Generally, a sufficient number of complementary nucleotide bases are chosen to bind to, or otherwise interact with, the target RNA, but the degree of complementarity can be modulated to accommodate siNA duplexes or varying length or base composition. By using such methodologies, siNA molecules can be designed to target sites within any known RNA sequence, for example those RNA sequences corresponding to the any gene transcript.

Chemically modified siNA constructs are designed to provide nuclease stability for systemic administration *in vivo* and/or improved pharmacokinetic, localization, and delivery properties while preserving the ability to mediate RNAi activity. Chemical modifications as described herein are introduced synthetically using synthetic methods described herein and those generally known in the art. The synthetic siNA constructs are then assayed for

nuclease stability in serum and/or cellular/tissue extracts (e.g. liver extracts). The synthetic siNA constructs are also tested in parallel for RNAi activity using an appropriate assay, such as a luciferase reporter assay as described herein or another suitable assay that can quantify RNAi activity. Synthetic siNA constructs that possess both nuclease stability and RNAi activity can be further modified and re-evaluated in stability and activity assays. The chemical modifications of the stabilized active siNA constructs can then be applied to any siNA sequence targeting any chosen RNA and used, for example, in target screening assays to pick lead siNA compounds for therapeutic development (see for example **Figure 11**).

Example 5: Chemical Synthesis and Purification of siNA

siNA molecules can be designed to interact with various sites in the RNA message, for example, target sequences within the RNA sequences described herein. The sequence of one strand of the siNA molecule(s) is complementary to the target site sequences described above. The siNA molecules can be chemically synthesized using methods described herein. Inactive siNA molecules that are used as control sequences can be synthesized by scrambling the sequence of the siNA molecules such that it is not complementary to the target sequence. Generally, siNA constructs can be synthesized using solid phase oligonucleotide synthesis methods as described herein (see for example Usman *et al.*, US Patent Nos. 5,804,683; 5,831,071; 5,998,203; 6,117,657; 6,353,098; 6,362,323; 6,437,117; 6,469,158; Scaringe *et al.*, US Patent Nos. 6,111,086; 6,008,400; 6,111,086 all incorporated by reference herein in their entirety).

In a non-limiting example, RNA oligonucleotides are synthesized in a stepwise fashion using the phosphoramidite chemistry as is known in the art. Standard phosphoramidite chemistry involves the use of nucleosides comprising any of 5'-O-dimethoxytrityl, 2'-O-tert-butyldimethylsilyl, 3'-O-2-Cyanoethyl N,N-diisopropylphosphoroamidite groups, and exocyclic amine protecting groups (e.g. N6-benzoyl adenosine, N4 acetyl cytidine, and N2-isobutyryl guanosine). Alternately, 2'-O-Silyl Ethers can be used in conjunction with acid-labile 2'-O-orthoester protecting groups in the synthesis of RNA as described by Scaringe *supra*. Differing 2' chemistries can require different protecting groups, for example 2'-deoxy-2'-amino nucleosides can utilize N-phthaloyl

protection as described by Usman *et al.*, US Patent 5,631,360, incorporated by reference herein in its entirety).

During solid phase synthesis, each nucleotide is added sequentially (3'- to 5'- direction) to the solid support-bound oligonucleotide. The first nucleoside at the 3'-end of the chain is covalently attached to a solid support (e.g., controlled pore glass or polystyrene) using various linkers. The nucleotide precursor, a ribonucleoside phosphoramidite, and activator are combined resulting in the coupling of the second nucleoside phosphoramidite onto the 5'-end of the first nucleoside. The support is then washed and any unreacted 5'-hydroxyl groups are capped with a capping reagent such as acetic anhydride to yield inactive 5'-acetyl moieties. The trivalent phosphorus linkage is then oxidized to a more stable phosphate linkage. At the end of the nucleotide addition cycle, the 5'-O-protecting group is cleaved under suitable conditions (e.g., acidic conditions for trityl-based groups and Fluoride for silyl-based groups). The cycle is repeated for each subsequent nucleotide.

Modification of synthesis conditions can be used to optimize coupling efficiency, for example by using differing coupling times, differing reagent/phosphoramidite concentrations, differing contact times, differing solid supports and solid support linker chemistries depending on the particular chemical composition of the siNA to be synthesized. Deprotection and purification of the siNA can be performed as is generally described in Usman *et al.*, US 5,831,071, US 6,353,098, US 6,437,117, and Bellon *et al.*, US 6,054,576, US 6,162,909, US 6,303,773, or Scaringe *supra*, incorporated by reference herein in their entireties. Additionally, deprotection conditions can be modified to provide the best possible yield and purity of siNA constructs. For example, applicant has observed that oligonucleotides comprising 2'-deoxy-2'-fluoro nucleotides can degrade under inappropriate deprotection conditions. Such oligonucleotides are deprotected using aqueous methylamine at about 35°C for 30 minutes. If the 2'-deoxy-2'-fluoro containing oligonucleotide also comprises ribonucleotides, after deprotection with aqueous methylamine at about 35°C for 30 minutes, TEA-HF is added and the reaction maintained at about 65°C for an additional 15 minutes.

Example 6: RNAi *in vitro* assay to assess siNA activity

An *in vitro* assay that recapitulates RNAi in a cell-free system is used to evaluate siNA constructs targeting VEGF and/or VEGFr RNA targets. The assay comprises the system described by Tuschl *et al.*, 1999, *Genes and Development*, 13, 3191-3197 and Zamore *et al.*,
5 2000, *Cell*, 101, 25-33 adapted for use with VEGF and/or VEGFr target RNA. A *Drosophila* extract derived from syncytial blastoderm is used to reconstitute RNAi activity *in vitro*. Target RNA is generated via *in vitro* transcription from an appropriate VEGF and/or VEGFr expressing plasmid using T7 RNA polymerase or via chemical synthesis as described herein. Sense and antisense siNA strands (for example 20 uM each) are annealed
10 by incubation in buffer (such as 100 mM potassium acetate, 30 mM HEPES-KOH, pH 7.4, 2 mM magnesium acetate) for 1 min. at 90°C followed by 1 hour at 37°C, then diluted in lysis buffer (for example 100 mM potassium acetate, 30 mM HEPES-KOH at pH 7.4, 2mM magnesium acetate). Annealing can be monitored by gel electrophoresis on an agarose gel in TBE buffer and stained with ethidium bromide. The *Drosophila* lysate is prepared using
15 zero to two-hour-old embryos from Oregon R flies collected on yeasted molasses agar that are dechorionated and lysed. The lysate is centrifuged and the supernatant isolated. The assay comprises a reaction mixture containing 50% lysate [vol/vol], RNA (10-50 pM final concentration), and 10% [vol/vol] lysis buffer containing siNA (10 nM final concentration). The reaction mixture also contains 10 mM creatine phosphate, 10 ug/ml creatine
20 phosphokinase, 100 uM GTP, 100 uM UTP, 100 uM CTP, 500 uM ATP, 5 mM DTT, 0.1 U/uL RNasin (Promega), and 100 uM of each amino acid. The final concentration of potassium acetate is adjusted to 100 mM. The reactions are pre-assembled on ice and preincubated at 25° C for 10 minutes before adding RNA, then incubated at 25° C for an additional 60 minutes. Reactions are quenched with 4 volumes of 1.25 x Passive Lysis
25 Buffer (Promega). Target RNA cleavage is assayed by RT-PCR analysis or other methods known in the art and are compared to control reactions in which siNA is omitted from the reaction.

Alternately, internally-labeled target RNA for the assay is prepared by *in vitro* transcription in the presence of [α -³²P] CTP, passed over a G 50 Sephadex column by
30 spin chromatography and used as target RNA without further purification. Optionally,

target RNA is 5'-³²P-end labeled using T4 polynucleotide kinase enzyme. Assays are performed as described above and target RNA and the specific RNA cleavage products generated by RNAi are visualized on an autoradiograph of a gel. The percentage of cleavage is determined by Phosphor Imager[®] quantitation of bands representing intact control RNA or RNA from control reactions without siNA and the cleavage products generated by the assay.

In one embodiment, this assay is used to determine target sites the VEGF and/or VEGFr RNA target for siNA mediated RNAi cleavage, wherein a plurality of siNA constructs are screened for RNAi mediated cleavage of the VEGF and/or VEGFr RNA target, for example, by analyzing the assay reaction by electrophoresis of labeled target RNA, or by northern blotting, as well as by other methodology well known in the art.

Example 7: Nucleic acid inhibition of VEGF and/or VEGFr target RNA *in vivo*

siNA molecules targeted to the huma VEGF and/or VEGFr RNA are designed and synthesized as described above. These nucleic acid molecules can be tested for cleavage activity *in vivo*, for example, using the following procedure. The target sequences and the nucleotide location within the VEGF and/or VEGFr RNA are given in **Table II and III**.

Two formats are used to test the efficacy of siNAs targeting VEGF and/or VEGFr. First, the reagents are tested in cell culture using, for example, HUVEC, HMVEC, or A375 cells to determine the extent of RNA and protein inhibition. siNA reagents (*e.g.*; see **Tables II and III**) are selected against the VEGF and/or VEGFr target as described herein. RNA inhibition is measured after delivery of these reagents by a suitable transfection agent to, for example, HUVEC, HMVEC, or A375 cells. Relative amounts of target RNA are measured versus actin using real-time PCR monitoring of amplification (*eg.*, ABI 7700 Taqman[®]). A comparison is made to a mixture of oligonucleotide sequences made to unrelated targets or to a randomized siNA control with the same overall length and chemistry, but randomly substituted at each position. Primary and secondary lead reagents are chosen for the target and optimization performed. After an optimal transfection agent concentration is chosen, a

RNA time-course of inhibition is performed with the lead siNA molecule. In addition, a cell-plating format can be used to determine RNA inhibition.

Delivery of siNA to Cells

Cells (e.g., HUVEC, HMVEC, or A375 cells) are seeded, for example, at 1×10^5 cells per well of a six-well dish in EGM-2 (BioWhittaker) the day before transfection. siNA (final concentration, for example 20nM) and cationic lipid (e.g., final concentration $2 \mu\text{g/ml}$) are complexed in EGM basal media (Biowhittaker) at 37°C for 30 mins in polystyrene tubes. Following vortexing, the complexed siNA is added to each well and incubated for the times indicated. For initial optimization experiments, cells are seeded, for example, at 1×10^3 in 96 well plates and siNA complex added as described. Efficiency of delivery of siNA to cells is determined using a fluorescent siNA complexed with lipid. Cells in 6-well dishes are incubated with siNA for 24 hours, rinsed with PBS and fixed in 2% paraformaldehyde for 15 minutes at room temperature. Uptake of siNA is visualized using a fluorescent microscope.

15 Taqman and Lightcycler quantification of mRNA

Total RNA is prepared from cells following siNA delivery, for example, using Qiagen RNA purification kits for 6-well or Rneasy extraction kits for 96-well assays. For Taqman analysis, dual-labeled probes are synthesized with the reporter dye, FAM or JOE, covalently linked at the 5'-end and the quencher dye TAMRA conjugated to the 3'-end. One-step RT-PCR amplifications are performed on, for example, an ABI PRISM 7700 Sequence Detector using 50 μl reactions consisting of 10 μl total RNA, 100 nM forward primer, 900 nM reverse primer, 100 nM probe, 1X TaqMan PCR reaction buffer (PE-Applied Biosystems), 5.5 mM MgCl_2 , 300 μM each dATP, dCTP, dGTP, and dTTP, 10U RNase Inhibitor (Promega), 1.25U AmpliTaq Gold (PE-Applied Biosystems) and 10U M-MLV Reverse Transcriptase (Promega). The thermal cycling conditions can consist of 30 min at 48°C , 10 min at 95°C , followed by 40 cycles of 15 sec at 95°C and 1 min at 60°C . Quantitation of mRNA levels is determined relative to standards generated from serially diluted total cellular RNA (300, 100, 33, 11 ng/rxn) and normalizing to β -actin or GAPDH mRNA in

parallel TaqMan reactions. For each gene of interest an upper and lower primer and a fluorescently labeled probe are designed. Real time incorporation of SYBR Green I dye into a specific PCR product can be measured in glass capillary tubes using a lightcycler. A standard curve is generated for each primer pair using control cRNA. Values are represented as relative expression to GAPDH in each sample.

Western blotting

Nuclear extracts can be prepared using a standard micro preparation technique (see for example Andrews and Faller, 1991, *Nucleic Acids Research*, 19, 2499). Protein extracts from supernatants are prepared, for example using TCA precipitation. An equal volume of 20% TCA is added to the cell supernatant, incubated on ice for 1 hour and pelleted by centrifugation for 5 minutes. Pellets are washed in acetone, dried and resuspended in water. Cellular protein extracts are run on a 10% Bis-Tris NuPage (nuclear extracts) or 4-12% Tris-Glycine (supernatant extracts) polyacrylamide gel and transferred onto nitro-cellulose membranes. Non-specific binding can be blocked by incubation, for example, with 5% non-fat milk for 1 hour followed by primary antibody for 16 hour at 4°C. Following washes, the secondary antibody is applied, for example (1:10,000 dilution) for 1 hour at room temperature and the signal detected with SuperSignal reagent (Pierce).

Example 8: Animal Models useful to evaluate the down-regulation of VEGF and/or VEGFr gene expression

There are several animal models in which the anti-angiogenesis effect of nucleic acids of the present invention, such as siRNA, directed against VEGF, VEGFr1, VEGFr2 and/or VEGFr3 mRNAs can be tested. Typically a corneal model has been used to study angiogenesis in rat and rabbit since recruitment of vessels can easily be followed in this normally avascular tissue (Pandey *et al.*, 1995 *Science* 268: 567-569). In these models, a small Teflon or Hydron disk pretreated with an angiogenesis factor (e.g. bFGF or VEGF) is inserted into a pocket surgically created in the cornea. Angiogenesis is monitored 3 to 5 days later. siRNA directed against VEGF, VEGFr1, VEGFr2 and/or VEGFr3 mRNAs are delivered in the disk as well, or dropwise to the eye over the time course of the experiment.

In another eye model, hypoxia has been shown to cause both increased expression of VEGF and neovascularization in the retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909; Shweiki *et al.*, 1992 *J. Clin. Invest.* 91: 2235-2243).

5 In human glioblastomas, it has been shown that VEGF is at least partially responsible for tumor angiogenesis (Plate *et al.*, 1992 *Nature* 359, 845). Animal models have been developed in which glioblastoma cells are implanted subcutaneously into nude mice and the progress of tumor growth and angiogenesis is studied (Kim *et al.*, 1993 *supra*; Millauer *et al.*, 1994 *supra*).

10 Another animal model that addresses neovascularization involves Matrigel, an extract of basement membrane that becomes a solid gel when injected subcutaneously (Passaniti *et al.*, 1992 *Lab. Invest.* 67: 519-528). When the Matrigel is supplemented with angiogenesis factors such as VEGF, vessels grow into the Matrigel over a period of 3 to 5 days and angiogenesis can be assessed. Again, nucleic acids directed against VEGF mRNAs are delivered in the Matrigel.

15 Several animal models exist for screening of anti-angiogenic agents. These include corneal vessel formation following corneal injury (Burger *et al.*, 1985 *Cornea* 4: 35-41; Lepri, *et al.*, 1994 *J. Ocular Pharmacol.* 10: 273-280; Ormerod *et al.*, 1990 *Am. J. Pathol.* 137: 1243-1252) or intracorneal growth factor implant (Grant *et al.*, 1993 *Diabetologia* 36: 282-291; Pandey *et al.* 1995 *supra*; Zieche *et al.*, 1992 *Lab. Invest.* 67: 711-715), vessel growth into Matrigel matrix containing growth factors (Passaniti *et al.*, 1992 *supra*), female reproductive organ neovascularization following hormonal manipulation (Shweiki *et al.*, 1993 *Clin. Invest.* 91: 2235-2243), several models involving inhibition of tumor growth in highly vascularized solid tumors (O'Reilly *et al.*, 1994 *Cell* 79: 315-328; Senger *et al.*, 1993 *Cancer and Metas. Rev.* 12: 303-324; Takahasi *et al.*, 1994 *Cancer Res.* 54: 4233-4237; Kim *et al.*, 1993 *supra*), and transient hypoxia-induced neovascularization in the mouse retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909).

The cornea model, described in Pandey *et al. supra*, is the most common and well characterized model for screening anti-angiogenic agent efficacy. This model involves an

avascular tissue into which vessels are recruited by a stimulating agent (growth factor, thermal or alkali burn, endotoxin). The corneal model utilizes the intrastromal corneal implantation of a Teflon pellet soaked in a VEGF-Hydron solution to recruit blood vessels toward the pellet, which can be quantitated using standard microscopic and image analysis techniques. To evaluate their anti-angiogenic efficacy, nucleic acids are applied topically to the eye or bound within Hydron on the Teflon pellet itself. This avascular cornea as well as the Matrigel (see below) provide for low background assays. While the corneal model has been performed extensively in the rabbit, studies in the rat have also been conducted.

The mouse model (Passaniti et al., *supra*) is a non-tissue model that utilizes Matrigel, an extract of basement membrane (Kleinman et al., 1986) or Millipore® filter disk, which can be impregnated with growth factors and anti-angiogenic agents in a liquid form prior to injection. Upon subcutaneous administration at body temperature, the Matrigel or Millipore® filter disk forms a solid implant. VEGF embedded in the Matrigel or Millipore® filter disk is used to recruit vessels within the matrix of the Matrigel or Millipore® filter disk which can be processed histologically for endothelial cell specific vWF (factor VIII antigen) immunohistochemistry, Trichrome-Masson stain, or hemoglobin content. Like the cornea, the Matrigel or Millipore® filter disk is avascular; however, it is not tissue. In the Matrigel or Millipore® filter disk model, nucleic acids are administered within the matrix of the Matrigel or Millipore® filter disk to test their anti-angiogenic efficacy. Thus, delivery issues in this model, as with delivery of nucleic acids by Hydron-coated Teflon pellets in the rat cornea model, may be less problematic due to the homogeneous presence of the nucleic acid within the respective matrix.

Other model systems to study tumor angiogenesis is reviewed by Folkman, 1985 *Adv. Cancer. Res.* 43, 175.

25 *Use of murine models*

For a typical systemic study involving 10 mice (20 g each) per dose group, 5 doses (1, 3, 10, 30 and 100 mg/kg daily over 14 days continuous administration), approximately 400

mg of siRNA, formulated in saline is used. A similar study in young adult rats (200 g) requires over 4 g. Parallel pharmacokinetic studies involve the use of similar quantities of siRNA further justifying the use of murine models.

Lewis lung carcinoma and B-16 melanoma murine models

- 5 Identifying a common animal model for systemic efficacy testing of nucleic acids is an efficient way of screening siRNA for systemic efficacy.

10 The Lewis lung carcinoma and B-16 murine melanoma models are well accepted models of primary and metastatic cancer and are used for initial screening of anti-cancer agents. These murine models are not dependent upon the use of immunodeficient mice, are relatively inexpensive, and minimize housing concerns. Both the Lewis lung and B-16 melanoma models involve subcutaneous implantation of approximately 10^6 tumor cells from metastatically aggressive tumor cell lines (Lewis lung lines 3LL or D122, LLc-LN7; B-16-BL6 melanoma) in C57BL/6J mice. Alternatively, the Lewis lung model can be produced by the surgical implantation of tumor spheres (approximately 0.8 mm in diameter).
15 Metastasis also can be modeled by injecting the tumor cells directly intravenously. In the Lewis lung model, microscopic metastases can be observed approximately 14 days following implantation with quantifiable macroscopic metastatic tumors developing within 21-25 days. The B-16 melanoma exhibits a similar time course with tumor neovascularization beginning 4 days following implantation. Since both primary and
20 metastatic tumors exist in these models after 21-25 days in the same animal, multiple measurements can be taken as indices of efficacy. Primary tumor volume and growth latency as well as the number of micro- and macroscopic metastatic lung foci or number of animals exhibiting metastases can be quantitated. The percent increase in lifespan can also be measured. Thus, these models provide suitable primary efficacy assays for screening
25 systemically administered siRNA nucleic acids and siRNA nucleic acid formulations .

In the Lewis lung and B-16 melanoma models, systemic pharmacotherapy with a wide variety of agents usually begins 1-7 days following tumor implantation/inoculation with either continuous or multiple administration regimens. Concurrent pharmacokinetic studies

can be performed to determine whether sufficient tissue levels of siRNA can be achieved for pharmacodynamic effect to be expected. Furthermore, primary tumors and secondary lung metastases can be removed and subjected to a variety of *in vitro* studies (*i.e.* target RNA reduction).

5 In addition, animal models are useful in screening compounds, eg. siRNA molecules, for efficacy in treating renal failure, such as a result of autosomal dominant polycystic kidney disease (ADPKD). The Han:SPRD rat model, mice with a targeted mutation in the Pkd2 gene and congenital polycystic kidney (cpk) mice, closely resemble human ADPKD and provide animal models to evaluate the therapeutic effect of siRNA constructs that have
10 the potential to interfere with one or more of the pathogenic elements of ADPKD mediated renal failure, such as angiogenesis. Angiogenesis may be necessary in the progression of ADPKD for growth of cyst cells as well as increased vascular permeability promoting fluid secretion into cysts. Proliferation of cystic epithelium is also a feature of ADPKD because cyst cells in culture produce soluble vascular endothelial growth factor (VEGF). VEGFr1
15 has also been detected in epithelial cells of cystic tubules but not in endothelial cells in the vasculature of cystic kidneys or normal kidneys. VEGFr2 expression is increased in endothelial cells of cyst vessels and in endothelial cells during renal ischemia-reperfusion. It is proposed that inhibition of VEGF receptors with anti-VEGFr1 and anti-VEGFr2 siRNA molecules would attenuate cyst formation, renal failure and mortality in ADPKD. Anti-
20 VEGFr2 siRNA molecules would therefore be designed to inhibit angiogenesis involved in cyst formation. As VEGFr1 is present in cystic epithelium and not in vascular endothelium of cysts, it is proposed that anti-VEGFr1 siRNA molecules would attenuate cystic epithelial cell proliferation and apoptosis which would in turn lead to less cyst formation. Further, it is proposed that VEGF produced by cystic epithelial cells is one of the stimuli for angiogenesis
25 as well as epithelial cell proliferation and apoptosis. The use of Han:SPRD rats (see for example Kaspareit-Rittinghausen *et al.*, 1991, *Am.J.Pathol.* 139, 693-696), mice with a targeted mutation in the Pkd2 gene (Pkd2^{-/-} mice, see for example Wu *et al.*, 2000, *Nat.Genet.* 24, 75-78) and cpk mice (see for example Woo *et al.*, 1994, *Nature*, 368, 750-753) all provide animal models to study the efficacy of siRNA molecules of the invention
30 against VEGFr1 and VEGFr2 mediated renal failure.

VEGF, VEGFr1 VGFR2 and/or VEGFr3 protein levels can be measured clinically or experimentally by FACS analysis. VEGF, VEGFr1 VGFR2 and/or VEGFr3 encoded mRNA levels are assessed by Northern analysis, RNase-protection, primer extension analysis and/or quantitative RT-PCR. siRNA nucleic acids that block VEGF, VEGFr1 VGFR2 and/or VEGFr3 protein encoding mRNAs and therefore result in decreased levels of VEGF, VEGFr1 VGFR2 and/or VEGFr3 activity by more than 20% *in vitro* can be identified.

Example 9: siNA-mediated inhibition of angiogenesis *in vivo*

The purpose of this study was to assess the anti-angiogenic activity of siNA targeted against VEGFr1 in the rat cornea model of VEGF induced angiogenesis (see above). The siNA molecules have matched inverted controls, which are inactive since they are not able to interact with the RNA target. The siNA molecules and VEGF were co-delivered using the filter disk method: Nitrocellulose filter disks (Millipore®) of 0.057 diameter were immersed in appropriate solutions and were surgically implanted in rat cornea as described by Pandey *et al.*, *supra*.

The stimulus for angiogenesis in this study was the treatment of the filter disk with 30 μ M VEGF, which is implanted within the cornea's stroma. This dose yields reproducible neovascularization stemming from the pericorneal vascular plexus growing toward the disk in a dose-response study 5 days following implant. Filter disks treated only with the vehicle for VEGF show no angiogenic response. The siNA were co-administered with VEGF on a disk in two different siNA concentrations. One concern with the simultaneous administration is that the siNA would not be able to inhibit angiogenesis since VEGF receptors could be stimulated. However, Applicant has observed that in low VEGF doses, the neovascular response reverts to normal, suggesting that the VEGF stimulus is essential for maintaining the angiogenic response. Blocking the production of VEGF receptors using simultaneous administration of anti-VEGF-R mRNA siNA could attenuate the normal neovascularization induced by the filter disk treated with VEGF.

Materials and Methods:

Test Compounds and Controls

R&D Systems VEGF, carrier free at 75 μ M in 82 mM Tris-Cl, pH 6.9

5 siNA, 1.67 μ G/ μ L, SITE 2340 (SEQ ID NO: 2; SEQ ID NO: 6) sense/antisense

siNA, 1.67 μ G/ μ L, INVERTED CONTROL FOR SITE 2340 (SEQ ID NO: 19; SEQ ID NO: 20) sense/antisense

siNA 1.67 μ g/ μ L, Site 2340 (SEQ ID NO: 419; SEQ ID NO: 420) sense/antisense

Animals

10

Harlan Sprague-Dawley Rats, Approximately 225-250g

45 males, 5 animals per group.

Husbandry

15

Animals are housed in groups of two. Feed, water, temperature and humidity are determined according to Pharmacology Testing Facility performance standards (SOP's) which are in accordance with the 1996 Guide for the Care and Use of Laboratory Animals (NRC). Animals are acclimated to the facility for at least 7 days prior to experimentation. During this time, animals are observed for overall health and sentinels are bled for baseline serology.

20

Experimental Groups

Each solution (VEGF and siNAs) was prepared as a 1X solution for final concentrations shown in the experimental groups described in Table III.

25 *siNA Annealing Conditions*

siNA sense and antisense strands are annealed for 1 minute in H₂O at 1.67mg/mL/strand followed by a 1 hour incubation at 37°C producing 3.34 mg/mL of duplexed siNA. For the 20µg/eye treatment, 6 µLs of the 3.34 mg/mL duplex is injected into the eye (see below). The 3.34 mg/mL duplex siNA can then be serially diluted for dose response assays.

Preparation of VEGF Filter Disk

For corneal implantation, 0.57 mm diameter nitrocellulose disks, prepared from 0.45 µm pore diameter nitrocellulose filter membranes (Millipore Corporation), were soaked for 30 min in 1 µL of 75 µM VEGF in 82 mM Tris·HCl (pH 6.9) in covered petri dishes on ice. Filter disks soaked only with the vehicle for VEGF (83 mM Tris-Cl pH 6.9) elicit no angiogenic response.

Corneal surgery

The rat corneal model used in this study was a modified from Koch *et al. Supra* and Pandey *et al., supra*. Briefly, corneas were irrigated with 0.5% povidone iodine solution followed by normal saline and two drops of 2% lidocaine. Under a dissecting microscope (Leica MZ-6), a stromal pocket was created and a presoaked filter disk (see above) was inserted into the pocket such that its edge was 1 mm from the corneal limbus.

Intraconjunctival injection of test solutions

Immediately after disk insertion, the tip of a 40-50 µm OD injector (constructed in our laboratory) was inserted within the conjunctival tissue 1 mm away from the edge of the corneal limbus that was directly adjacent to the VEGF-soaked filter disk. Six hundred nanoliters of test solution (siNA, inverted control or sterile water vehicle) were dispensed at a rate of 1.2 µL/min using a syringe pump (Kd Scientific). The injector was then removed, serially rinsed in 70% ethanol and sterile water and immersed in sterile water between each injection. Once the test solution was injected, closure of the eyelid was maintained using

microaneurism clips until the animal began to recover gross motor activity. Following treatment, animals were warmed on a heating pad at 37°C.

Quantitation of angiogenic response

5 Five days after disk implantation, animals were euthanized following administration of 0.4 mg/kg atropine and corneas were digitally imaged. The neovascular surface area (NSA, expressed in pixels) was measured *postmortem* from blood-filled corneal vessels using computerized morphometry (Image Pro Plus, Media Cybernetics, v2.0). The individual mean NSA was determined in triplicate from three regions of identical size in the area of
10 maximal neovascularization between the filter disk and the limbus. The number of pixels corresponding to the blood-filled corneal vessels in these regions was summated to produce an index of NSA. A group mean NSA was then calculated. Data from each treatment group were normalized to VEGF/siNA vehicle-treated control NSA and finally expressed as percent inhibition of VEGF-induced angiogenesis.

15 *Statistics*

After determining the normality of treatment group means, group mean percent inhibition of VEGF-induced angiogenesis was subjected to a one-way analysis of variance. This was followed by two post-hoc tests for significance including Dunnett's (comparison to
20 VEGF control) and Tukey-Kramer (all other group mean comparisons) at alpha = 0.05. Statistical analyses were performed using JMP v.3.1.6 (SAS Institute).

Results are graphically represented in **Figure 12**. As shown in **Figure 12**, VEGFr1 site 4229 active siNA (RPI 29695/29699) at three concentrations were effective at inhibiting angiogenesis compared to the inverted siNA control (RPI 2983/29984) and the VEGF
25 control. A chemically modified version of the VEGFr1 site 4229 active siNA comprising a sense strand having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with 5' and 3' terminal inverted deoxybasic residues (RPI 30196) and an antisense strand having having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with a terminal 3'-phosphorothioate internucleotide linkage (RPI 30416), showed similar inhibition. (Data not shown) This result shows siNA

molecules of differing chemically modified composition of the invention are capable of significantly inhibiting angiogenesis *in vivo*.

Example 10: RNAi mediated inhibition of VEGF and/or VEGFr RNA expression

siNA constructs (Table III) are tested for efficacy in reducing VEGF and/or VEGFr RNA expression in, for example, HUVEC, HMVEC, or A375 cells. Cells are plated approximately 24h before transfection in 96-well plates at 5,000-7,500 cells/well, 100 µl/well, such that at the time of transfection cells are 70-90% confluent. For transfection, annealed siNAs are mixed with the transfection reagent (Lipofectamine 2000, Invitrogen) in a volume of 50 µl/well and incubated for 20 min. at room temperature. The siNA transfection mixtures are added to cells to give a final siNA concentration of 25 nM in a volume of 150 µl. Each siNA transfection mixture is added to 3 wells for triplicate siNA treatments. Cells are incubated at 37° for 24h in the continued presence of the siNA transfection mixture. At 24h, RNA is prepared from each well of treated cells. The supernatants with the transfection mixtures are first removed and discarded, then the cells are lysed and RNA prepared from each well. Target gene expression following treatment is evaluated by RT-PCR for the target gene and for a control gene (36B4, an RNA polymerase subunit) for normalization. The triplicate data is averaged and the standard deviations determined for each treatment. Normalized data are graphed and the percent reduction of target mRNA by active siNAs in comparison to their respective inverted control siNAs is determined.

Figure 13 shows a non-limiting example of reduction of VEGFr1 mRNA in A375 cells mediated by chemically-modified siNAs that target VEGFr1 mRNA. A549 cells were transfected with 0.25 µg/well of lipid complexed with 25 nM siNA. A screen of siNA constructs (Stabilization "Stab" chemistries are shown in Table IV, constructs are referred to by RPI number, see Table III) comprising Stab 4/5 chemistry (RPI 31190/31193), Stab 1/2 chemistry (RPI 31183/31186 and RPI 31184/31187), and unmodified RNA (RPI 30075/30076) were compared to untreated cells, matched chemistry inverted control siNA constructs (RPI 31208/31211, RPI 31201/31204, RPI 31202/31205, and RPI 30077/30078), scrambled siNA control constructs (Scram1 and Scram2), and cells transfected with lipid

alone (transfection control). As shown in the figure, all of the siNA constructs significantly reduce VEGFr1 RNA expression. Additional stabilization chemistries as described in **Table IV** are similarly assayed for activity. These siNA constructs are compared to appropriate matched chemistry inverted controls. In addition, the siNA constructs are also compared to
5 untreated cells, cells transfected with lipid and scrambled siNA constructs, and cells transfected with lipid alone (transfection control).

Example 11: Indications

The present body of knowledge in VEGF and/or VEGFr research indicates the need for methods to assay VEGF and/or VEGFr activity and for compounds that can regulate
10 VEGF and/or VEGFr expression for research, diagnostic, and therapeutic use. As described herein, the nucleic acid molecules of the present invention can be used in assays to diagnose disease state related of VEGF and/or VEGFr levels. In addition, the nucleic acid molecules can be used to treat disease state related to VEGF and/or VEGFr levels.

Particular conditions and disease states that can be associated with VEGF and/or
15 VEGFr expression modulation include, but are not limited to:

1) Tumor angiogenesis: Angiogenesis has been shown to be necessary for tumors to grow into pathological size (Folkman, 1971, *PNAS* 76, 5217-5221; Wellstein & Czubayko, 1996, *Breast Cancer Res and Treatment* 38, 109-119). In addition, it allows tumor cells to travel through the circulatory system during metastasis. Increased levels of gene expression
20 of a number of angiogenic factors such as vascular endothelial growth factor (VEGF) have been reported in vascularized and edema-associated brain tumors (Berkman *et al.*, 1993 *J. Clin. Invest.* 91, 153). A more direct demonstration of the role of VEGF in tumor angiogenesis was demonstrated by Jim Kim *et al.*, 1993 *Nature* 362,841 wherein, monoclonal antibodies against VEGF were successfully used to inhibit the growth of
25 rhabdomyosarcoma, glioblastoma multiforme cells in nude mice. Similarly, expression of a dominant negative mutated form of the flt-1 VEGF receptor inhibits vascularization induced by human glioblastoma cells in nude mice (Millauer *et al.*, 1994, *Nature* 367, 576). Specific

tumor/cancer types that can be targeted using the nucleic acid molecules of the invention include but are not limited to the tumor/cancer types described herein.

2) Ocular diseases: Neovascularization has been shown to cause or exacerbate ocular diseases including, but not limited to, macular degeneration, neovascular glaucoma, diabetic
5 retinopathy, myopic degeneration, and trachoma (Norrby, 1997, *APMIS* 105, 417-437). Aiello *et al.*, 1994 *New Engl. J. Med.* 331, 1480, showed that the ocular fluid of a majority of patients suffering from diabetic retinopathy and other retinal disorders contains a high concentration of VEGF. Miller *et al.*, 1994 *Am. J. Pathol.* 145, 574, reported elevated levels of VEGF mRNA in patients suffering from retinal ischemia. These observations support a
10 direct role for VEGF in ocular diseases. Other factors, including those that stimulate VEGF synthesis, may also contribute to these indications.

3) Dermatological Disorders: Many indications have been identified which may be angiogenesis dependent, including but not limited to, psoriasis, verruca vulgaris, angiofibroma of tuberous sclerosis, port-wine stains, Sturge Weber syndrome, Kippel-
15 Trenaunay-Weber syndrome, and Osler-Weber-Rendu syndrome (Norrby, *supra*). Intradermal injection of the angiogenic factor b-FGF demonstrated angiogenesis in nude mice (Weckbecker *et al.*, 1992, *Angiogenesis: Key principles-Science-Technology-Medicine*, ed R. Steiner). Detmar *et al.*, 1994 *J. Exp. Med.* 180, 1141 reported that VEGF and its receptors were over-expressed in psoriatic skin and psoriatic dermal microvessels,
20 suggesting that VEGF plays a significant role in psoriasis.

4) Rheumatoid arthritis: Immunohistochemistry and *in situ* hybridization studies on tissues from the joints of patients suffering from rheumatoid arthritis show an increased level of VEGF and its receptors (Fava *et al.*, 1994 *J. Exp. Med.* 180, 341). Additionally, Koch *et al.*, 1994 *J. Immunol.* 152, 4149, found that VEGF-specific antibodies were able to
25 significantly reduce the mitogenic activity of synovial tissues from patients suffering from rheumatoid arthritis. These observations support a direct role for VEGF in rheumatoid arthritis. Other angiogenic factors including those of the present invention may also be involved in arthritis.

5) Endometriosis: Various studies indicate that VEGF is directly implicated in endometriosis. In one study, VEGF concentrations measured by ELISA in peritoneal fluid were found to be significantly higher in women with endometriosis than in women without endometriosis (24.1 ± 15 ng/ml vs 13.3 ± 7.2 ng/ml in normals). In patients with endometriosis, higher concentrations of VEGF were detected in the proliferative phase of the menstrual cycle (33 ± 13 ng/ml) compared to the secretory phase (10.7 ± 5 ng/ml). The cyclic variation was not noted in fluid from normal patients (McLaren *et al.*, 1996, *Human Reprod.* 11, 220-223). In another study, women with moderate to severe endometriosis had significantly higher concentrations of peritoneal fluid VEGF than women without endometriosis. There was a positive correlation between the severity of endometriosis and the concentration of VEGF in peritoneal fluid. In human endometrial biopsies, VEGF expression increased relative to the early proliferative phase approximately 1.6-, 2-, and 3.6-fold in midproliferative, late proliferative, and secretory endometrium (Shifren *et al.*, 1996, *J. Clin. Endocrinol. Metab.* 81, 3112-3118). In a third study, VEGF-positive staining of human ectopic endometrium was shown to be localized to macrophages (double immunofluorescent staining with CD14 marker). Peritoneal fluid macrophages demonstrated VEGF staining in women with and without endometriosis. However, increased activation of macrophages (acid phosphatase activity) was demonstrated in fluid from women with endometriosis compared with controls. Peritoneal fluid macrophage conditioned media from patients with endometriosis resulted in significantly increased cell proliferation ($[^3\text{H}]$ thymidine incorporation) in HUVEC cells compared to controls. The percentage of peritoneal fluid macrophages with VEGFr2 mRNA was higher during the secretory phase, and significantly higher in fluid from women with endometriosis ($80 \pm 15\%$) compared with controls ($32 \pm 20\%$). Flt-mRNA was detected in peritoneal fluid macrophages from women with and without endometriosis, but there was no difference between the groups or any evidence of cyclic dependence (McLaren *et al.*, 1996, *J. Clin. Invest.* 98, 482-489). In the early proliferative phase of the menstrual cycle, VEGF has been found to be expressed in secretory columnar epithelium (estrogen-responsive) lining both the oviducts and the uterus in female mice. During the secretory phase, VEGF expression was shown to have shifted to the underlying stroma composing the functional endometrium. In addition to examining the endometrium, neovascularization of ovarian

follicles and the corpus luteum, as well as angiogenesis in embryonic implantation sites have been analyzed. For these processes, VEGF was expressed in spatial and temporal proximity to forming vasculature (Shweiki *et al.*, 1993, *J. Clin. Invest.* 91, 2235-2243).

6) Kidney disease: Autosomal dominant polycystic kidney disease (ADPKD) is the most common life threatening hereditary disease in the USA. It affects about 1:400 to 1:1000 people and approximately 50% of people with ADPKD develop renal failure. ADPKD accounts for about 5-10% of end-stage renal failure in the USA, requiring dialysis and renal transplantation. Angiogenesis is implicated in the progression of ADPKD for growth of cyst cells, as well as increased vascular permeability promoting fluid secretion into cysts. Proliferation of cystic epithelium is a feature of ADPKD because cyst cells in culture produce soluble vascular endothelial growth factor (VEGF). VEGFr1 has been detected in epithelial cells of cystic tubules but not in endothelial cells in the vasculature of cystic kidneys or normal kidneys. VEGFr2 expression is increased in endothelial cells of cyst vessels and in endothelial cells during renal ischemia-reperfusion.

The use of radiation treatments and chemotherapeutics, such as Gemcytabine and cyclophosphamide, are non-limiting examples of chemotherapeutic agents that can be combined with or used in conjunction with the nucleic acid molecules (*e.g.* siNA molecules) of the instant invention. Those skilled in the art will recognize that other anti-cancer compounds and therapies can similarly be readily combined with the nucleic acid molecules of the instant invention (*e.g.* siNA molecules) and are hence within the scope of the instant invention. Such compounds and therapies are well known in the art (see for example *Cancer: Principles and Practice of Oncology*, Volumes 1 and 2, eds Devita, V.T., Hellman, S., and Rosenberg, S.A., J.B. Lippincott Company, Philadelphia, USA; incorporated herein by reference) and include, without limitation, folates, antifolates, pyrimidine analogs, fluoropyrimidines, purine analogs, adenosine analogs, topoisomerase I inhibitors, anthracyclins, retinoids, antibiotics, anthracyclins, platinum analogs, alkylating agents, nitrosoureas, plant derived compounds such as vinca alkaloids, epipodophyllotoxins, tyrosine kinase inhibitors, taxols, radiation therapy, surgery, nutritional supplements, gene therapy, radiotherapy, for example 3D-CRT, immunotoxin therapy, for example ricin, and

monoclonal antibodies. Specific examples of chemotherapeutic compounds that can be combined with or used in conjunction with the nucleic acid molecules of the invention include, but are not limited to, Paclitaxel; Docetaxel; Methotrexate; Doxorubin; Edatrexate; Vinorelbine; Tomaxifen; Leucovorin; 5-fluoro uridine (5-FU); Ionotecan; Cisplatin; Carboplatin; Amsacrine; Cytarabine; Bleomycin; Mitomycin C; Dactinomycin; Mithramycin; Hexamethylmelamine; Dacarbazine; L-asparaginase; Nitrogen mustard; Melphalan, Chlorambucil; Busulfan; Ifosfamide; 4-hydroperoxycyclophosphamide; Thiotepa; Irinotecan (CAMPTOSAR®, CPT-11, Camptothecin-11, Campto) Tamoxifen; Herceptin; IMC C225; ABX-EGF; and combinations thereof. The above list of compounds are non-limiting examples of compounds and/or methods that can be combined with or used in conjunction with the nucleic acid molecules (e.g. siNA) of the instant invention. Those skilled in the art will recognize that other drug compounds and therapies can similarly be readily combined with the nucleic acid molecules of the instant invention (e.g., siNA molecules) are hence within the scope of the instant invention.

Example 12: Diagnostic uses

The siNA molecules of the invention can be used in a variety of diagnostic applications, such as in the identification of molecular targets (e.g., RNA) in a variety of applications, for example, in clinical, industrial, environmental, agricultural and/or research settings. Such diagnostic use of siNA molecules involves utilizing reconstituted RNAi systems, for example, using cellular lysates or partially purified cellular lysates. siNA molecules of this invention can be used as diagnostic tools to examine genetic drift and mutations within diseased cells or to detect the presence of endogenous or exogenous, for example viral, RNA in a cell. The close relationship between siNA activity and the structure of the target RNA allows the detection of mutations in any region of the molecule, which alters the base-pairing and three-dimensional structure of the target RNA. By using multiple siNA molecules described in this invention, one can map nucleotide changes, which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with siNA molecules can be used to inhibit gene expression and define the role of specified gene products in the progression of disease or infection. In this manner, other genetic targets can be defined as important mediators of the disease. These experiments will

lead to better treatment of the disease progression by affording the possibility of combination therapies (e.g., multiple siNA molecules targeted to different genes, siNA molecules coupled with known small molecule inhibitors, or intermittent treatment with combinations siNA molecules and/or other chemical or biological molecules). Other *in vitro* uses of siNA molecules of this invention are well known in the art, and include detection of the presence of mRNAs associated with a disease, infection, or related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a siNA using standard methodologies, for example, fluorescence resonance emission transfer (FRET).

10 In a specific example, siNA molecules that cleave only wild-type or mutant forms of the target RNA are used for the assay. The first siNA molecules (*i.e.*, those that cleave only wild-type forms of target RNA) are used to identify wild-type RNA present in the sample and the second siNA molecules (*i.e.*, those that cleave only mutant forms of target RNA) are used to identify mutant RNA in the sample. As reaction controls, synthetic substrates of both wild-type and mutant RNA are cleaved by both siNA molecules to demonstrate the relative siNA efficiencies in the reactions and the absence of cleavage of the "non-targeted" RNA species. The cleavage products from the synthetic substrates also serve to generate size markers for the analysis of wild-type and mutant RNAs in the sample population. Thus, each analysis requires two siNA molecules, two substrates and one unknown sample, which is combined into six reactions. The presence of cleavage products is determined using an RNase protection assay so that full-length and cleavage fragments of each RNA can be analyzed in one lane of a polyacrylamide gel. It is not absolutely required to quantify the results to gain insight into the expression of mutant RNAs and putative risk of the desired phenotypic changes in target cells. The expression of mRNA whose protein product is implicated in the development of the phenotype (*i.e.*, disease related or infection related) is adequate to establish risk. If probes of comparable specific activity are used for both transcripts, then a qualitative comparison of RNA levels is adequate and decreases the cost of the initial diagnosis. Higher mutant form to wild-type ratios are correlated with higher risk whether RNA levels are compared qualitatively or quantitatively.

All patents and publications mentioned in the specification are indicative of the levels of skill of those skilled in the art to which the invention pertains. All references cited in this disclosure are incorporated by reference to the same extent as if each reference had been incorporated by reference in its entirety individually.

5 One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The methods and compositions described herein as presently representative of preferred embodiments are exemplary and are not intended as limitations
10 on the scope of the invention. Changes therein and other uses will occur to those skilled in the art, which are encompassed within the spirit of the invention, are defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications can be made to the invention disclosed herein without departing from the scope and spirit of the invention. Thus, such additional embodiments are within the scope of
15 the present invention and the following claims. The present invention teaches one skilled in the art to test various combinations and/or substitutions of chemical modifications described herein toward generating nucleic acid constructs with improved activity for mediating RNAi activity. Such improved activity can comprise improved stability, improved bioavailability, and/or improved activation of cellular responses mediating RNAi. Therefore, the specific
20 embodiments described herein are not limiting and one skilled in the art can readily appreciate that specific combinations of the modifications described herein can be tested without undue experimentation toward identifying siNA molecules with improved RNAi activity.

The invention illustratively described herein suitably can be practiced in the absence
25 of any element or elements, limitation or limitations that are not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of", and "consisting of" may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and

expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments, optional features, modification and
5 variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the description and the appended claims.

In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that
10 the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

Table I: VEGF and VEGFr Accession Numbers

NM_005429	Homo sapiens vascular endothelial growth factor C (VEGFC), mRNA gi 19924300 ref NM_005429.2 [19924300]
NM_003376	Homo sapiens vascular endothelial growth factor (VEGF), mRNA gi 19923239 ref NM_003376.2 [19923239]
AF095785	Homo sapiens vascular endothelial growth factor (VEGF) gene, promoter region and partial cds gi 4154290 gb AF095785.1 [4154290]
NM_003377	Homo sapiens vascular endothelial growth factor B (VEGFB), mRNA gi 20070172 ref NM_003377.2 [20070172]
AF486837	Homo sapiens vascular endothelial growth factor isoform VEGF165 (VEGF) mRNA, complete cds gi 19909064 gb AF486837.1 [19909064]
AF468110	

Homo sapiens vascular endothelial growth factor B isoform (VEGFB) gene, complete cds, alternatively spliced
gi|18766397|gb|AF468110.1|[18766397]

AF437895
Homo sapiens vascular endothelial growth factor (VEGF) gene, partial cds
gi|16660685|gb|AF437895.1|AF437895[16660685]

AY047581
Homo sapiens vascular endothelial growth factor (VEGF) mRNA, complete cds
gi|15422108|gb|AY047581.1|[15422108]

AF063657
Homo sapiens vascular endothelial growth factor receptor (FLT1) mRNA, complete cds
gi|3132830|gb|AF063657.1|AF063657[3132830]

AF092127
Homo sapiens vascular endothelial growth factor (VEGF) gene, partial sequence
gi|4139168|gb|AF092127.1|AF092127[4139168]

AF092126
Homo sapiens vascular endothelial growth factor (VEGF) gene, 5' UTR
gi|4139167|gb|AF092126.1|AF092126[4139167]

AF092125
Homo sapiens vascular endothelial growth factor (VEGF) gene, partial cds
gi|4139165|gb|AF092125.1|AF092125[4139165]

E15157
Human VEGF mRNA
gi|5709840|dbj|E15157.1||pat|JP|1998052285|2[5709840]

E15156
Human VEGF mRNA
gi|5709839|dbj|E15156.1||pat|JP|1998052285|1[5709839]

E14233
Human mRNA for vascular endothelial growth factor (VEGF), complete cds
gi|5708916|dbj|E14233.1||pat|JP|1997286795|1[5708916]

AF024710
Homo sapiens vascular endothelial growth factor (VEGF) mRNA, 3'UTR
gi|2565322|gb|AF024710.1|AF024710[2565322]

AJ010438
Homo sapiens mRNA for vascular endothelial growth factor, splicing variant
VEGF183
gi|3647280|emb|AJ010438.1|HSA010438[3647280]

AF098331
Homo sapiens vascular endothelial growth factor (VEGF) gene, promoter, partial
sequence
gi|4235431|gb|AF098331.1|AF098331[4235431]

AF022375
Homo sapiens vascular endothelial growth factor mRNA, complete cds
gi|3719220|gb|AF022375.1|AF022375[3719220]

AH006909
vascular endothelial growth factor {alternative splicing} [human, Genomic, 414
nt 5 segments]
gi|1680143|gb|AH006909.1||bbm|191843[1680143]

U01134
Human soluble vascular endothelial cell growth factor receptor (sflt) mRNA,
complete cds
gi|451321|gb|U01134.1|U01134[451321]

E14000
Human mRNA for FLT
gi|3252767|dbj|E14000.1||pat|JP|1997255700|1[3252767]

E13332
cDNA encoding vascular endothelial cell growth factor VEGF
gi|3252137|dbj|E13332.1||pat|JP|1997173075|1[3252137]

E13256
Human mRNA for FLT, complete cds
gi|3252061|dbj|E13256.1||pat|JP|1997154588|1[3252061]

AF063658
Homo sapiens vascular endothelial growth factor receptor 2 (KDR) mRNA, complete cds
gi|3132832|gb|AF063658.1|AF063658[3132832]

AJ000185
Homo Sapiens mRNA for vascular endothelial growth factor-D
gi|2879833|emb|AJ000185.1|HSAJ185[2879833]

D89630
Homo sapiens mRNA for VEGF-D, complete cds
gi|2780339|dbj|D89630.1|[2780339]

AF035121
Homo sapiens KDR/flk-1 protein mRNA, complete cds
gi|2655411|gb|AF035121.1|AF035121[2655411]

AF020393
Homo sapiens vascular endothelial growth factor C gene, partial cds and 5' upstream region
gi|2582366|gb|AF020393.1|AF020393[2582366]

Y08736

H.sapiens vegf gene, 3'UTR
gi|1619596|emb|Y08736.1|HSVEGF3UT[1619596]

X62568

H.sapiens vegf gene for vascular endothelial growth factor
gi|37658|emb|X62568.1|HSVEGF[37658]

X94216

H.sapiens mRNA for VEGF-C protein
gi|1177488|emb|X94216.1|HSVEGFC[1177488]

NM_002020

Homo sapiens fms-related tyrosine kinase 4 (FLT4), mRNA
gi|4503752|ref|NM_002020.1|[4503752]

NM_002253

Homo sapiens kinase insert domain receptor (a type III receptor tyrosine kinase)
(KDR), mRNA
gi|11321596|ref|NM_002253.1|[11321596]

Table II: VEGF and VEGFr siNA and Target Sequences

VEGFR1 g[4503748]ref[NM_002019.1

Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPos	Lower seq	Seq ID
1	GCGGACACUCCUCUGGCU	1	1	GCGGACACUCCUCUGGCU	1	23	AGCCGAGAGGAGUCCGC	428
19	UCCUCCCCGGCAGCGCGG	2	19	UCCUCCCCGGCAGCGCGG	2	41	CCGCCGUCGCGGGGAGG	429
37	GCGGUCGGAGCGGGCUCC	3	37	GCGGUCGGAGCGGGCUCC	3	59	GGAGCCCGCUCGCGAGCCG	430
55	CGGGGUCGCGGUGCAGCGG	4	55	CGGGGUCGCGGUGCAGCGG	4	77	CCGUCGACCCCGAGCCCG	431
73	GCCAGCGGGCCUGGCGCG	5	73	GCCAGCGGGCCUGGCGCG	5	95	CGCCGCCAGGCCCGCUGGC	432
91	GAGGAUUAACCGGGGAAGU	6	91	GAGGAUUAACCGGGGAAGU	6	113	ACUCCCCGGGUAAUCCUC	433
109	UGGUUGUCUCCUGGCU	7	109	UGGUUGUCUCCUGGCU	7	131	UCCAGCCAGGAGACAACA	434
127	AGCCGCGAGAGCGGCGCUC	8	127	AGCCGCGAGAGCGGCGCUC	8	149	GAGCGCCCGUCUCGCGGCU	435
145	CAGGCGCGGGCCGCGCGG	9	145	CAGGCGCGGGCCGCGCGG	9	167	CCGCCGGCCCGCGCCCGC	436
163	GCGGCGAACGAGAGACCG	10	163	GCGGCGAACGAGAGACCG	10	185	CCGUCCUCUCGUCGCGCG	437
181	GACUCUGCGCGCGGCGG	11	181	GACUCUGCGCGCGGCGG	11	203	CGACCCGGCCCGCCAGAGUC	438
199	GUUGCGCGGGGAGCGCGG	12	199	GUUGCGCGGGGAGCGCGG	12	221	CCGCGCUCGCGCCCGCCCAAC	439
217	GGCACCGGGCGAGCAGGCC	13	217	GGCACCGGGCGAGCAGGCC	13	239	GGCCUGCUCGCGCCCGGUGCC	440
235	CGCGUCGCGCUCACCAUGG	14	235	CGCGUCGCGCUCACCAUGG	14	257	CCAUGGUGAGCGCGACGCG	441
253	GUCAGCUACUGGGACACCG	15	253	GUCAGCUACUGGGACACCG	15	275	CGGUGUCCAGUAGCUGAC	442
271	GGGUCCUGCUGUGCGCGC	16	271	GGGUCCUGCUGUGCGCGC	16	293	GCGGCGACAGCAGGACCCC	443
289	CUGCUCAGCUCUGCUCUUC	17	289	CUGCUCAGCUCUGCUCUUC	17	311	GAAGCAGACAGCUGAGCAG	444
307	CUCACAGGAUCUAGUUCAG	18	307	CUCACAGGAUCUAGUUCAG	18	329	CUGAACUAGAUCCUGAG	445
325	GGUUCAAAUAUAAAGAU	19	325	GGUUCAAAUAUAAAGAU	19	347	GAUCUUUAUUUUUAACC	446
343	CCUGAACUGAGUUUAAAAG	20	343	CCUGAACUGAGUUUAAAAG	20	365	CUUUUAAACUACGUUCAGG	447
361	GGCACCCAGCACAUCAUGC	21	361	GGCACCCAGCACAUCAUGC	21	383	GCAUGAUGUGGUGGUGCC	448
379	CAAGCAGGCCAGACACUGC	22	379	CAAGCAGGCCAGACACUGC	22	401	GCAUGUGUGGCGGUGCUUG	449
397	CAUCUCCAAUGCAGGGGGG	23	397	CAUCUCCAAUGCAGGGGGG	23	419	CCCCCUGCAUUGGAGAU	450
415	GAAGCAGCCCAUAAUUGGU	24	415	GAAGCAGCCCAUAAUUGGU	24	437	ACCAUUUAUGGCGUCUUC	451
433	UCUUUGCCUGAAUUGGUGA	25	433	UCUUUGCCUGAAUUGGUGA	25	455	UCACCAUUUCAGGCAAGA	452
451	AGUAAGGAAAGCGAAAGGC	26	451	AGUAAGGAAAGCGAAAGGC	26	473	GCCUUUGCUUCCUUACU	453
469	CUGAGCAUAACUAAUUCUG	27	469	CUGAGCAUAACUAAUUCUG	27	491	CAGAUUAGUUAUGCUCAG	454
487	GCCUGUGGAAGAAUUGGCA	28	487	GCCUGUGGAAGAAUUGGCA	28	509	UGCCAUUUCUCCACAGGC	455
505	AAACAAUUCGACGACUUCU	29	505	AAACAAUUCGACGACUUCU	29	527	AAGUACUGCAGAAUUGUUU	456
523	UUAAACCUUGAACACAGCUC	30	523	UUAAACCUUGAACACAGCUC	30	545	GAGCUGUGUUAAGGUA	457

541	CAAGCAAAACCCACACUGGCU	31	541	CAAGCAAAACCCACACUGGCU	31	563	AGCCAGUGUGGUUUGCUUG	458
559	UUCUACAGCUGCAAAUAUC	32	559	UUCUACAGCUGCAAAUAUC	32	581	GAUUAUUUGCAGCUGUAGAA	459
577	CUAGCUGUACCUUACUCAA	33	577	CUAGCUGUACCUUACUCAA	33	599	UUGAAGUAGGUACAGCUAG	460
595	AAGAAGAAAGGAAACAGAAU	34	595	AAGAAGAAAGGAAACAGAAU	34	617	AUUUCUUUCCUUCUUCUU	461
613	UCUGCAAAUCUAUAUAUUA	35	613	UCUGCAAAUCUAUAUAUUA	35	635	UAAAUAUAUAGAUUGCAGA	462
631	AUUAGUGAUACAGGUAGAC	36	631	AUUAGUGAUACAGGUAGAC	36	653	GUCUACCCUGUAUCACUAAU	463
649	CCUUUCGUAGAGAUUUA	37	649	CCUUUCGUAGAGAUUUA	37	671	UGUACAUUCUUAACGAAAGG	464
667	AGUGAAAUCCCGAAAUUA	38	667	AGUGAAAUCCCGAAAUUA	38	689	UAAUUUCGGGGAUUAUCACU	465
685	AUACCAUAGACUGAAGGAA	39	685	AUACCAUAGACUGAAGGAA	39	707	UCCCUACAGUACUGUGUUA	466
703	AGGAGCUCUGCAUUCUCCU	40	703	AGGAGCUCUGCAUUCUCCU	40	725	AGGGAUUGACGAGCUCUCCU	467
721	UGCCGGGUUACGUCACCUA	41	721	UGCCGGGUUACGUCACCUA	41	743	UAGGUGACGUAACCCCGCA	468
739	AACAUCACUGUUAUUA	42	739	AACAUCACUGUUAUUA	42	761	UAAAAGUAACAGUGAUUU	469
757	AAAAAGUUUCCACUUGACA	43	757	AAAAAGUUUCCACUUGACA	43	779	UGUCAAGUGGAAACUUUUU	470
775	ACUUUGAUCCCUUGAUGGAA	44	775	ACUUUGAUCCCUUGAUGGAA	44	797	UCCCAUCAGGGAUCAAAGU	471
793	AAACGCAUUAUCUGGGACA	45	793	AAACGCAUUAUCUGGGACA	45	815	UGUCCCAGAUUAUGCGUUU	472
811	AGUAGAAAGGGCUUUAUCA	46	811	AGUAGAAAGGGCUUUAUCA	46	833	UGAUGAAGCCCUUUCUACU	473
829	AUAUCAAUUGCAACGUACA	47	829	AUAUCAAUUGCAACGUACA	47	851	UGUACGUUGCAUUUUGAUU	474
847	AAAGAAUAGGGCUUCUGA	48	847	AAAGAAUAGGGCUUCUGA	48	869	UCAGAAGCCCUAUAUUCUUU	475
865	ACCUGUGAAGCAACAGUCA	49	865	ACCUGUGAAGCAACAGUCA	49	887	UGACUGUUGCUUCACAGGU	476
883	AAUGGGCAUUUGUAUAGA	50	883	AAUGGGCAUUUGUAUAGA	50	905	UCUUAUACAAUUGCCCAUU	477
901	ACAAACUAUCUCACACAUC	51	901	ACAAACUAUCUCACACAUC	51	923	GAUGUGAGAUAGUUUGU	478
919	CGACAAACCAAUACAUAUCA	52	919	CGACAAACCAAUACAUAUCA	52	941	UGAUUGUAUUUGGUUUUGCG	479
937	AUAGAUUGUCCAAUAAGCA	53	937	AUAGAUUGUCCAAUAAGCA	53	959	UGCUUAUUUGGACAUUAU	480
955	ACACGACGCCAGUCAAU	54	955	ACACGACGCCAGUCAAU	54	977	AUUUGACUGGGCGUGGUGU	481
973	UUACUUAGAGGCCAUACUC	55	973	UUACUUAGAGGCCAUACUC	55	995	GAGUAUGGCCCUUAAGUAA	482
991	CUUGUCCUCAAUUGUACUG	56	991	CUUGUCCUCAAUUGUACUG	56	1013	CAGUACAAUUGAGGACAAG	483
1009	GCUACCACUCCCUUGAACA	57	1009	GCUACCACUCCCUUGAACA	57	1031	UGUJCAAGGGAGUGGUAGC	484
1027	ACGAGAGUUAUUAUGACCU	58	1027	ACGAGAGUUAUUAUGACCU	58	1049	AGGUCAUUUGAACUCUCGU	485
1045	UGGAGUUUACCCUGAUGAAA	59	1045	UGGAGUUUACCCUGAUGAAA	59	1067	UUUCAUCAGGGUAACUCCA	486
1063	AAAAUAGAGAGCUUCCG	60	1063	AAAAUAGAGAGCUUCCG	60	1085	CGGAAGCUCUCUUAUUUUU	487
1081	GUAGGCGACGAUUUGACC	61	1081	GUAGGCGACGAUUUGACC	61	1103	GGUCAAUUUGCGCCUUAAC	488
1099	CAAAGCAUUCCCAUGCCA	62	1099	CAAAGCAUUCCCAUGCCA	62	1121	UGGCAUGGGAAUUGCUUUG	489
1117	AACAUUUUACAGUGUUC	63	1117	AACAUUUUACAGUGUUC	63	1139	GAACACUGUAGAAUUGU	490
1135	CUUACUAUUUGACAAAUGC	64	1135	CUUACUAUUUGACAAAUGC	64	1157	GCAUUUUGCAAUAGUAG	491
1153	CAGAACAAAGACAAAGGAC	65	1153	CAGAACAAAGACAAAGGAC	65	1175	GUCCUUUGUCUUGUUCUG	492
1171	CUUUAUACUUUGUCUGUAA	66	1171	CUUUAUACUUUGUCUGUAA	66	1193	UUACACGACAAGUAUAAAG	493

1189	AGGAGUGGACCAUCAUUA	67	1189	AGGAGUGGACCAUCAUUA	67	1211	UGAAUGAUGGUCCACUCCU	494
1207	AAAUUGUUAACACCCUCAG	68	1207	AAAUUGUUAACACCCUCAG	68	1229	CUGAGGUGUUAACAGAUUU	495
1225	GUGCAUAUAUAUGAUAAG	69	1225	GUGCAUAUAUAUGAUAAG	69	1247	CUUUUAUAUAUAUGCAC	496
1243	GCAUUAUCACUGUGAAAC	70	1243	GCAUUAUCACUGUGAAAC	70	1265	GUUUCACAGUGAUGAUGC	497
1261	CAUCGAAACAGCAGGUGC	71	1261	CAUCGAAACAGCAGGUGC	71	1283	GCACCUGUGUUUUCGAUG	498
1279	CUUGAAACCGUAGCUGGCA	72	1279	CUUGAAACCGUAGCUGGCA	72	1301	UGCCAGCUACGGUUUUAAG	499
1297	AAGCGGCUUACCGGCUCU	73	1297	AAGCGGCUUACCGGCUCU	73	1319	AGAGCCGGUAAGACCGCUU	500
1315	UCUAUGAAAGUGAAGGCAU	74	1315	UCUAUGAAAGUGAAGGCAU	74	1337	AUGCCUUCACUUUUAUAGA	501
1333	UUUCCUCCGCGGAAGUUG	75	1333	UUUCCUCCGCGGAAGUUG	75	1355	CAACUUCGCGCGAGGGAAA	502
1351	GUAUGGUUAAAGAUUGGU	76	1351	GUAUGGUUAAAGAUUGGU	76	1373	ACCAUCUUUUUAACCAUAC	503
1369	UUACCUGCGACUGAGAAU	77	1369	UUACCUGCGACUGAGAAU	77	1391	AUUUCUCAGUCGCAGGUAA	504
1387	UCUGCUCGCUAUUUGACUC	78	1387	UCUGCUCGCUAUUUGACUC	78	1409	GAGUCAAAUAGCGAGCAGA	505
1405	CGUGGCUACUGUUAUUA	79	1405	CGUGGCUACUGUUAUUA	79	1427	UAAUUAACGAGUAGCCACG	506
1423	AUCAAGGACGUAACUGAAG	80	1423	AUCAAGGACGUAACUGAAG	80	1445	CUUCAGUUACGUCCUUGAU	507
1441	GAGGAUGCAGGGAUUUAUA	81	1441	GAGGAUGCAGGGAUUUAUA	81	1463	UAUAAUCCCGUGCAUCCUC	508
1459	ACAAUCUUGCUGAGCAUAA	82	1459	ACAAUCUUGCUGAGCAUAA	82	1481	UUAUGCUCAGCAAGAUUGU	509
1477	AAACAGUCAAAUGUGUUUA	83	1477	AAACAGUCAAAUGUGUUUA	83	1499	UAAACACAUUUGACUGUUU	510
1495	AAAAACCUACUGCCACUC	84	1495	AAAAACCUACUGCCACUC	84	1517	GAGUGCAGUGAGGUUUUU	511
1513	CUAAUUGUCAAUUGUGAAAC	85	1513	CUAAUUGUCAAUUGUGAAAC	85	1535	GUUUCACAUUGACAAUUA	512
1531	CCCAGAUUUACGAAAAGG	86	1531	CCCAGAUUUACGAAAAGG	86	1553	CCUUUUCGUAAUUCUGGG	513
1549	GCCGUGUCAUCGUUUCAG	87	1549	GCCGUGUCAUCGUUUCAG	87	1571	CUGGAAACGAUACACGGC	514
1567	GACCCGGCUCUCUACCCAC	88	1567	GACCCGGCUCUCUACCCAC	88	1589	GUGGUAAGAGAGCGGGUC	515
1585	CUGGGCAGCAGACAAUCC	89	1585	CUGGGCAGCAGACAAUCC	89	1607	GGUUUUGUCUGUCCCCAG	516
1603	CUGACUUGUACCGCAUUG	90	1603	CUGACUUGUACCGCAUUG	90	1625	CAUAUGCGGUACAAGUCAG	517
1621	GGUAUCCCUCAACCUACAA	91	1621	GGUAUCCCUCAACCUACAA	91	1643	UUGUAGGUUGAGGGAUACC	518
1639	AUCAAGUGGUUCUGGCACC	92	1639	AUCAAGUGGUUCUGGCACC	92	1661	GGUGCCAGAACCAUUGAU	519
1657	CCUGUAACCAUAAUUAU	93	1657	CCUGUAACCAUAAUUAU	93	1679	AAUGAUUUGGUUACAGGG	520
1675	UCCGAAGCAAGGUGUGACU	94	1675	UCCGAAGCAAGGUGUGACU	94	1697	AGUCACACCUUGCUUGGA	521
1693	UUUUGUCCAAUAAUUAAG	95	1693	UUUUGUCCAAUAAUUAAG	95	1715	CUUCAUUAUUGGAACAAA	522
1711	GAGUCCUUUAUCCUGGAUG	96	1711	GAGUCCUUUAUCCUGGAUG	96	1733	CAUCCAGGAUAAAGACUC	523
1729	GCUGACAGCAACAUUGGAA	97	1729	GCUGACAGCAACAUUGGAA	97	1751	UUCCCAUGUUGCUGUCAG	524
1747	AACAGAAUUGAGAGCAUCA	98	1747	AACAGAAUUGAGAGCAUCA	98	1769	UGAUGCUCUCAAUUCUGU	525
1765	ACUCAGCGCAUGGCAUUA	99	1765	ACUCAGCGCAUGGCAUUA	99	1787	UUAAUUGCCAUUGCGCUGAGU	526
1783	AUAGAAGGAAGAAUUAAGA	100	1783	AUAGAAGGAAGAAUUAAGA	100	1805	UCUUAUUCUUUCCUUCUUAU	527
1801	AUGGCUAGCACCUUGGUUG	101	1801	AUGGCUAGCACCUUGGUUG	101	1823	CAACCAAGGUGCUAGCCAU	528
1819	GUGGCUAGACUCUAGAAUUU	102	1819	GUGGCUAGACUCUAGAAUUU	102	1841	AAAUUCUAGAGUACGCCAC	529

1837	UCUGGAUUCUACAUUUUGCA	103	1837	UCUGGAUUCUACAUUUUGCA	103	1859	UGCAAAUUGUAGAUUCCAGA	530
1855	AUAGCUUCCAAUAAAGUUG	104	1855	AUAGCUUCCAAUAAAGUUG	104	1877	CAACUUAUUGGAAGCUAU	531
1873	GGGACUGUGGGAAGAAACA	105	1873	GGGACUGUGGGAAGAAACA	105	1895	UGUUAUUCUCCACAGUCCC	532
1891	AUAAGCUUUUAUAUCACAG	106	1891	AUAAGCUUUUAUAUCACAG	106	1913	CUGUGAUUAAAGGCUUUAU	533
1909	GAUGGCCAAUUGGGUUC	107	1909	GAUGGCCAAUUGGGUUC	107	1931	GAAACCCAUUUGGCACAU	534
1927	CAUGUUAACUUGGAAAAA	108	1927	CAUGUUAACUUGGAAAAA	108	1949	UUUUUCCAAGUUAACAUG	535
1945	AUGCCGACGGAAGGAGAG	109	1945	AUGCCGACGGAAGGAGAG	109	1967	CCUCUCCUCCGUCGGCAU	536
1963	GACCUGAACUUGUUGCA	110	1963	GACCUGAACUUGUUGCA	110	1985	UGCAAGACAGUUAUCAGGUC	537
1981	ACAGUUAACAAGUUCUUAU	111	1981	ACAGUUAACAAGUUCUUAU	111	2003	AUAAGAACUUGUUAACUGU	538
1999	UACAGAGACGUUACUUGGA	112	1999	UACAGAGACGUUACUUGGA	112	2021	UCCAAGUAAACGUCUCUGA	539
2017	AUUUAACUGCGGACAGUUA	113	2017	AUUUAACUGCGGACAGUUA	113	2039	UAACUGUCCGAGUAAAAU	540
2035	AUAACAGAAACAAGCACU	114	2035	AUAACAGAAACAAGCACU	114	2057	AGUGCAUUGUUCUGUUAUU	541
2053	UACAGUUAUAGCAAGCAAA	115	2053	UACAGUUAUAGCAAGCAAA	115	2075	UUUGCUUGCUAAUACUGUA	542
2071	AAAUGGCCAUCACUAAGG	116	2071	AAAUGGCCAUCACUAAGG	116	2093	CCUUAUGUGAUGGCCAUUUU	543
2089	GAGCACUCCAUACUCUUA	117	2089	GAGCACUCCAUACUCUUA	117	2111	UAAGAGUGAUGGAGUGCUC	544
2107	AAUCUUAACCAUCAUGAUG	118	2107	AAUCUUAACCAUCAUGAUG	118	2129	CAUUAUGAUGGUUAGAAU	545
2125	GUUUCUCCUGCAAGAUUCAG	119	2125	GUUUCUCCUGCAAGAUUCAG	119	2147	CUGAAUCUUGCAGGGAAC	546
2143	GGCACCUAUGCCUGCAGAG	120	2143	GGCACCUAUGCCUGCAGAG	120	2165	CUCUGCAGCAUAGGUGCC	547
2161	GCCAGGAUUGUAUACACAG	121	2161	GCCAGGAUUGUAUACACAG	121	2183	CUGUGUAUACAUUCCUGGC	548
2179	GGGAAAGAAUCCUCCAGA	122	2179	GGGAAAGAAUCCUCCAGA	122	2201	UCUGGAGGAUUAUCCUCCC	549
2197	AAGAAAGAAUUAACAUA	123	2197	AAGAAAGAAUUAACAUA	123	2219	UGAUUGUAUUUUAUUCUU	550
2215	AGAGAUACAGGAAGCACCAU	124	2215	AGAGAUACAGGAAGCACCAU	124	2237	AUGGUGCUUCCUGAUUCU	551
2233	UACCUCCUGCGAAACCUCA	125	2233	UACCUCCUGCGAAACCUCA	125	2255	UGAGGUUUUCGAGGAGGUA	552
2251	AGUGAUCACACAGUGGCCA	126	2251	AGUGAUCACACAGUGGCCA	126	2273	UGGCCACUGUGUGAUCAU	553
2269	AUCAGCAGUCCACCACUU	127	2269	AUCAGCAGUCCACCACUU	127	2291	AAGUGGUGGAACUUGCUGAU	554
2287	UUAGACUGUCAUGCUAAUG	128	2287	UUAGACUGUCAUGCUAAUG	128	2309	CAUUAUGCAUGACAGUCUAA	555
2305	GGUGUCCCGAGCCUCAGA	129	2305	GGUGUCCCGAGCCUCAGA	129	2327	UCUGAGGCUCGGGACACC	556
2323	AUCACUUGGUUUUAAAACA	130	2323	AUCACUUGGUUUUAAAACA	130	2345	UGUUUUUAAAACCAAGUGAU	557
2341	AACCACAAAUACAACAAG	131	2341	AACCACAAAUACAACAAG	131	2363	CUUGUUGUAUUUUUGUGGU	558
2359	GAGCCUGGAUUUUUUUAG	132	2359	GAGCCUGGAUUUUUUUAG	132	2381	CUAAAUAUUUCCAGGCUC	559
2377	GGACCAGGAAGCAGCACGC	133	2377	GGACCAGGAAGCAGCACGC	133	2399	GCUGUGCUGUUCUUGGUCC	560
2395	CUGUUUAUUGAAAGAGUCA	134	2395	CUGUUUAUUGAAAGAGUCA	134	2417	UGACUCUUUCAAUAAACAG	561
2413	ACAGAAGAGGAUGAAGGUG	135	2413	ACAGAAGAGGAUGAAGGUG	135	2435	CACCUUACUCCUCUUCUGU	562
2431	GUCUAUCACUGCAAAAGCCA	136	2431	GUCUAUCACUGCAAAAGCCA	136	2453	UGGCUUUUGCAGUUAUUGU	563
2449	ACCAACCCAGAAAGGUCUCUG	137	2449	ACCAACCCAGAAAGGUCUCUG	137	2471	CAGAGCCUUCUUGGUUGGU	564
2467	GUGGAAAGUUCAGCAUACC	138	2467	GUGGAAAGUUCAGCAUACC	138	2489	GGUAUGCUGAAACUUAUCCAC	565

2485	CUCACUGUUAAGGAACCU	139	2485	CUCACUGUUAAGGAACCU	139	2507	AGGUUCCUUUAACAGUGAG	566
2503	UCGGACAAGUCUUAUCUGG	140	2503	UCGGACAAGUCUUAUCUGG	140	2525	CCAGAUUAGACUUGUCCGA	567
2521	GAGCUGAUCACUCUAACAU	141	2521	GAGCUGAUCACUCUAACAU	141	2543	AUGUUAAGAGUAGUACGUC	568
2539	UGCACCUUGUGGCGCGA	142	2539	UGCACCUUGUGGCGCGA	142	2561	UCGCAGCCACACAGGUGCA	569
2557	ACUCUCUUCUGGCUCCUUA	143	2557	ACUCUCUUCUGGCUCCUUA	143	2579	AUAGGAGCCAGAAAGAGAGU	570
2575	UUAACCCUCCUUAUCCGAA	144	2575	UUAACCCUCCUUAUCCGAA	144	2597	UUCGGAUAAAGGAGGGUUA	571
2593	AAAUGAAAAGGUCUUCUU	145	2593	AAAUGAAAAGGUCUUCUU	145	2615	AAGAAGACCUUUUUAUUUU	572
2611	UCUGAAUUAAGACUGACU	146	2611	UCUGAAUUAAGACUGACU	146	2633	AGUCAGUCUUUAUUUCAGA	573
2629	UACCUAUCAAUUAUUAUGG	147	2629	UACCUAUCAAUUAUUAUGG	147	2651	CCAUUAUUAUUUAUAGGUA	574
2647	GACCCAGAUAGAUUCCUU	148	2647	GACCCAGAUAGAUUCCUU	148	2669	AAGGAACUUAUCUGGGUC	575
2665	UUGGAUGAGCAGUGAGC	149	2665	UUGGAUGAGCAGUGAGC	149	2687	GCUCACACUGCUCAUCCAA	576
2683	CGGCUCUUAUUAUGGCGA	150	2683	CGGCUCUUAUUAUGGCGA	150	2705	UGGCAUCAUAAAGGGAGCCG	577
2701	AGCAAGUGGGAGUUUGCCC	151	2701	AGCAAGUGGGAGUUUGCCC	151	2723	GGGCAAAACUCCACUUGCU	578
2719	CGGGAGAGACUUAACUGG	152	2719	CGGGAGAGACUUAACUGG	152	2741	CCAGUUUAAGUCUCUCOCG	579
2737	GGCAAUACACUUGGAAGAG	153	2737	GGCAAUACACUUGGAAGAG	153	2759	CUCUCCAAAGUGAUUUGCC	580
2755	GGGCUUUUUGGAAAAGUGG	154	2755	GGGCUUUUUGGAAAAGUGG	154	2777	CCACUUUUCGAAAAGCCCC	581
2773	GUUCAAGCAUCAGCAUUUG	155	2773	GUUCAAGCAUCAGCAUUUG	155	2795	CAAAUGCUGAUGCUUGAAC	582
2791	GGCAUUAAGAAUACACCUA	156	2791	GGCAUUAAGAAUACACCUA	156	2813	UAGGUGAUUUUUAUAGCC	583
2809	ACGUGCCGACUGUGGCUG	157	2809	ACGUGCCGACUGUGGCUG	157	2831	CAGCCACAGUCCGCGACGU	584
2827	GUGAAAUGCUGAAAGAGG	158	2827	GUGAAAUGCUGAAAGAGG	158	2849	CCUCUUUCAGCAUUUUCAC	585
2845	GGGGCCACGGCCAGCGAGU	159	2845	GGGGCCACGGCCAGCGAGU	159	2867	ACUCGUGGGCGUGGCCCC	586
2863	UACAAAGCUCUGAUGACUG	160	2863	UACAAAGCUCUGAUGACUG	160	2885	CAGUCAUCAGAGCUUUGUA	587
2881	GAGCUAAAAUUCUUGACCC	161	2881	GAGCUAAAAUUCUUGACCC	161	2903	GGGUCAAAGAUUUUUAAGCUC	588
2899	CACAUUGGCCACCAUCUGA	162	2899	CACAUUGGCCACCAUCUGA	162	2921	UCAGAUUGGUGGCCAAUGUG	589
2917	AACGUGGUUAACCUUGCUG	163	2917	AACGUGGUUAACCUUGCUG	163	2939	CCAGAGGUUAACCAAGUU	590
2935	GGAGCCUGCACCAAGCAAG	164	2935	GGAGCCUGCACCAAGCAAG	164	2957	CUUGCUUGGUGCAGGCCUCC	591
2953	GGAGGCCUCUGAUGGUGA	165	2953	GGAGGCCUCUGAUGGUGA	165	2975	UCACCAUCAGAGGCCUCCUCC	592
2971	AUUGUUUUAUACUGCAAAU	166	2971	AUUGUUUUAUACUGCAAAU	166	2993	AUUUGCAGUAUUAACAACAAU	593
2989	UAUGGAAUUCUCCCAACU	167	2989	UAUGGAAUUCUCCCAACU	167	3011	AGUUGGAGAGAUUUAUCCAU	594
3007	UACCUCAAGAGCAACACGUG	168	3007	UACCUCAAGAGCAACACGUG	168	3029	CACGUUUGCUCUUGAGGUA	595
3025	GACUUAUUUUUUCUACA	169	3025	GACUUAUUUUUUCUACA	169	3047	UGUUGAGAAAAAUAAAGUC	596
3043	AAGGAUGCAGCACUACACA	170	3043	AAGGAUGCAGCACUACACA	170	3065	UGUGUAGUGCUGCAUCCUU	597
3061	AUGGAGCCUAAAGAAAGAA	171	3061	AUGGAGCCUAAAGAAAGAA	171	3083	UUUCUUUUCUUAAGGCCUCCAU	598
3079	AAAUGGAGCCAGGCCUUGG	172	3079	AAAUGGAGCCAGGCCUUGG	172	3101	CCAGGCCUUGGCCUCCAUUUU	599
3097	GAACAAGGCAAGAAACCAA	173	3097	GAACAAGGCAAGAAACCAA	173	3119	UUGGUUUCUUGCCUUGUUC	600
3115	AGACUAGAUAGCGUCACCA	174	3115	AGACUAGAUAGCGUCACCA	174	3137	UGGUGACGCUAUCUAGUCU	601

3133	AGCAGCGAAAGCUUUGCGA	175	3133	AGCAGCGAAAGCUUUGCGA	175	3155	UCGAAAGCUUUCGUGCU	602
3151	AGCUCGGCUUUCAGGAAG	176	3151	AGCUCGGCUUUCAGGAAG	176	3173	CUUCCUGAAAGCCGAGCU	603
3169	GAUAAAAGUCUGAGUGAUG	177	3169	GAUAAAAGUCUGAGUGAUG	177	3191	CAUCACUCAGACUUAUUC	604
3187	GUUGAGGAAGAGGAGGAUU	178	3187	GUUGAGGAAGAGGAGGAUU	178	3209	AUCCUCCUUCUCCUCAAC	605
3205	UCUGACGGUUCUACAAAG	179	3205	UCUGACGGUUCUACAAAG	179	3227	CCUUGAAGAAACCGUCAG	606
3223	GAGCCCAUCACUAGGAAG	180	3223	GAGCCCAUCACUAGGAAG	180	3245	CUUCCAUAGUGAUGGCGUC	607
3241	GAUCUGAUUUUCUACAGUU	181	3241	GAUCUGAUUUUCUACAGUU	181	3263	AACUGAAGAAUACAGUC	608
3259	UUUCAAGUGGCCAGAGGCA	182	3259	UUUCAAGUGGCCAGAGGCA	182	3281	UGCCUCUGGCCACUUGAAA	609
3277	AUGGAGUUCUGUCUCCCA	183	3277	AUGGAGUUCUGUCUCCCA	183	3299	UGGAAGACAGGAACUCCAU	610
3295	AGAAAGUGCAUUAUCGGG	184	3295	AGAAAGUGCAUUAUCGGG	184	3317	CCCAGUAAUGCACUUCU	611
3313	GACCUGGCAGCGAGAAACA	185	3313	GACCUGGCAGCGAGAAACA	185	3335	UGUUUCUGCGUGCCAGGUC	612
3331	AUUCUUUAUCUGAGAAACA	186	3331	AUUCUUUAUCUGAGAAACA	186	3353	UGUUUCAGAUAAAAGAAU	613
3349	AACGUGGUGAAGAUUUGUG	187	3349	AACGUGGUGAAGAUUUGUG	187	3371	CACAAUUCUCCACCACGUU	614
3367	GAUUUUGCCUUGCCCCGGG	188	3367	GAUUUUGCCUUGCCCCGGG	188	3389	CCCGGCAAGGCCAAAAUUC	615
3385	GAUUAUUUAAGAAACCCCG	189	3385	GAUUAUUUAAGAAACCCCG	189	3407	CGGGGUUCUUAUAAAUAUC	616
3403	GAUUAUGUGAGAAAGGAG	190	3403	GAUUAUGUGAGAAAGGAG	190	3425	CUCUUUUCUCACAUAAUC	617
3421	GAUACUGACUUCUCUGA	191	3421	GAUACUGACUUCUCUGA	191	3443	UCAGAGGAAGUCGAGUAUC	618
3439	AAUUGGAUGGCUCCCGAAU	192	3439	AAUUGGAUGGCUCCCGAAU	192	3461	AUUCGGGAGCCAUCCAUUU	619
3457	UCUAUCUUUGACAAAUAUCU	193	3457	UCUAUCUUUGACAAAUAUCU	193	3479	AGAUUUUGUCAAAGAUAGA	620
3475	UACAGCACCAAGAGCGACG	194	3475	UACAGCACCAAGAGCGACG	194	3497	CGUCGCUCUUGGUGCUGUA	621
3493	GUGUGGUCUUAACGGAGUAU	195	3493	GUGUGGUCUUAACGGAGUAU	195	3515	AUACUCCGUAAGACCACAC	622
3511	UUGCUGUGGGAAAUUCUUCU	196	3511	UUGCUGUGGGAAAUUCUUCU	196	3533	AGAAGAUUUCACACAGCAA	623
3529	UCCUUAAGGUGGUCUCCAU	197	3529	UCCUUAAGGUGGUCUCCAU	197	3551	AUGGAGACCCACCUAAGGA	624
3547	UACCCAGGAGUACAAAUGG	198	3547	UACCCAGGAGUACAAAUGG	198	3569	CCAUUUUUAUCCUUGGGUA	625
3565	GAUGAGGACUUUUGCAGUC	199	3565	GAUGAGGACUUUUGCAGUC	199	3587	GACUGCAAAAGUCCUCAUC	626
3583	CGCCUGAGGGAAGGCAUGA	200	3583	CGCCUGAGGGAAGGCAUGA	200	3605	UCAUGCCUUCUCCUAGGCG	627
3601	AGGAUGAGAGCUCCUGAGU	201	3601	AGGAUGAGAGCUCCUGAGU	201	3623	ACUCAGGAGCUUCUACUCCU	628
3619	UACUCUACUCCUGAAAUCU	202	3619	UACUCUACUCCUGAAAUCU	202	3641	AGAUUUCAGGAGUAGAGUA	629
3637	UAUCAGAUCAUGCUGGACU	203	3637	UAUCAGAUCAUGCUGGACU	203	3659	AGUCCAGCAUGAUUCUGUA	630
3655	UGCUGGCACAGAGACCCAA	204	3655	UGCUGGCACAGAGACCCAA	204	3677	UUGGGUCUCUGUGCCAGCA	631
3673	AAAGAAAGGCCAAGAUUUG	205	3673	AAAGAAAGGCCAAGAUUUG	205	3695	CAAAUUCUUGGCCUUCUUCU	632
3691	GCAGAACUUGUGGAAAAAC	206	3691	GCAGAACUUGUGGAAAAAC	206	3713	GUUUUCCACAAAGUUCUGC	633
3709	CUAGGUGAUUUGCUUCAAG	207	3709	CUAGGUGAUUUGCUUCAAG	207	3731	CUUGAAGCAAAUACCCUAG	634
3727	GCAAAUGUACAACAGGAUG	208	3727	GCAAAUGUACAACAGGAUG	208	3749	CAUCCUGUUGUACAUUUGC	635
3745	GGUAAAGACUACAUCCCAA	209	3745	GGUAAAGACUACAUCCCAA	209	3767	UUGGGAUGUAGUCUUAUCC	636
3763	AUCAUUGCCAUACUGACAG	210	3763	AUCAUUGCCAUACUGACAG	210	3785	CUGUCAGUAUGGCAUUGAU	637

3781	GGAAUAGUGGGUUUACAU	211	3781	GGAAUAGUGGGUUUACAU	211	3803	AUGAAACCCACUUAUUC	638
3799	UACUACAUCUCCUCCUUCU	212	3799	UACUACAUCUCCUCCUUCU	212	3821	AGAAGCAGGAGUUGAGUA	639
3817	UCUGAGGACUUCUUAAGG	213	3817	UCUGAGGACUUCUUAAGG	213	3839	CCUUGAAGAGUCCUCAGA	640
3835	GAAAGUUAUUCAGCUCGGA	214	3835	GAAAGUUAUUCAGCUCGGA	214	3857	UCGGAGCUGAAUUAUUC	641
3853	AAGUUAAUUCAGGAAGCU	215	3853	AAGUUAAUUCAGGAAGCU	215	3875	AGCUUCCUGAAUUAACUU	642
3871	UCUGAUGAUGUCAGAUUG	216	3871	UCUGAUGAUGUCAGAUUG	216	3893	CAUAUCUGACAUCAUCAGA	643
3889	GUAAUUGCUUUAAGUUUA	217	3889	GUAAUUGCUUUAAGUUUA	217	3911	UGAACUUGAAAGCAUUUAC	644
3907	AUGAGCCUGGAAAGAAUA	218	3907	AUGAGCCUGGAAAGAAUA	218	3929	UGAUUCUUCAGGCUCAU	645
3925	AAACCUUUGAAGAACUUU	219	3925	AAACCUUUGAAGAACUUU	219	3947	AAAGUUCUUAAGGCUUUU	646
3943	UUAACGAAUCCACCUCCA	220	3943	UUAACGAAUCCACCUCCA	220	3965	UGGAGGUGGCAUUCGGUAA	647
3961	AUGUUUUGAUGACUACCGG	221	3961	AUGUUUUGAUGACUACCGG	221	3983	CCUGGUAGUACAUAACAU	648
3979	GGCGACAGCAGCACUCUGU	222	3979	GGCGACAGCAGCACUCUGU	222	4001	ACAGAGUGCUGCUGCGCC	649
3997	UUGGCCUCUCCCAUGCUGA	223	3997	UUGGCCUCUCCCAUGCUGA	223	4019	UCAGCAUGGGAGAGGCCAA	650
4015	AAGCGCUUCACCUUGGACUG	224	4015	AAGCGCUUCACCUUGGACUG	224	4037	CAGUCCAGGUGAAGCGCUU	651
4033	GACAGCAAAACCCCAAGGCCU	225	4033	GACAGCAAAACCCCAAGGCCU	225	4055	AGGCCUUGGGUUUGCUGUC	652
4051	UCGCUCAAGAUUGACUUGA	226	4051	UCGCUCAAGAUUGACUUGA	226	4073	UCAAGUCAAUUCUUGAGCGA	653
4069	AGAGUAAACCAAGUAAAGUA	227	4069	AGAGUAAACCAAGUAAAGUA	227	4091	UACUUUACUGGUUACUCU	654
4087	AAGGAGUCGGGCGUGUCUG	228	4087	AAGGAGUCGGGCGUGUCUG	228	4109	CAGACAGCCCGACUCUUCU	655
4105	GAUGUCAGCAGGCCAGUUU	229	4105	GAUGUCAGCAGGCCAGUUU	229	4127	AACUGGGCCUGCUGACAU	656
4123	UUCUGCCAUUCCAGCUGUG	230	4123	UUCUGCCAUUCCAGCUGUG	230	4145	CACAGCUGGAUUGGCAGAA	657
4141	GGCACGUCAGCGAAGGCA	231	4141	GGCACGUCAGCGAAGGCA	231	4163	UGCCUUGCUGACGUGCCC	658
4159	AAGCGCAGGUUACCCUACG	232	4159	AAGCGCAGGUUACCCUACG	232	4181	CGUAGGUGAACCCUGCGCUU	659
4177	GACCACGUCAGCUGGAAA	233	4177	GACCACGUCAGCUGGAAA	233	4199	UUUCCAGCUGACGUGGUC	660
4195	AGGAAAUUCGCGUGCUGCU	234	4195	AGGAAAUUCGCGUGCUGCU	234	4217	AGCAGCAGCGAUUUUCCU	661
4213	UCCCGCCCCCAGACUACA	235	4213	UCCCGCCCCCAGACUACA	235	4235	UGUAGUCUGGGGGCGGGGA	662
4231	AACUCGGUGGUCUCCUACU	236	4231	AACUCGGUGGUCUCCUACU	236	4253	AGUACAGGACCACCGAGUU	663
4249	UCCACCCACCCAUUCUAGA	237	4249	UCCACCCACCCAUUCUAGA	237	4271	UCUAGAUUGGUGGGGUGGA	664
4267	AGUUUGACACGAAGCCUUA	238	4267	AGUUUGACACGAAGCCUUA	238	4289	UAAAGCUUCUGUCUCAAACU	665
4285	AUUUCUAGAAAGCACAUUG	239	4285	AUUUCUAGAAAGCACAUUG	239	4307	CACAUUGUCUUCUAGAAAU	666
4303	GUUUUUAUACCCCGAGGAA	240	4303	GUUUUUAUACCCCGAGGAA	240	4325	UUCUUGGGGUGUUAUUAUAC	667
4321	AACUAGCUUUUUGCCAGUUA	241	4321	AACUAGCUUUUUGCCAGUUA	241	4343	AUACUGGCAAAAGCUAGUU	668
4339	UUUAGCAUAUAUAAGUUUA	242	4339	UUUAGCAUAUAUAAGUUUA	242	4361	UAAACUUAUAUAUGCAUAA	669
4357	ACACCUUUAUCUUCUCCAU	243	4357	ACACCUUUAUCUUCUCCAU	243	4379	CAUGGAAAGAUAAAGGUGU	670
4375	GGGAGCCAGCUGCUUUUUG	244	4375	GGGAGCCAGCUGCUUUUUG	244	4397	CAAAAGCAGCUGGCUCCC	671
4393	GUGAUUUUUUUAUAUAGUGC	245	4393	GUGAUUUUUUUAUAUAGUGC	245	4415	GCACUUAUUAUUAUUAUAC	672
4411	CUUUUUUUUUUUGACUAAAC	246	4411	CUUUUUUUUUUUGACUAAAC	246	4433	GUUAGUCAAAAAAAG	673

4429	CAAGAAUGUAACUCCAGAU	247	4429	CAAGAAUGUAACUCCAGAU	247	4451	AUCUGGAGUUACAUUCUUG	674
4447	UAGAGAAUAGUGACAAGU	248	4447	UAGAGAAUAGUGACAAGU	248	4469	ACUUGUCACUAAUUCUCUA	675
4465	UGAAGAACACUACUGCUAA	249	4465	UGAAGAACACUACUGCUAA	249	4487	UUAGCAGUAGUUUCUUA	676
4483	AAUCCUCAUGUUACUCAGU	250	4483	AAUCCUCAUGUUACUCAGU	250	4505	ACUGAGUAACAUAGGAGUU	677
4501	UGUUAGAGAAAUCCUCCU	251	4501	UGUUAGAGAAAUCCUCCU	251	4523	AGGAAGAUUUCUCUAACA	678
4519	UAAACCCAAUGACUCCCU	252	4519	UAAACCCAAUGACUCCCU	252	4541	AGGAAGUACAUUGGUUUA	679
4537	UGCUCACACCCCGCCACC	253	4537	UGCUCACACCCCGCCACC	253	4559	GGUGCGGGGUUGGAGCA	680
4555	CUCAGGCACGCAGGACCA	254	4555	CUCAGGCACGCAGGACCA	254	4577	UGGUCCUGCGUGCCUGAG	681
4573	AGUUUGAUUGAGGAGCUGC	255	4573	AGUUUGAUUGAGGAGCUGC	255	4595	GCAGCUCUCAAUCAAACU	682
4591	CACUGAUCACCCAAUGCAU	256	4591	CACUGAUCACCCAAUGCAU	256	4613	AUGCAUUGGUUGAUACAGUG	683
4609	UCACGUACCCACUGGGCC	257	4609	UCACGUACCCACUGGGCC	257	4631	GGCCAGUGGGGUACGUGA	684
4627	CAGCCUUGCAGCCCAAAAC	258	4627	CAGCCUUGCAGCCCAAAAC	258	4649	GUUUUGGCGUCAGGGCUG	685
4645	CCAGGGCAACAAGCCCGU	259	4645	CCAGGGCAACAAGCCCGU	259	4667	ACGGCUUGUUGCCCGGG	686
4663	UUAGCCCCAGGGUACACU	260	4663	UUAGCCCCAGGGUACACU	260	4685	AGUGAUCCCGUGGGCUAA	687
4681	UGGCUGGCCUAGCAACAU	261	4681	UGGCUGGCCUAGCAACAU	261	4703	AUGUUGCUCAGGGCCAGCCA	688
4699	UCUCGGGAGUCCUAGCA	262	4699	UCUCGGGAGUCCUAGCA	262	4721	UGCUAGAGGACUCCCGAGA	689
4717	AGGCCUAGACAUUGAGG	263	4717	AGGCCUAGACAUUGAGG	263	4739	CCUCACAUUGCUUAGGCCU	690
4735	GAGGAAAGGAAAAAAGC	264	4735	GAGGAAAGGAAAAAAGC	264	4757	GCUUUUUUCCUUUUCUC	691
4753	CAAAAAGCAAGGAGAAAA	265	4753	CAAAAAGCAAGGAGAAAA	265	4775	UUUUCUCCCUUGCUUUUUG	692
4771	AGAGAACCGGGAGAAAGC	266	4771	AGAGAACCGGGAGAAAGC	266	4793	GCCUUCUCCCGGUUUCUCU	693
4789	CAUGAGAAAGAAUUUAGAG	267	4789	CAUGAGAAAGAAUUUAGAG	267	4811	UCUCAAAUUCUUUUCUACUG	694
4807	ACGACCAUUGGGCACGG	268	4807	ACGACCAUUGGGCACGG	268	4829	CCGUGCCACAUUGGUGCGU	695
4825	GAGGGGACGGGGCUCAGC	269	4825	GAGGGGACGGGGCUCAGC	269	4847	GCUGAGCCCGUCCCCUC	696
4843	CAAUGCCAUUUCAGUGGCU	270	4843	CAAUGCCAUUUCAGUGGCU	270	4865	AGCCACUGAAUUGGCAUUG	697
4861	UUCGCAGCUCUGACCCUUC	271	4861	UUCGCAGCUCUGACCCUUC	271	4883	GAAGGUCACAGCUGGGAA	698
4879	CUACAUUUGAGGGCCACGC	272	4879	CUACAUUUGAGGGCCACGC	272	4901	GCUGGGCCCUCAAUUGUAG	699
4897	CCAGGAGCAGAUUGGACAGC	273	4897	CCAGGAGCAGAUUGGACAGC	273	4919	GCUGUCCAUUGCUCUCCUGG	700
4915	CGAUGAGGGGACAUUUUCU	274	4915	CGAUGAGGGGACAUUUUCU	274	4937	AGAAAUGUCCCUCAUCG	701
4933	UGGAUUCUGGGAGGCAAGA	275	4933	UGGAUUCUGGGAGGCAAGA	275	4955	UCUUGCCUCCCGAUAUCCA	702
4951	AAAAGGACAAUUAUCUUUU	276	4951	AAAAGGACAAUUAUCUUUU	276	4973	AAAAGAUUUUGUCCUJU	703
4969	UUUGGAACUAAAGCAAAU	277	4969	UUUGGAACUAAAGCAAAU	277	4991	AAUUGCUUUAUAGUCCAAA	704
4987	UUUAGACCUUUUACCUAUGG	278	4987	UUUAGACCUUUUACCUAUGG	278	5009	CCAUAGGUAAAAGGUCUAAA	705
5005	GAAGUGGUUCUAUGUCCAU	279	5005	GAAGUGGUUCUAUGUCCAU	279	5027	AUGGACAUAGAACCACUUC	706
5023	UUCUCAUUCGUGGCAUGUU	280	5023	UUCUCAUUCGUGGCAUGUU	280	5045	AACAUGCCACGAUAGAGAA	707
5041	UUUGAUUUGUAGCACUGAG	281	5041	UUUGAUUUGUAGCACUGAG	281	5063	CUCAGUGCUACAAUCAA	708
5059	GGGUGGCACUCAAUCUCUGA	282	5059	GGGUGGCACUCAAUCUCUGA	282	5081	UCAGAGUUUGAGUCCACCC	709

5077	AGCCCAUACUUUUGGCUC	283	5077	AGCCCAUACUUUUGGCUC	283	5099	GGAGCCAAAAGUAGGGCU	710
5095	CUCUAGUAGAUGCACUGA	284	5095	CUCUAGUAGAUGCACUGA	284	5117	UCAGUGCAUCUUACUAGAG	711
5113	AAAACUUAGCCAGAGUUAG	285	5113	AAAACUUAGCCAGAGUUAG	285	5135	CUAACUCUGGCUAAGUUUU	712
5131	GGUUGUCUCCAGGCCAUGA	286	5131	GGUUGUCUCCAGGCCAUGA	286	5153	UCAUGGCCUGGAGACAACC	713
5149	AUGGCCUUAACACUGAAAAU	287	5149	AUGGCCUUAACACUGAAAAU	287	5171	AUUUUCAGUGUAGGCCCAU	714
5167	UGUCACAUUCUAAUUUUGG	288	5167	UGUCACAUUCUAAUUUUGG	288	5189	CCCAAAUAGAAUGUGACA	715
5185	GUUUAAUUAUAGUCCAG	289	5185	GUUUAAUUAUAGUCCAG	289	5207	CUGGACUUAUUAUUAUAC	716
5203	GACACUUAAUCUAAUUUCU	290	5203	GACACUUAAUCUAAUUUCU	290	5225	AGAAUUGAGUUAAUGUGC	717
5221	UUGGUAAUUAUCUGUUUUG	291	5221	UUGGUAAUUAUCUGUUUUG	291	5243	CAAAACAGAAUAAUACCAA	718
5239	GCACAGUUAUUGUGAAAG	292	5239	GCACAGUUAUUGUGAAAG	292	5261	CUUUCACAACUAAUCUGUC	719
5257	GAAGCUGAGAAAGAAUGAA	293	5257	GAAGCUGAGAAAGAAUGAA	293	5279	UUCAUUCUUCUCAGCUUUC	720
5275	AAUUGCAGUCCUGAGGAGA	294	5275	AAUUGCAGUCCUGAGGAGA	294	5297	UCUCCUCAGGACUGCAUUU	721
5293	AGUUUUCUCCAUUAUCAA	295	5293	AGUUUUCUCCAUUAUCAA	295	5315	UUUUGAUUGGAGAAACU	722
5311	ACGAGGCUAGUAGGAGGAA	296	5311	ACGAGGCUAGUAGGAGGAA	296	5333	UUCUCCUACAGCCCUUGU	723
5329	AAAAGGUCAAAUAGGUCAA	297	5329	AAAAGGUCAAAUAGGUCAA	297	5351	UUGACCUUAUUGACCUUUU	724
5347	AGGGAAGACCCGUCUCUA	298	5347	AGGGAAGACCCGUCUCUA	298	5369	UAGAGACGGGGUCUCCCU	725
5365	AUACCAACCAACCAAUUC	299	5365	AUACCAACCAACCAAUUC	299	5387	GAUUUGUUUGGUUGGUUU	726
5383	CACCAACACAGUUGGGACC	300	5383	CACCAACACAGUUGGGACC	300	5405	GGUCCCAACUGUUGUUGG	727
5401	CCAAAACACAGGAAGUCAG	301	5401	CCAAAACACAGGAAGUCAG	301	5423	CUGACUUCUUGUUGUUGG	728
5419	GUCACGUUUCCUUUUCAUU	302	5419	GUCACGUUUCCUUUUCAUU	302	5441	AAUGAAAAGGAAACGUGAC	729
5437	UUAAUGGGGAUUCACAUU	303	5437	UUAAUGGGGAUUCACAUU	303	5459	AUAGUGGAUUCUCCAUUAA	730
5455	UCUCACACUAAUCUGAAAG	304	5455	UCUCACACUAAUCUGAAAG	304	5477	CUUUCAGAUUAGUGUGAGA	731
5473	GAUGUGGGAAGAGCAUUAG	305	5473	GAUGUGGGAAGAGCAUUAG	305	5495	CUAAUGCUCUCCACAUCC	732
5491	GCUGGCGCAUUAUAAAGCAC	306	5491	GCUGGCGCAUUAUAAAGCAC	306	5513	GUGCUUAAUUAUGCGCCAGC	733
5509	CUUUAAGCUCUUGAGUAA	307	5509	CUUUAAGCUCUUGAGUAA	307	5531	UUACUCAAGGAGCUUAAAG	734
5527	AAAAGGUGGUUUGUAAUUU	308	5527	AAAAGGUGGUUUGUAAUUU	308	5549	AAAUUACAUACCAACCUUUU	735
5545	UAUGCAAGGUUUUUCUCCA	309	5545	UAUGCAAGGUUUUUCUCCA	309	5567	UGGAGAAUACCUUGCAUA	736
5563	AGUUUGGACUCAGGAUUAU	310	5563	AGUUUGGACUCAGGAUUAU	310	5585	AAUAUCCUGAGUCCCAACU	737
5581	UAGUUAAUGAGCCAUACU	311	5581	UAGUUAAUGAGCCAUACU	311	5603	AGUGAUGGCUCAUUUAACUA	738
5599	UAGAAAGAAAGCCCAUUUU	312	5599	UAGAAAGAAAGCCCAUUUU	312	5621	AAAAUGGGCUUUUCUUCUA	739
5617	UAAACUGCUUUUGAAACUUG	313	5617	UAAACUGCUUUUGAAACUUG	313	5639	CAAGUUUCAAAGCAGUUGA	740
5635	GCCUGGGUUGAGAGCAUGA	314	5635	GCCUGGGUUGAGAGCAUGA	314	5657	UCAUGCUCAGACCCAGGC	741
5653	AUGGGAUAGGGAGACAGG	315	5653	AUGGGAUAGGGAGACAGG	315	5675	CCUGUCUCCUUAUUCUCCAU	742
5671	GGUAGGAAGGGCGCCUAC	316	5671	GGUAGGAAGGGCGCCUAC	316	5693	GUAGGCGCCUUAUUCUAC	743
5689	CUCUUCAGGGUUAUAAAGAU	317	5689	CUCUUCAGGGUUAUAAAGAU	317	5711	AUCUUUAGACCCUAGAGAG	744
5707	UCAAGUGGGCCUUGGAUCG	318	5707	UCAAGUGGGCCUUGGAUCG	318	5729	CGAUCCAAAGGCCCAUUGA	745

5725	GCUAAGCUGGCUUCUUG	319	5725	GCUAAGCUGGCUUCUUG	319	5747	CAACAGAGCCAGCUUAGC	746
5743	GAUGCUAUUUAUGCAAGUU	320	5743	GAUGCUAUUUAUGCAAGUU	320	5765	AACUUGCAUAAUAGCAUC	747
5761	UAGGGUCUAUGUAUUUAGG	321	5761	UAGGGUCUAUGUAUUUAGG	321	5783	CCUAAUACAUAGACCCUA	748
5779	GAUGCGCCUACUCUUCAGG	322	5779	GAUGCGCCUACUCUUCAGG	322	5801	CCUGAAGAGUAGGCGCAUC	749
5797	GGUCUAAAGAUCAAGUGGG	323	5797	GGUCUAAAGAUCAAGUGGG	323	5819	CCCACUUGAUUUAUAGACC	750
5815	GCCUUGGAUCGCUAAGCUG	324	5815	GCCUUGGAUCGCUAAGCUG	324	5837	CAGCUUAGCGAUCCAAGGC	751
5833	GGCUCUGUUUGAUGCUAUU	325	5833	GGCUCUGUUUGAUGCUAUU	325	5855	AAUAGCAUCAAACAGAGCC	752
5851	UUAUGCAAGUUAAGGUCUA	326	5851	UUAUGCAAGUUAAGGUCUA	326	5873	UAGACCCUAAUUGCAUAA	753
5869	AUGUAUUUAGGAUGUCUG	327	5869	AUGUAUUUAGGAUGUCUG	327	5891	GCAGACAUCCUAAAUACAU	754
5887	CACCUUCUGCAGCCAGUCA	328	5887	CACCUUCUGCAGCCAGUCA	328	5909	UGACUGGCGUGCAGAAAGGUG	755
5905	AGAAGCUGGAGGCAACA	329	5905	AGAAGCUGGAGGCAACA	329	5927	UGUUGCCUCUCCAGCUUCU	756
5923	AGUGGAUUGCUGCUUCUUG	330	5923	AGUGGAUUGCUGCUUCUUG	330	5945	CAAGAAGCAGCAAUCCACU	757
5941	GGGAGAAAGUAUGCUUC	331	5941	GGGAGAAAGUAUGCUUC	331	5963	GAAAGCAUACUCUUCUCCCC	758
5959	CCUUUAUCCAUUAUUUU	332	5959	CCUUUAUCCAUUAUUUU	332	5981	AAAUUACAUGGAUAAAGG	759
5977	UAACUGUAACCCUGAGCU	333	5977	UAACUGUAACCCUGAGCU	333	5999	AGCUCAGGUUCUACAGUUA	760
5995	UCUAAGUAACCGAAGAU	334	5995	UCUAAGUAACCGAAGAU	334	6017	CAUUCUUCGGUUAUUAUA	761
6013	GAUGCCUCUGUUCUUAUG	335	6013	GAUGCCUCUGUUCUUAUG	335	6035	CAUAAGAACAGAGGCAUAC	762
6031	GUGCCACAUCUUGUUAUA	336	6031	GUGCCACAUCUUGUUAUA	336	6053	UUAACAAGGAUGUGGCAC	763
6049	AAGGCUCUCUGUAUGAAGA	337	6049	AAGGCUCUCUGUAUGAAGA	337	6071	UCUUAUACAGAGAGCCUU	764
6067	AGAUGGACCGUCAUCAGC	338	6067	AGAUGGACCGUCAUCAGC	338	6089	GCUGAUGACGGUCCCAUCU	765
6085	CACAUUCCUAGUGAGCCU	339	6085	CACAUUCCUAGUGAGCCU	339	6107	AGCUCACUAGGGAUUG	766
6103	UACUUGGCUCCUGGCAGCG	340	6103	UACUUGGCUCCUGGCAGCG	340	6125	CCGCUGCCAGGAGCCAGUA	767
6121	GCUUUUUGGGAAGACUCAC	341	6121	GCUUUUUGGGAAGACUCAC	341	6143	GUGAGUCUCCACAAAAGC	768
6139	CUAGCCAGAAGAGAGGAGU	342	6139	CUAGCCAGAAGAGAGGAGU	342	6161	ACUCCUCUCUUCUGGCUAG	769
6157	UGGACAGUCUUCUCCACC	343	6157	UGGACAGUCUUCUCCACC	343	6179	GGUGGAGAGGACUGUCCCA	770
6175	CAAGAUUAAUCCAAACA	344	6175	CAAGAUUAAUCCAAACA	344	6197	UGUUUGGAUUUAGAUUUG	771
6193	AAAAGCAGGCUAGAGCCAG	345	6193	AAAAGCAGGCUAGAGCCAG	345	6215	CUGGCUCUAGCCUGCUUUU	772
6211	GAAGAGAGGACAAUUCUU	346	6211	GAAGAGAGGACAAUUCUU	346	6233	AAAGAUUUUGUCCUCUUC	773
6229	UGUUUUCUUCUUCUUAAC	347	6229	UGUUUUCUUCUUCUUAAC	347	6251	GUAAAGAGAGGGAACAACA	774
6247	CACAUACGCAACCCACUUG	348	6247	CACAUACGCAACCCACUUG	348	6269	CAGGUGGUUUGCGUAUGUG	775
6265	GUGACAGCUGGCAAUUUUA	349	6265	GUGACAGCUGGCAAUUUUA	349	6287	UAAAAUUGCCAGCUGUCAC	776
6283	AUAAUACAGGUAAUUGGAA	350	6283	AUAAUACAGGUAAUUGGAA	350	6305	UUCAGUUUACCUUUAUUAU	777
6301	AGGAGGUUAAACUCAGAAA	351	6301	AGGAGGUUAAACUCAGAAA	351	6323	UUUCUGAGUUUAACCUCCU	778
6319	AAAAAGACCCUCAGUCAAA	352	6319	AAAAAGACCCUCAGUCAAA	352	6341	UUGACUGAGGCUUCUUCUUU	779
6337	AUUCUCUACUUUUUUUUUU	353	6337	AUUCUCUACUUUUUUUUUU	353	6359	AAAAAAAAGUAGAGAAU	780
6355	UUUUUUUCCAAAUACAGUA	354	6355	UUUUUUUCCAAAUACAGUA	354	6377	UAUCUGAUUUUGGAAAAAAA	781

6373	AAUAGCCCGAGCAAAUAGUG	355	6373	AAUAGCCCGAGCAAAUAGUG	355	6395	CACUAAUUUGCGGCUAUU	782
6391	GAUAAACAAUAAACCUUA	356	6391	GAUAAACAAUAAACCUUA	356	6413	UAAGGUUUUAUUUGUUAUC	783
6409	AGCUGUUCAGUCUUGAUU	357	6409	AGCUGUUCAGUCUUGAUU	357	6431	AAUCAAGACAUAGAACAGCU	784
6427	UUCAAUAAUUAUUUCUUA	358	6427	UUCAAUAAUUAUUUCUUA	358	6449	UUAAGAAUUAUUUAUUGAA	785
6445	AUCAUUAAGAGACCAUAU	359	6445	AUCAUUAAGAGACCAUAU	359	6467	AUUAUGGUCUCUUAUUGAU	786
6463	UAAUACUCCUUUUAAGA	360	6463	UAAUACUCCUUUUAAGA	360	6485	UCUUGAAAAAGGAGUUAUUA	787
6481	AGAAAAGCAAAACCAUAG	361	6481	AGAAAAGCAAAACCAUAG	361	6503	CUAAUGGUUUUGCUUUUCU	788
6499	GAUUUGUUAUCAGCUCU	362	6499	GAUUUGUUAUCAGCUCU	362	6521	AGGAGCUGAGUAACAAUUC	789
6517	UUCAAACUCAGGUUUGUAG	363	6517	UUCAAACUCAGGUUUGUAG	363	6539	CUACAAACCGAGGUUUGAA	790
6535	GCAUACAUAGAUCCA	364	6535	GCAUACAUAGAUCCA	364	6557	UGGAUGGACUCAUGUAUGC	791
6553	AUCAGUCAAAAGAUUGUUC	365	6553	AUCAGUCAAAAGAUUGUUC	365	6575	GAACCAUUCUUUGACUGAU	792
6571	CCAUCUGGAGUCUUAUUGU	366	6571	CCAUCUGGAGUCUUAUUGU	366	6593	ACAUUAAGACUCCAGAUUG	793
6589	UAGAAAGAAAAUGGAGAC	367	6589	UAGAAAGAAAAUGGAGAC	367	6611	GUCCCAUUUUUCUUUCUA	794
6607	CUUGUAAUUAUGAGCUAGU	368	6607	CUUGUAAUUAUGAGCUAGU	368	6629	ACUAGCUCAUUAUUAACAAG	795
6625	UUCAAAGUGCUUUGUUAU	369	6625	UUCAAAGUGCUUUGUUAU	369	6647	AUGAACAAAGCACUUAUUA	796
6643	UUAUUUAAGCACUGAAAAU	370	6643	UUAUUUAAGCACUGAAAAU	370	6665	AUUUUCAGUGCUAUUUUAA	797
6661	UUGAAACAUGAUUUAACUG	371	6661	UUGAAACAUGAUUUAACUG	371	6683	CAGUUAUUUAUUGUUAUCAA	798
6679	GAUAAUUAUCCAUAUUAU	372	6679	GAUAAUUAUCCAUAUUAU	372	6701	AAUUGAUUGGAAUUAUUAUC	799
6697	UGCCAUUUAUGACAAAAU	373	6697	UGCCAUUUAUGACAAAAU	373	6719	AUUUUUGUCAUAAAUGGCA	800
6715	UGGUUGGCACUAAACAAAGA	374	6715	UGGUUGGCACUAAACAAAGA	374	6737	UCUUUGUUAUGUGCCAAACCA	801
6733	AACGAGCACUCCUUUCAG	375	6733	AACGAGCACUCCUUUCAG	375	6755	CUGAAAAGGAGUGUCUCGUU	802
6751	GAGUUUCUGAGAUAAUUGUA	376	6751	GAGUUUCUGAGAUAAUUGUA	376	6773	UACAUUAUCUCAGAAAACUC	803
6769	ACGUGGAACAGUCUGGGUG	377	6769	ACGUGGAACAGUCUGGGUG	377	6791	CACCCAGACUUGUCCACGU	804
6787	GGAAUGGGGCGUAAACCAU	378	6787	GGAAUGGGGCGUAAACCAU	378	6809	AUGGUUUCAGCCCCAUUCC	805
6805	UGUGCAAGUCUGUGUCUUG	379	6805	UGUGCAAGUCUGUGUCUUG	379	6827	CAAGACACAGACUUGCACA	806
6823	GUCAGUCCAGAAAGUGACA	380	6823	GUCAGUCCAGAAAGUGACA	380	6845	UGUCACUUCUUGGACUGAC	807
6841	ACCGAGAUUUAAUUUUAAG	381	6841	ACCGAGAUUUAAUUUUAAG	381	6863	CUAAAAUUAAACAUUCUGGU	808
6859	GGGACCCGUGCCUUGUUUC	382	6859	GGGACCCGUGCCUUGUUUC	382	6881	GAACAAAGGCACGGGUCCC	809
6877	CCUAGCCCAAGAAUUGCA	383	6877	CCUAGCCCAAGAAUUGCA	383	6899	UGCAUUCUUGUGGGCUAGG	810
6895	AAACAUCAAACAGAUACUC	384	6895	AAACAUCAAACAGAUACUC	384	6917	GAGUAUCUGUUUGAUGUUU	811
6913	CGCUAGCCUUAUUAAAUU	385	6913	CGCUAGCCUUAUUAAAUU	385	6935	AAUUUAAUUGAGGCUAGCG	812
6931	UGAUUAAAGGAGGUGCA	386	6931	UGAUUAAAGGAGGUGCA	386	6953	UGCACUCCUCCUUUUAUUA	813
6949	AUCUUUGGCCGACAGUGGU	387	6949	AUCUUUGGCCGACAGUGGU	387	6971	ACCACUGUCGGCCAAAGAU	814
6967	UGUAACUGUGUGUGUGUGU	388	6967	UGUAACUGUGUGUGUGUGU	388	6989	ACACACACACACAGUUAUA	815
6985	UGUGUGUGUGUGUGUGUGU	389	6985	UGUGUGUGUGUGUGUGUGU	389	7007	ACACACACACACACACA	816
7003	UGUGUGUGUGUGUGUGUGG	390	7003	UGUGUGUGUGUGUGUGUGG	390	7025	CCACACCCACACACACACA	817

7021	GGUGUAUGUGUUUUUGUG	391	7021	GGUGUAUGUGUUUUUGUG	391	7043	CACAAAACACACAUACACC	818
7039	GCAUAACUUAUUUAGGAAA	392	7039	GCAUAACUUAUUUAGGAAA	392	7061	UUUCCUUUAAAUAUUUAGC	819
7057	ACUGGAAUUUUUAAAGUAC	393	7057	ACUGGAAUUUUUAAAGUAC	393	7079	GUAACUUUUAAAUAUUUCCAGU	820
7075	CUUUUAUACAAACCAAGAA	394	7075	CUUUUAUACAAACCAAGAA	394	7097	UUCUUGGUUUUUAUAAAAG	821
7093	AUAUAUGCUACAGAUUAA	395	7093	AUAUAUGCUACAGAUUAA	395	7115	UUUAUCUGUAUUAUUAU	822
7111	AGACAGACAUUUUUGGUC	396	7111	AGACAGACAUUUUUGGUC	396	7133	GACCAACCAUGUCUGUCU	823
7129	CCUAUAUUUCUAGUCAUGA	397	7129	CCUAUAUUUCUAGUCAUGA	397	7151	UCAUGACUAGAAAUAUAGG	824
7147	AUGAAUGUAUUUUGUAUAC	398	7147	AUGAAUGUAUUUUGUAUAC	398	7169	GUAUAACAAUAUUAUUAU	825
7165	CCAUCUUAUUAUUAUUAU	399	7165	CCAUCUUAUUAUUAUUAU	399	7187	GUAUAUUAUUAUUAUUAU	826
7183	CUUAAAAUAUUUCUUAU	400	7183	CUUAAAAUAUUUCUUAU	400	7205	AUUAAGAAUAUUUUUAAG	827
7201	UUGGGAUUUUAUUAUCGUAC	401	7201	UUGGGAUUUUAUUAUCGUAC	401	7223	GUACGAUUAUUAUUAUUAU	828
7219	CCAACUUAUUUAUUAUUAU	402	7219	CCAACUUAUUUAUUAUUAU	402	7241	AGUUUAUUAUUAUUAUUAU	829
7237	UUGGCAACUGCUUUUAUUGU	403	7237	UUGGCAACUGCUUUUAUUGU	403	7259	ACAUAAAAGCAGUUUCCAA	830
7255	UUCUGUCUGCUUCCAUAAA	404	7255	UUCUGUCUGCUUCCAUAAA	404	7277	UUUAUGGAAAGGAGACAGAA	831
7273	AUUUUUAUUAUUAUUAUUAU	405	7273	AUUUUUAUUAUUAUUAUUAU	405	7295	AAUUAUAUUUUUUAUUAU	832
7291	UCAACAAAGAAAGCUCU	406	7291	UCAACAAAGAAAGCUCU	406	7313	AGAGCUUUUUUUUUUUUAU	833
7309	UUUUUUUCCUUAUUAUUAU	407	7309	UUUUUUUCCUUAUUAUUAU	407	7331	UUUAUUUUUAGGAAAAAAUA	834
7327	ACUCAAAUUUAUUAUUAUUAU	408	7327	ACUCAAAUUUAUUAUUAUUAU	408	7349	AACAAGGAUUAUUAUUAUUAU	835
7345	UAGAAACUUUUAUUAUUAU	409	7345	UAGAAACUUUUAUUAUUAU	409	7367	UAAUUUUUUUUUUUUUUUAU	836
7363	AAGAAACUUUUAUUAUUAU	410	7363	AAGAAACUUUUAUUAUUAU	410	7385	CCAUUUCAAAGUUUUUUUUUAU	837
7381	GUUCUAAAAUUUUAUUAUUAU	411	7381	GUUCUAAAAUUUUAUUAUUAU	411	7403	UUUAAGCAUUUUUUUUUUUAU	838
7399	AUAUUUAUUAUUAUUAUUAU	412	7399	AUAUUUAUUAUUAUUAUUAU	412	7421	AGUUUUCCAUUAUUAUUAUUAU	839
7417	UAAUUGUUUAUUAUUAUUAU	413	7417	UAAUUGUUUAUUAUUAUUAU	413	7439	UCAGCUAAACUUAUUAUUAUUAU	840
7435	AUUUAUUGGGUUUUUUAUUAU	414	7435	AUUUAUUGGGUUUUUUAUUAU	414	7457	UUCGAAAAACCCAUUAUUAUUAU	841
7453	ACCUUUACUUUUUUUUUUUAUUAU	415	7453	ACCUUUACUUUUUUUUUUUAUUAU	415	7475	CAACAAAAAGUGAAAGGU	842
7471	GUUUUAACCUUAUUAUUAUUAU	416	7471	GUUUUAACCUUAUUAUUAUUAU	416	7493	GUUGUGAAUAUUAUUAUUAUUAU	843
7489	CUGUGUAAAUUUGCCAAUUAU	417	7489	CUGUGUAAAUUUGCCAAUUAU	417	7511	UUUAUUGGCAUUUAUUAUUAUUAU	844
7507	AUUCUGUCCAUUUAUUAUUAUUAU	418	7507	AUUCUGUCCAUUUAUUAUUAUUAU	418	7529	CAUUUUAUUAUUAUUAUUAUUAU	845
7525	GCAAAUUAUUAUUAUUAUUAUUAU	419	7525	GCAAAUUAUUAUUAUUAUUAUUAU	419	7547	UCUACACUGGAUUAUUAUUAUUAU	846
7543	AUAUAUUUUAUUAUUAUUAUUAU	420	7543	AUAUAUUUUAUUAUUAUUAUUAU	420	7565	GGGUGAUGGUAUUAUUAUUAUUAU	847
7561	CUAUGGAUUAUUAUUAUUAUUAU	421	7561	CUAUGGAUUAUUAUUAUUAUUAU	421	7583	AACUAGCCAAUUAUUAUUAUUAUUAU	848
7579	UUUGCCUUUUUAUUAUUAUUAUUAU	422	7579	UUUGCCUUUUUAUUAUUAUUAUUAU	422	7601	UUUGCUUUUAUUAUUAUUAUUAUUAU	849
7597	AUUCAUUUAUUAUUAUUAUUAUUAU	423	7597	AUUCAUUUAUUAUUAUUAUUAUUAU	423	7619	CAUUCAGGCUUUAUUAUUAUUAUUAU	850
7615	GUCUGCCUUAUUAUUAUUAUUAUUAU	424	7615	GUCUGCCUUAUUAUUAUUAUUAUUAU	424	7637	AGAGAAUAUUAUUAUUAUUAUUAUUAU	851
7633	UGCUCUUUUAUUAUUAUUAUUAUUAU	425	7633	UGCUCUUUUAUUAUUAUUAUUAUUAU	425	7655	AAGGAGAAUAUUAUUAUUAUUAUUAUUAU	852
7651	UUGAACCCCGUUUAUUAUUAUUAUUAU	426	7651	UUGAACCCCGUUUAUUAUUAUUAUUAU	426	7673	GAUGUUUUUAUUAUUAUUAUUAUUAUUAU	853

7662	AAAACAUCUUGGACACUC	427	7662	AAAACAUCUUGGACACUC	427	7684	GAGUGCCACAGGAUUGUUU	854
------	--------------------	-----	------	--------------------	-----	------	---------------------	-----

VEGFR2 gi|11321596|ref|NM_002253.1

Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPos	Lower seq	Seq ID
1	ACUGAGUCCCGGACCCCG	855	1	ACUGAGUCCCGGACCCCG	855	23	CGGGUCCCGGACUCAGU	1179
19	GGGAGAGCGGUCAGUGU	856	19	GGGAGAGCGGUCAGUGU	856	41	ACACACUGACCGCUCUCCC	1180
37	UGGUCGUCGCUUUCUCU	857	37	UGGUCGUCGCUUUCUCU	857	59	AGAGAAACGCAGCGACCA	1181
55	UGCCUGCGCGGCAUCAC	858	55	UGCCUGCGCGGCAUCAC	858	77	GUGAUGCCCGCGCAGGCA	1182
73	CUUGCGCGCGCAGAAAGU	859	73	CUUGCGCGCGCAGAAAGU	859	95	ACUUCUCGCGCGCGCAAG	1183
91	UCCGUCUGGCAGCCUGGAU	860	91	UCCGUCUGGCAGCCUGGAU	860	113	AUCCAGGCUCCAGACGGA	1184
109	UAUCCUCUCCUACCGGCAC	861	109	UAUCCUCUCCUACCGGCAC	861	131	GUGCCGGUAGGAGAGGAUA	1185
127	CCCGCAGACCCCGCAG	862	127	CCCGCAGACCCCGCAG	862	149	CUGCAGGGGCUUCGCGG	1186
145	GCCGCCGUCGCGCCCGG	863	145	GCCGCCGUCGCGCCCGG	863	167	CCGGCGCCGACCGCGGC	1187
163	GGCUCUAGCCCGUGCG	864	163	GGCUCUAGCCCGUGCG	864	185	CGCACAGGGCUAGGGAGCC	1188
181	GCUCAUCUGUCCUGCGCUG	865	181	GCUCAUCUGUCCUGCGCUG	865	203	CAGCGCAGGACAGUUGAGC	1189
199	GCGGGUGCGCGAGUJCC	866	199	GCGGGUGCGCGAGUJCC	866	221	GGAACUCGCGGACCCCGC	1190
217	CACCUCCGCGCCUCCUUCU	867	217	CACCUCCGCGCCUCCUUCU	867	239	AGAAGGAGGCGCGGAGGUG	1191
235	UCUAGACAGGCGCUGGGAG	868	235	UCUAGACAGGCGCUGGGAG	868	257	CUCCAGCGCCUUCUAGA	1192
253	GAAAGAACCGGCUCCCGAG	869	253	GAAAGAACCGGCUCCCGAG	869	275	CUCCGAGCGGCUUCUJUC	1193
271	GUUCUGGGCAUUCGCCCG	870	271	GUUCUGGGCAUUCGCCCG	870	293	CGGGCGAAUUGCCAGAAC	1194
289	GGCUCGAGGUGCAGGAUGC	871	289	GGCUCGAGGUGCAGGAUGC	871	311	GCAUCCUGCACCUUCGAGCC	1195
307	CAGAGCAAGGUGCUGCUGG	872	307	CAGAGCAAGGUGCUGCUGG	872	329	CCAGCAGCACCUUGCUCUG	1196
325	GCCGUCGCCUGUGGGUCU	873	325	GCCGUCGCCUGUGGGUCU	873	347	AGAGCCACAGGGCGACGGC	1197
343	UGCGUGGAGACCCGGGCGG	874	343	UGCGUGGAGACCCGGGCGG	874	365	CGGCCCGGGUCCUCCACGCA	1198
361	GCCUCUGUGGGUUGGCCUA	875	361	GCCUCUGUGGGUUGGCCUA	875	383	UAGGCAAAACCCACAGAGGC	1199
379	AGUGUUUCUCUUGAUCUGC	876	379	AGUGUUUCUCUUGAUCUGC	876	401	GCAGAUCAAGAGAAACACU	1200
397	CCCAGGCUAGCAUACAAA	877	397	CCCAGGCUAGCAUACAAA	877	419	UUUGUAUGCUGAGCCUGGG	1201
415	AAAGACAUACUUAUUA	878	415	AAAGACAUACUUAUUA	878	437	UAUUUGUAAGUAUGUCUUU	1202
433	AAGGCUAAUACACUCUUC	879	433	AAGGCUAAUACACUCUUC	879	455	GAAGAGUUUAUUAGCCUU	1203
451	CAAAUACUUGCAGGGGAC	880	451	CAAAUACUUGCAGGGGAC	880	473	GUCCCCUGCAAGUAUUUG	1204
469	CAGAGGACUUGGACUGGC	881	469	CAGAGGACUUGGACUGGC	881	491	GCCAGUCCAAAGUCCUCUG	1205
487	CUUUGGCCCAUAUUCAGA	882	487	CUUUGGCCCAUAUUCAGA	882	509	UCUGAUUAUUGGGCCAAAG	1206
505	AGUGGCAGUGAGCAAAAGG	883	505	AGUGGCAGUGAGCAAAAGG	883	527	CCCUUUGCUCACUGCCACU	1207
523	GUGGAGGUGACUGAGUGCA	884	523	GUGGAGGUGACUGAGUGCA	884	545	UGCACUCAGUACCCUCCAC	1208

541	AGCGAUGGGCCUUCUUGUA	885	541	AGCGAUGGGCCUUCUUGUA	885	563	UACAGAAGAGGCCAUCGCU	1209
559	AAGACACUCACAAUCCAA	886	559	AAGACACUCACAAUCCAA	886	581	UUGGAAUUGUGAGUGUCUU	1210
577	AAAGUGAUCGGAUUGACA	887	577	AAAGUGAUCGGAUUGACA	887	599	UGUCAUUUCCGAUCACUUU	1211
595	ACUGGAGCCUACAAGUGCU	888	595	ACUGGAGCCUACAAGUGCU	888	617	AGCACUUGUAGGCUCGAGU	1212
613	UUCUACCGGGAACUGACU	889	613	UUCUACCGGGAACUGACU	889	635	AGUCAGUUUCCCGGUAGAA	1213
631	UUGGCCUCGGUUCUUUAUG	890	631	UUGGCCUCGGUUCUUUAUG	890	653	CAUAAAUAGACCCGAGGCCAA	1214
649	GUCUAUGUUCAAGAUUACA	891	649	GUCUAUGUUCAAGAUUACA	891	671	UGUAAUUCUUGAACAUAGAC	1215
667	AGAUUCUCCAUUUUUGCUU	892	667	AGAUUCUCCAUUUUUGCUU	892	689	AAGCAAUAAAUUGGAGAUUCU	1216
685	UCUGUAGUGACCAACAU	893	685	UCUGUAGUGACCAACAU	893	707	CAUGUUGGUCACUAAACAGA	1217
703	GGAGUCGUGUACAUUACUG	894	703	GGAGUCGUGUACAUUACUG	894	725	CAGUAAUUGUACACGACUCC	1218
721	GAGAACAAAAACAAAACUG	895	721	GAGAACAAAAACAAAACUG	895	743	CAGUUUUUGUUUUUGUUCUC	1219
739	GUGGUGAUUCCAUUGUCUG	896	739	GUGGUGAUUCCAUUGUCUG	896	761	CGAGACAUUGAAUACCCAC	1220
757	GGGUCCAUUUCAAAUCUCA	897	757	GGGUCCAUUUCAAAUCUCA	897	779	UGAGAUUUUAAAUGGACCC	1221
775	AACGUGUCACUUUGUGCAA	898	775	AACGUGUCACUUUGUGCAA	898	797	UUGCACAAAGUGACACGUU	1222
793	AGAUACCCAGAAAAGAGAU	899	793	AGAUACCCAGAAAAGAGAU	899	815	AUCUCUUUUCUGGGUAUCU	1223
811	UUUJUUCUUGAGGUUACA	900	811	UUUJUUCUUGAGGUUACA	900	833	UGUUACCAUCAGGAAUAAA	1224
829	AGAAUUUCUUGGACAGCA	901	829	AGAAUUUCUUGGACAGCA	901	851	UGCUGUCCAGGAAAUUCU	1225
847	AAGAAGGCUUUUACUUAUC	902	847	AAGAAGGCUUUUACUUAUC	902	869	GAAUAGUAAAGCCCUUCUU	1226
865	CCAGCUACAUGAUCAGCU	903	865	CCAGCUACAUGAUCAGCU	903	887	AGCUGAUCUAGUAGCUGGG	1227
883	UAUGCUGGCAUGGUCUUCU	904	883	UAUGCUGGCAUGGUCUUCU	904	905	AGAAGACCAUGCCGCAUA	1228
901	UGUGAAGCAAAAAUUUAUG	905	901	UGUGAAGCAAAAAUUUAUG	905	923	CAUUAAUUUUUUGCUUCACA	1229
919	GAUGAAAGUUUACAGUCUA	906	919	GAUGAAAGUUUACAGUCUA	906	941	UAGACUGGUAACUUUUAUC	1230
937	AUUUAGUACAUAGUUGUCG	907	937	AUUUAGUACAUAGUUGUCG	907	959	CGACAACUUAUGACAUAAU	1231
955	GUUGUAGGGUUAUAGGAUUU	908	955	GUUGUAGGGUUAUAGGAUUU	908	977	AAAUCCUUAUACCCUACAAC	1232
973	UAUGAUGUGGUUCUGAGUC	909	973	UAUGAUGUGGUUCUGAGUC	909	995	GACUCAGAACCCACAUCAUA	1233
991	CCGUCUCAUGGAAUUUGAAC	910	991	CCGUCUCAUGGAAUUUGAAC	910	1013	GUUCAUUUCCAUUGAGACGG	1234
1009	CUAUCUGUUGGAGAAAAGC	911	1009	CUAUCUGUUGGAGAAAAGC	911	1031	GCUUUUUCUCCAAACAGAUAG	1235
1027	CUUGUCUUAAAUUUGUACAG	912	1027	CUUGUCUUAAAUUUGUACAG	912	1049	CUGUACAAUUUUAAGACAAG	1236
1045	GCAAGAACUGAACUAAUUG	913	1045	GCAAGAACUGAACUAAUUG	913	1067	CAUUUAGUUCAGUUCUUGC	1237
1063	GUGGGAUUGACUUAACU	914	1063	GUGGGAUUGACUUAACU	914	1085	AGUUGAAGUCAAUCCCCAC	1238
1081	UGGGAUUAACCUUUCUUCGA	915	1081	UGGGAUUAACCUUUCUUCGA	915	1103	UCGAAGAAGGGUUAUCCCA	1239
1099	AAGCAUCAGCAUAAGAAAC	916	1099	AAGCAUCAGCAUAAGAAAC	916	1121	GUUUCUUUUGCUGAUGCUU	1240
1117	CUUGUAAACCCGAGACCUAA	917	1117	CUUGUAAACCCGAGACCUAA	917	1139	UUAGGUCUCGGUUUUAACAG	1241
1135	AAACCCAGUCUGGGAGUG	918	1135	AAACCCAGUCUGGGAGUG	918	1157	CACUCCAGACUGGGUUUUU	1242
1153	GAGAUGAAGAAUUUUUGA	919	1153	GAGAUGAAGAAUUUUUGA	919	1175	UCAAAAAUUUUCUUAUCUC	1243
1171	AGCACCUUAACUUAUAGAU	920	1171	AGCACCUUAACUUAUAGAU	920	1193	CAUCUAUAGUUAAGGUGCU	1244

1189	GGUGUAACCCGGAGUGACC	921	1189	GGUGUAACCCGGAGUGACC	921	1211	GGUCACUCCGGGUUACACC	1245
1207	CAAGGAUUGUACACCUUG	922	1207	CAAGGAUUGUACACCUUG	922	1229	CACAGGUGUACAAUCCUUG	1246
1225	GCAGCAUCCAGUGGCGUGA	923	1225	GCAGCAUCCAGUGGCGUGA	923	1247	UCAGCCACUGGAUGCUGC	1247
1243	AUGACCAAGAAGAACAGCA	924	1243	AUGACCAAGAAGAACAGCA	924	1265	UGCUGUUCUUCUUGGUGAU	1248
1261	ACAUUUGUCAGGUGCCAU	925	1261	ACAUUUGUCAGGUGCCAU	925	1283	CAUGGACCCUGACAAAUGU	1249
1279	GA AAAACCUUUGUUGCUU	926	1279	GA AAAACCUUUGUUGCUU	926	1301	AAGCAACAAAAGGUUUUUC	1250
1297	UUUGGAAGUGGCAUGGAAU	927	1297	UUUGGAAGUGGCAUGGAAU	927	1319	AUUCCAUGCCACUCCAAA	1251
1315	UCUCUGGUGGAAGCCACGG	928	1315	UCUCUGGUGGAAGCCACGG	928	1337	CCGUGGCUUCCACCCAGAGA	1252
1333	GUGGGGAGGUGUCAGAA	929	1333	GUGGGGAGGUGUCAGAA	929	1355	UUCUGACACGCUCCCCAC	1253
1351	AUCCUGCGAAGUACCUUG	930	1351	AUCCUGCGAAGUACCUUG	930	1373	CAAGGUACUUCGACGGAU	1254
1369	GGUACCCACCCCCAGAAA	931	1369	GGUACCCACCCCCAGAAA	931	1391	UUUCUGGGGUGGGUAAACC	1255
1387	AUAAAUGGUUAAAAAUG	932	1387	AUAAAUGGUUAAAAAUG	932	1409	CAUUUUUAUACCAUUUUU	1256
1405	GGAUACCCCUUGAGUCCA	933	1405	GGAUACCCCUUGAGUCCA	933	1427	UGGACUCAAAGGGGUUUUCC	1257
1423	AUACACAAUUAAAGCGG	934	1423	AUACACAAUUAAAGCGG	934	1445	CCGCUUUAUUUGUGUGAUU	1258
1441	GGGCAUGUACUGACGUAUA	935	1441	GGGCAUGUACUGACGUAUA	935	1463	UAAUCGUCAGUACAUGCC	1259
1459	AUGGAAGUGAGUGAAAGAG	936	1459	AUGGAAGUGAGUGAAAGAG	936	1481	CUCUUUCACUCACUUCUCCAU	1260
1477	GACACAGGAUUUACACUG	937	1477	GACACAGGAUUUACACUG	937	1499	CAGUGUAUUUCCUGUGUC	1261
1495	GUCAUCCUUACCAUCCCA	938	1495	GUCAUCCUUACCAUCCCA	938	1517	UGGGAUUGGUAGGGAUGAC	1262
1513	AUUUCAAAGGAGAAAGCAGA	939	1513	AUUUCAAAGGAGAAAGCAGA	939	1535	UCUGCUUCUCCUUUGAAAAU	1263
1531	AGCAUGUGGUCUCUCUGG	940	1531	AGCAUGUGGUCUCUCUGG	940	1553	CCAGAGAGACCAUUGGCU	1264
1549	GUUGUGUAUGUCCACCCC	941	1549	GUUGUGUAUGUCCACCCC	941	1571	GGGUGGGACAUACACAAC	1265
1567	CAGAUUGGUGAGAAUUCUC	942	1567	CAGAUUGGUGAGAAUUCUC	942	1589	GAGAUUUCUCACCAAUUCUG	1266
1585	CUAAUCUCUCCUGUGGAUU	943	1585	CUAAUCUCUCCUGUGGAUU	943	1607	AAUCCACAGGAGAGAUUAG	1267
1603	UCCUACACGUCGACACCA	944	1603	UCCUACACGUCGACACCA	944	1625	UGGUGCCGUACUGGUGGGA	1268
1621	ACUCAACGCUACAUUGUA	945	1621	ACUCAACGCUACAUUGUA	945	1643	UACAUUGUCAGCGUUUGAGU	1269
1639	ACGGUCUAUGCCAUUCCUC	946	1639	ACGGUCUAUGCCAUUCCUC	946	1661	GAGGAUUGGCAUAGACCGU	1270
1657	CCCCGGCAUACAUUCCACU	947	1657	CCCCGGCAUACAUUCCACU	947	1679	AGUGGAUGUGAUUGCGGGG	1271
1675	UGGUUUUGGCAGUUGGAGG	948	1675	UGGUUUUGGCAGUUGGAGG	948	1697	CCUCCAAUUGCCAAUACCA	1272
1693	GAAGAGUUGCGCAACGAGC	949	1693	GAAGAGUUGCGCAACGAGC	949	1715	GCUCGUUGGCGCACUCUUC	1273
1711	CCAGCCAAAGCUGUCUCAG	950	1711	CCAGCCAAAGCUGUCUCAG	950	1733	CUGAGACAGCUUGGUGGG	1274
1729	GUGACAAAACCAUACCCU	951	1729	GUGACAAAACCAUACCCU	951	1751	AAGGUAUUGGUUUUGUCAC	1275
1747	UGUGAAGAAUUGGAGAGUG	952	1747	UGUGAAGAAUUGGAGAGUG	952	1769	CACUUCUCCAUUCUUCACA	1276
1765	GUGGAGGACUCCAGGGAG	953	1765	GUGGAGGACUCCAGGGAG	953	1787	CUCCUGGGAAGUCCUCCAC	1277
1783	GGAAUAAAAUUUGAAGUUA	954	1783	GGAAUAAAAUUUGAAGUUA	954	1805	UAACUUCAAUUUUUUAUCC	1278
1801	AAUAAAAUCAAUUUUGCUC	955	1801	AAUAAAAUCAAUUUUGCUC	955	1823	GAGCAAAUUGAUUUUUUUAU	1279
1819	CUAAUUGAAGGAAAAAACA	956	1819	CUAAUUGAAGGAAAAAACA	956	1841	UGUUUUUUUCCUUCAAUUAUAG	1280

1837	AAAACUGUAAGUACCCUUG	957	1837	AAAACUGUAAGUACCCUUG	957	1859	CAAGGGUACUACAGUUUU	1281
1855	GUUAUCCAAGCGGCAAAUG	958	1855	GUUAUCCAAGCGGCAAAUG	958	1877	CAUUUGCCGUUGGAUAC	1282
1873	GUGUCAGCUUUGUACAAAU	959	1873	GUGUCAGCUUUGUACAAAU	959	1895	AUUUGUACAAAGCUGACAC	1283
1891	UGUGAAGCGGUCAACAAAG	960	1891	UGUGAAGCGGUCAACAAAG	960	1913	CUUUUGUACCCGUUCACA	1284
1909	GUCCGGAGAGGAGAGAGGG	961	1909	GUCCGGAGAGGAGAGAGGG	961	1931	CCCUCUCUCUCCUCCCGAC	1285
1927	GUGAUCUCUUCUCCACGUGA	962	1927	GUGAUCUCUUCUCCACGUGA	962	1949	UCACGUGGAAGGAGACAC	1286
1945	ACCAGGGGUCCUGAAAUA	963	1945	ACCAGGGGUCCUGAAAUA	963	1967	UAAUUUCAGGACCCUUGU	1287
1963	ACUUUGCAACCUUGACAUUC	964	1963	ACUUUGCAACCUUGACAUUC	964	1985	GCAUUGCAGGUUGCAAAGU	1288
1981	CAGCCACUGAGCAGGAGA	965	1981	CAGCCACUGAGCAGGAGA	965	2003	UCUCCUGCUCAGUGGGCUG	1289
1999	AGCGUGUCUUUGUGGUGCA	966	1999	AGCGUGUCUUUGUGGUGCA	966	2021	UGCACCACAAAGACACGCU	1290
2017	ACUGCAGACAGAUUACGU	967	2017	ACUGCAGACAGAUUACGU	967	2039	ACGUAGAUUCUGUCUGCAGU	1291
2035	UUUGAGAACCUCACAUUGU	968	2035	UUUGAGAACCUCACAUUGU	968	2057	ACCAUGUGAGGUUCUCAAA	1292
2053	UACAAGCUUGGCCCCACAGC	969	2053	UACAAGCUUGGCCCCACAGC	969	2075	GCUGUGGGCCAAGCUUGUA	1293
2071	CCUCUGCCAAUCCAUGUGG	970	2071	CCUCUGCCAAUCCAUGUGG	970	2093	CCACAUGGAUUGGCAGAGG	1294
2089	GGAGAGUUGCCACACACUG	971	2089	GGAGAGUUGCCACACACUG	971	2111	CAGGUGUGGGCAACUCUCC	1295
2107	GUUUUGAAGAUAUUGGAUA	972	2107	GUUUUGAAGAUAUUGGAUA	972	2129	UAUCCAAGUUCUUGCAAAC	1296
2125	ACUUCUUUGGAAUUAUUGA	973	2125	ACUUCUUUGGAAUUAUUGA	973	2147	CAUUCAAUUUCCAAGAGU	1297
2143	GCCACCAUUGUUCUUAUA	974	2143	GCCACCAUUGUUCUUAUA	974	2165	UAUUAGAGAACAUGGUGGC	1298
2161	AGCACAAUUGACAUUUUGA	975	2161	AGCACAAUUGACAUUUUGA	975	2183	UCAAAUUGCAUUUUGUGCU	1299
2179	AUCAUGGAGCUUAAGAAUG	976	2179	AUCAUGGAGCUUAAGAAUG	976	2201	CAUUCUUAAAGCUCUUGAU	1300
2197	GCAUCCUUGCAGGACCAAG	977	2197	GCAUCCUUGCAGGACCAAG	977	2219	CUUGGUCCUGCAAGGAUGC	1301
2215	GGAGACUAUUGUCUGCCUUG	978	2215	GGAGACUAUUGUCUGCCUUG	978	2237	CAAGGCAGACAUAGUCUCC	1302
2233	GCUCAAGACAGGAAGACCA	979	2233	GCUCAAGACAGGAAGACCA	979	2255	UGGUCUUCUGUCUUGAGC	1303
2251	AAGAAAGACAUUGCGUGG	980	2251	AAGAAAGACAUUGCGUGG	980	2273	CCACGCAUUGUCUUUUCUU	1304
2269	GUCAGGAGCUCACAGUCC	981	2269	GUCAGGAGCUCACAGUCC	981	2291	GGACUGUGAGCUGCCUGAC	1305
2287	CUAGAGCGUGUGGCACCCA	982	2287	CUAGAGCGUGUGGCACCCA	982	2309	UGGGUGCCACACGCUUAG	1306
2305	ACGAUCACAGGAACCUUG	983	2305	ACGAUCACAGGAACCUUG	983	2327	CCAGGUUUCUGUGAUCGU	1307
2323	GAGAAUCAGACGACAAGUA	984	2323	GAGAAUCAGACGACAAGUA	984	2345	UACUUGUCUGUCUGAUUCUC	1308
2341	AUUUGGGAAAGCAUCGAAG	985	2341	AUUUGGGAAAGCAUCGAAG	985	2363	CUUCGAUGCUUUCGCCAAU	1309
2359	GUCUCAUGCAGGCAUCUG	986	2359	GUCUCAUGCAGGCAUCUG	986	2381	CAGAUCCCGUGCAUGAGAC	1310
2377	GGGAUCCCCCUCCACAGA	987	2377	GGGAUCCCCCUCCACAGA	987	2399	UCUGUGGAGGGGGAUUCUCC	1311
2395	AUCAUGUGGUUUAAGAUUA	988	2395	AUCAUGUGGUUUAAGAUUA	988	2417	UAUCUUUAACCAACAUAGU	1312
2413	AAUGAGACCCUUGUAGAAG	989	2413	AAUGAGACCCUUGUAGAAG	989	2435	CUUCUACAAAGGUCUUAUU	1313
2431	GACUCAGGCAUUGUAUUGA	990	2431	GACUCAGGCAUUGUAUUGA	990	2453	UCAUUAACAAUGCCUGAGUC	1314
2449	AAGGAUGGGAAACCGGAACC	991	2449	AAGGAUGGGAAACCGGAACC	991	2471	GGUUCGGUUCUCCAUCCUU	1315
2467	CUCACUAUCCGACAGAGUGA	992	2467	CUCACUAUCCGACAGAGUGA	992	2489	UCACUCUGCGGAUAGUGAG	1316

2485	AGGAAGGAGGACGAAGGCC	993	2485	AGGAAGGAGGACGAAGGCC	993	2507	GGCCUUCGUCUCCUCCU	1317
2503	CUCUACACCGCCAGGCAU	994	2503	CUCUACACCGCCAGGCAU	994	2525	AUGCCUGGCAGGUGUAGAG	1318
2521	UGCAGUGUUCUUGGCUGUG	995	2521	UGCAGUGUUCUUGGCUGUG	995	2543	CACAGCCAAAGAACACUGCA	1319
2539	GCAAAAGUGGAGGCAUUUU	996	2539	GCAAAAGUGGAGGCAUUUU	996	2561	AAAAGCCUCCACUUUUUGC	1320
2557	UUCAUAAUAGAGGUGCCC	997	2557	UUCAUAAUAGAGGUGCCC	997	2579	GGCACCUUCUUAUUAUGAA	1321
2575	CAGGAAAAGACGAACUUGG	998	2575	CAGGAAAAGACGAACUUGG	998	2597	CCAAGUUCGUCUUUUCCUG	1322
2593	GAUAUCAUUAUUCUAGUAG	999	2593	GAUAUCAUUAUUCUAGUAG	999	2615	CUACUAGAAUAAUUAUUUC	1323
2611	GGCACGGCGGUGAUUGCCA	1000	2611	GGCACGGCGGUGAUUGCCA	1000	2633	UGGCAUACCCGCCUGGCC	1324
2629	AUGUUCUUCUGGCUACUUC	1001	2629	AUGUUCUUCUGGCUACUUC	1001	2651	GAAGUAGCCAGAAACAUAU	1325
2647	CUUGUCAUCAUCCUACGGA	1002	2647	CUUGUCAUCAUCCUACGGA	1002	2669	UCCGUAGGAUUAUAGACAAG	1326
2665	ACCGUUAAGCGGCCAAUUG	1003	2665	ACCGUUAAGCGGCCAAUUG	1003	2687	CAUUGCCCCGUUAACGGU	1327
2683	GGAGGGGAACUGAAGACAG	1004	2683	GGAGGGGAACUGAAGACAG	1004	2705	CUGUCUUCAGUUCUCCUCC	1328
2701	GGCUACUUCUCCUACUGUA	1005	2701	GGCUACUUCUCCUACUGUA	1005	2723	UGACGAUGGACAAAGUAGCC	1329
2719	AUGGAUCCAGAUAAACUCC	1006	2719	AUGGAUCCAGAUAAACUCC	1006	2741	GGAGUUAUCUUGGAUCCAU	1330
2737	CCAUUGGAUAAACAUUUGUG	1007	2737	CCAUUGGAUAAACAUUUGUG	1007	2759	CACAAUUAUUAUCCAAUUGG	1331
2755	GAACGACUGCCUUAUUAUG	1008	2755	GAACGACUGCCUUAUUAUG	1008	2777	CAUCAUAAGGCAGUCGUUC	1332
2773	GCCAGCAAAUGGAAUUC	1009	2773	GCCAGCAAAUGGAAUUC	1009	2795	GGAAUUAUUAUUAUUGGCGC	1333
2791	CCCAGAGACCGGCUAAGC	1010	2791	CCCAGAGACCGGCUAAGC	1010	2813	GCUUCAGCCGUCUCCUUGG	1334
2809	CUAGGUAAGCCUUAUUGGCC	1011	2809	CUAGGUAAGCCUUAUUGGCC	1011	2831	GGCCAAAGAGGCUUAACCUAG	1335
2827	CGUGGUGCCUUAUUGGCCAAG	1012	2827	CGUGGUGCCUUAUUGGCCAAG	1012	2849	CUUGGCCAAAGGACCCACG	1336
2845	GUGAUUGAAGCAGAUUGCCU	1013	2845	GUGAUUGAAGCAGAUUGCCU	1013	2867	AGGCAUCUGCUUCAAUUCAC	1337
2863	UUUGGAAUUGACAAGACAG	1014	2863	UUUGGAAUUGACAAGACAG	1014	2885	CUGUCUUAUGCAAUUCCAAA	1338
2881	GCAACUUGCAGGACAGUAG	1015	2881	GCAACUUGCAGGACAGUAG	1015	2903	CUACUGUCCUGCAAGUUGC	1339
2899	GCAGUCAAAAUUGUAAAG	1016	2899	GCAGUCAAAAUUGUAAAG	1016	2921	CUUUAACAUAUUUUGACUGC	1340
2917	GAAGGAGCAACACACAGUG	1017	2917	GAAGGAGCAACACACAGUG	1017	2939	CACUGUGUGUUGCUCCUUC	1341
2935	GAGCAUCGAGCUCUCAUGU	1018	2935	GAGCAUCGAGCUCUCAUGU	1018	2957	ACAUGAGAGCUCGAGUCCUC	1342
2953	UCUGAACUCAAGAUCCUCA	1019	2953	UCUGAACUCAAGAUCCUCA	1019	2975	UGAGGAUCUUGAGUUCAGA	1343
2971	AUUCAUUUUGGUCACCAUC	1020	2971	AUUCAUUUUGGUCACCAUC	1020	2993	GAUGGUGACCAUAUUGAAU	1344
2989	CUCAAUUGUGUUAACCUUC	1021	2989	CUCAAUUGUGUUAACCUUC	1021	3011	GAAGGUUGACCAUAUUGAG	1345
3007	CUAGGUGCCUGUACCAAGC	1022	3007	CUAGGUGCCUGUACCAAGC	1022	3029	GCUUGGUACAGGCACCUJAG	1346
3025	CCAGGAGGGCCACUCAUGG	1023	3025	CCAGGAGGGCCACUCAUGG	1023	3047	CCAUGAGUGGCCUCCUGG	1347
3043	GUGAUUGUGGAAUUCUGCA	1024	3043	GUGAUUGUGGAAUUCUGCA	1024	3065	UGCAGAAUUAUUAUUAUUCAC	1348
3061	AAAUUUGGAAACCUUGUCCA	1025	3061	AAAUUUGGAAACCUUGUCCA	1025	3083	UGGACAGGUUUUCCAAUUUU	1349
3079	ACUUAACUGAGGAGCAAGA	1026	3079	ACUUAACUGAGGAGCAAGA	1026	3101	UCUUGCUCCUACAGGUAAGU	1350
3097	AGAAUUGAAUUUUGUCCCU	1027	3097	AGAAUUGAAUUUUGUCCCU	1027	3119	AGGGACAAAUUAUUAUUCU	1351
3115	UACAAGACCAAAAGGGGCAC	1028	3115	UACAAGACCAAAAGGGGCAC	1028	3137	GUGCCCUUUUGGUCUUGUA	1352

3133	CGAUUCCGUAAGGGAAAG	1029	3133	CGAUUCCGUAAGGGAAAG	1029	3155	CUUUCCCUUACGGAAUCG	1353
3151	GACUACGUUGGAGCAUCC	1030	3151	GACUACGUUGGAGCAUCC	1030	3173	GGAUUGCUCCAAACGUAGUC	1354
3169	CCUGUGGAUCUGAAACGGC	1031	3169	CCUGUGGAUCUGAAACGGC	1031	3191	GCCGUUUCAGAUCCACAGG	1355
3187	CGCUUGGACAUCAACCA	1032	3187	CGCUUGGACAUCAACCA	1032	3209	UGGUGAUGCUGUCCAAAGC	1356
3205	AGUAGCCAGAGCUCAGCCA	1033	3205	AGUAGCCAGAGCUCAGCCA	1033	3227	UGGUGAGCUCUGGCUACU	1357
3223	AGCUCUGGAUUUGGAGG	1034	3223	AGCUCUGGAUUUGGAGG	1034	3245	CCUCCACAAUCCAGAGCU	1358
3241	GAGAAGUCCUCAGUGAUG	1035	3241	GAGAAGUCCUCAGUGAUG	1035	3263	CAUCACUGAGGGACUUCUC	1359
3259	GUAGAAGAAGGAAAGCUC	1036	3259	GUAGAAGAAGGAAAGCUC	1036	3281	GAGCUUCCUCUUCUUCUAC	1360
3277	CCUGAAGAUUGUAUAGG	1037	3277	CCUGAAGAUUGUAUAGG	1037	3299	CCUUUAACAGAUUUCAGG	1361
3295	GACUCCUGACCUUGGAGC	1038	3295	GACUCCUGACCUUGGAGC	1038	3317	GUCCAAAGGUCAGGAAGUC	1362
3313	CAUCUCAUCUGUACAGCU	1039	3313	CAUCUCAUCUGUACAGCU	1039	3335	AGCUGUAACAGAUAGAGU	1363
3331	UUCCAAGUGGCUAAGGGCA	1040	3331	UUCCAAGUGGCUAAGGGCA	1040	3353	UGCCUUAGCCACUUGGAA	1364
3349	AUGGAGUUCUUGGCAUCGC	1041	3349	AUGGAGUUCUUGGCAUCGC	1041	3371	GCGAUGCCAAAGAACUCCAU	1365
3367	CGAAAGUGUAUCCACAGGG	1042	3367	CGAAAGUGUAUCCACAGGG	1042	3389	CCUGUGGUAACACUUCUG	1366
3385	GACCUGGCGGCACGAAUA	1043	3385	GACCUGGCGGCACGAAUA	1043	3407	UAUUUCGUGCCGCCAGGUC	1367
3403	AUCCUCUUAUCCGAGAGA	1044	3403	AUCCUCUUAUCCGAGAGA	1044	3425	UCUUCUCCGAUAAAGAGAU	1368
3421	AACGUGGUUAAAUUCUGUG	1045	3421	AACGUGGUUAAAUUCUGUG	1045	3443	CACAGAUUUAAACCACGUU	1369
3439	GACUUUGGCUUGGCCCGGG	1046	3439	GACUUUGGCUUGGCCCGGG	1046	3461	CCCGGCCCAAGCCAAAGUC	1370
3457	GAUUAUUUAUAAAGAUCCAG	1047	3457	GAUUAUUUAUAAAGAUCCAG	1047	3479	CUGGAUCUUUAUAAAUUUC	1371
3475	GAUUAUGUCAGAAAGGAG	1048	3475	GAUUAUGUCAGAAAGGAG	1048	3497	CUCCUUUUCUGACAUAAUC	1372
3493	GAUGCUCGCCUCCUUUGA	1049	3493	GAUGCUCGCCUCCUUUGA	1049	3515	UCAAAGGGAGGCGGAGCAUC	1373
3511	AAUUGGAUGGCCCCAGAAA	1050	3511	AAUUGGAUGGCCCCAGAAA	1050	3533	UUUCUGGGGCAUCCAUUU	1374
3529	ACAAUUUUUACAGAGUGU	1051	3529	ACAAUUUUUACAGAGUGU	1051	3551	ACACUCUGUCAAAAUIUGU	1375
3547	UACACAAUCCAGAGUGACG	1052	3547	UACACAAUCCAGAGUGACG	1052	3569	CGUCACUCUGGAUUGUGUA	1376
3565	GUCUGGUCUUUUGGUGUUU	1053	3565	GUCUGGUCUUUUGGUGUUU	1053	3587	AAACACCAAAAGACCAGAC	1377
3583	UUGCUGUGGGAUUAUUUU	1054	3583	UUGCUGUGGGAUUAUUUU	1054	3605	AAAAUUAUUUCCACAGCAA	1378
3601	UCCUUAGGUGCUUCUCCAU	1055	3601	UCCUUAGGUGCUUCUCCAU	1055	3623	AUGGAGAAGCACCUAAGGA	1379
3619	UAUCCUGGGUUAAGAUUG	1056	3619	UAUCCUGGGUUAAGAUUG	1056	3641	CAUUCUUUACCCCGAGUA	1380
3637	GAUGAAGAUUUUGUAGGC	1057	3637	GAUGAAGAUUUUGUAGGC	1057	3659	GCCUACAAAUUUCUUCUUC	1381
3655	CGAUUGAAAGAGGAACUA	1058	3655	CGAUUGAAAGAGGAACUA	1058	3677	UAGUUCUUCUUCUUCUUC	1382
3673	AGAAUGAGGGCCCUUUAU	1059	3673	AGAAUGAGGGCCCUUUAU	1059	3695	AAUCAGGGGCCCUCAUUCU	1383
3691	UAUACUACACCAAGAAUUG	1060	3691	UAUACUACACCAAGAAUUG	1060	3713	ACAUUCUGGUGUAGUAUA	1384
3709	UACCAGACCAUGCUGGACU	1061	3709	UACCAGACCAUGCUGGACU	1061	3731	AGUCCAGUGGUGUGGUA	1385
3727	UGCUGGCACGGGAGCCCA	1062	3727	UGCUGGCACGGGAGCCCA	1062	3749	UGGCUCCCGGUGCCAGCA	1386
3745	AGUCAGAGACCCACGUUUU	1063	3745	AGUCAGAGACCCACGUUUU	1063	3767	AAACGUGGGUCUCUGACU	1387
3763	UCAGAGUUGGUGGAACAUU	1064	3763	UCAGAGUUGGUGGAACAUU	1064	3785	AAGUUUCCACCAACUCUGA	1388

3781	UUUGGAAAUUCUUGCAAG	1065	3781	UUUGGAAAUUCUUGCAAG	1065	3803	CUUGCAAGAGAUUCCCAA	1389
3799	GUUAUUGCUCAGCAGGAUG	1066	3799	GUUAUUGCUCAGCAGGAUG	1066	3821	CAUCCUGCUGAGCAUJAGC	1390
3817	GGCAAAGACUACAUUGUUC	1067	3817	GGCAAAGACUACAUUGUUC	1067	3839	GAACAAUGUAGUCUUGGCC	1391
3835	CUUCCGAUACAGAGACUU	1068	3835	CUUCCGAUACAGAGACUU	1068	3857	AAGUCUCUGAUUCCGGAAG	1392
3853	UUGAGCAUGGAAGAGGAUU	1069	3853	UUGAGCAUGGAAGAGGAUU	1069	3875	AAUCCUUCUCCAUUGCUCAA	1393
3871	UCUGGACUCUCUGGCCUA	1070	3871	UCUGGACUCUCUGGCCUA	1070	3893	UAGGCAGAGAGAGUCCAGA	1394
3889	ACCUCACCGUUGUCCUGUA	1071	3889	ACCUCACCGUUGUCCUGUA	1071	3911	UACAGGAAACAGGUGAGGU	1395
3907	AUGGAGGAGGAGGAAGUAU	1072	3907	AUGGAGGAGGAGGAAGUAU	1072	3929	AUACUCCUCCUCCUCCAU	1396
3925	UGUGACCCCAAUUCCAUU	1073	3925	UGUGACCCCAAUUCCAUU	1073	3947	AAUGGAAUUGGGGUCACA	1397
3943	UAUGACAACACAGCAGGAA	1074	3943	UAUGACAACACAGCAGGAA	1074	3965	UUCUGCUGUGUUGUCAUA	1398
3961	AUCAGUCAGUACUCUGCAGA	1075	3961	AUCAGUCAGUACUCUGCAGA	1075	3983	UCUGCAGAUACUGACUGAU	1399
3979	AACAGUAAGCGAAAGAGCC	1076	3979	AACAGUAAGCGAAAGAGCC	1076	4001	GGCUCUUUCGCUUACUGUU	1400
3997	CGGCCUGUGAGUGUAAAAA	1077	3997	CGGCCUGUGAGUGUAAAAA	1077	4019	UUUUUACACUCACAGGCCG	1401
4015	ACAUUUGAAGAUUCCCGU	1078	4015	ACAUUUGAAGAUUCCCGU	1078	4037	ACGGGAUAUCUUCAAUUGU	1402
4033	UUAGAAGAACCCAGAGUAA	1079	4033	UUAGAAGAACCCAGAGUAA	1079	4055	UUACUUCUGGUUUCUUCUAA	1403
4051	AAAGUAAUCCCAAGUACA	1080	4051	AAAGUAAUCCCAAGUACA	1080	4073	UGUCAUCUGGGAUUACUUU	1404
4069	AACAGACGGACAGUGGUA	1081	4069	AACAGACGGACAGUGGUA	1081	4091	UACCACUGCCGUCUGUUU	1405
4087	AUGGUUCUUGCCUCAGAAG	1082	4087	AUGGUUCUUGCCUCAGAAG	1082	4109	CUUCUGAGGCCAAGACCAU	1406
4105	GAGCUGAAAAAUUUGGAAG	1083	4105	GAGCUGAAAAAUUUGGAAG	1083	4127	CUUCCAAAGUUUUCAGCUC	1407
4123	GACAGAACCACAAUUAUCUC	1084	4123	GACAGAACCACAAUUAUCUC	1084	4145	GAGAUAAUUGGUUCUGUC	1408
4141	CCAUUUUUGGUGGAUUGG	1085	4141	CCAUCUUUUGGUGGAUUGG	1085	4163	CCAUUCCACCACAAAGAUUG	1409
4159	GUGCCCAAGCAAAAGCAGGG	1086	4159	GUGCCCAAGCAAAAGCAGGG	1086	4181	CCUUGCUUUUGCUGGGCAC	1410
4177	GAGUCUGUGGCAUCUGAAG	1087	4177	GAGUCUGUGGCAUCUGAAG	1087	4199	CUUCAGAUCCACACAGACUC	1411
4195	GGCUCAAACCAGACACAGCG	1088	4195	GGCUCAAACCAGACACAGCG	1088	4217	CGCUUGUCUGGUUUUGAGCC	1412
4213	GGCUACCAUCCGGAUAUC	1089	4213	GGCUACCAUCCGGAUAUC	1089	4235	GAUAUCCGGACUGGUAGCC	1413
4231	CACUCCGAUGACACAGACA	1090	4231	CACUCCGAUGACACAGACA	1090	4253	UGUCUGUGUCAUCGGAGUG	1414
4249	ACCACCGUGUACUCCAGUG	1091	4249	ACCACCGUGUACUCCAGUG	1091	4271	CACUGGAGUACACGGUGGU	1415
4267	GAGGAAGCAGAACUUUUA	1092	4267	GAGGAAGCAGAACUUUUA	1092	4289	UUAAAAGUUCUUCUUCUUC	1416
4285	AAGCUGAUAGAGAUUGGAG	1093	4285	AAGCUGAUAGAGAUUGGAG	1093	4307	CUCCAAUCUCUUAUCAGCUU	1417
4303	GUGCAAAACCGGUAGCAGAG	1094	4303	GUGCAAAACCGGUAGCAGAG	1094	4325	CUGUGCUACCCGUUUUGCAC	1418
4321	GCCCAGAUUCCAGCCUG	1095	4321	GCCCAGAUUCCAGCCUG	1095	4343	CAGGUGGAGAAUUCUGGCC	1419
4339	GACUCGGGGACACACACUGA	1096	4339	GACUCGGGGACACACACUGA	1096	4361	UCAGUGUGUCCCGGAGUC	1420
4357	AGCUCUCCUCCUGUUUUA	1097	4357	AGCUCUCCUCCUGUUUUA	1097	4379	UUUAAACAGGAGGAGAGCU	1421
4375	AAGGAAGCAUCCACACCCC	1098	4375	AAGGAAGCAUCCACACCCC	1098	4397	GGGGUGUGGAUUCUUCUUU	1422
4393	CAACUCCCGGACAUACAU	1099	4393	CAACUCCCGGACAUACAU	1099	4415	AUGUGAUGUCCCGGAGUUG	1423
4411	UGAGAGGUCUGCUCAGAUU	1100	4411	UGAGAGGUCUGCUCAGAUU	1100	4433	AAUCUGAGCAGACCCUCA	1424

4429	UUUGAAGUGUUGUUCUUUC	1101	4429	UUUGAAGUGUUGUUCUUUC	1101	4451	GAAGAACAACACUUCAAA	1425
4447	CCACCAGCAGGAAGUAGCC	1102	4447	CCACCAGCAGGAAGUAGCC	1102	4469	GGCUACUUCUGUGUGG	1426
4465	CGCAUUGAUUUUUAUUUC	1103	4465	CGCAUUGAUUUUUAUUUC	1103	4487	GAUAUGAAAAUUAUUGCG	1427
4483	CGACAACAGAAAAAGGACC	1104	4483	CGACAACAGAAAAAGGACC	1104	4505	GGUCCUUUUUUGUGUUGCG	1428
4501	CUCGGACUGCAGGGAGCCA	1105	4501	CUCGGACUGCAGGGAGCCA	1105	4523	UGGCUCCUGCAGUCCGAG	1429
4519	AGUCUUUJAGGCAUAUCCU	1106	4519	AGUCUUUJAGGCAUAUCCU	1106	4541	AGGAUAUGCCUAGAAAGACU	1430
4537	UGGAAGAGGCUUGUGACCC	1107	4537	UGGAAGAGGCUUGUGACCC	1107	4559	GGGUCACAAAGCCUUCUCCA	1431
4555	CAAGAAUGUGUCUGUGUCU	1108	4555	CAAGAAUGUGUCUGUGUCU	1108	4577	AGACACAGACACAUUCUUG	1432
4573	UUCUCCAGUGUUGACCUUG	1109	4573	UUCUCCAGUGUUGACCUUG	1109	4595	CAGGUCACACUGGGAGAA	1433
4591	GAUCCUUCUUUUUAUUA	1110	4591	GAUCCUUCUUUUUAUUA	1110	4613	UGAAUGAAAAAGAGGAUC	1434
4609	AUUUAAAAAGCAUUAUUA	1111	4609	AUUUAAAAAGCAUUAUUA	1111	4631	AUGAUAAUGCUUUUUAUUA	1435
4627	UGCCCCUGUGCGGGUCUC	1112	4627	UGCCCCUGUGCGGGUCUC	1112	4649	GAGACCCGAGCAGGGGCA	1436
4645	CACCAUGGGUUUJAGAACAA	1113	4645	CACCAUGGGUUUJAGAACAA	1113	4667	UUGUUUJAAACCCAUUGUG	1437
4663	AAGAGCUUCAAAGCAUUGGC	1114	4663	AAGAGCUUCAAAGCAUUGGC	1114	4685	GCCAUUUGUUGAAAGCUUUC	1438
4681	CCCCAUCCUCAAAGAAUUA	1115	4681	CCCCAUCCUCAAAGAAUUA	1115	4703	UACUUCUUUGAGGAUUGGG	1439
4699	AGCAGUACCUUGGGAGCUG	1116	4699	AGCAGUACCUUGGGAGCUG	1116	4721	CAGCUCCCGAGGUAUCUUC	1440
4717	GACACUUCUGUJAAACUAG	1117	4717	GACACUUCUGUJAAACUAG	1117	4739	CUAGUUUUAACAGAAUGUUC	1441
4735	GAAGAUAAACAGGCAACG	1118	4735	GAAGAUAAACAGGCAACG	1118	4757	CGUUGCCUGGUUUAUCUUC	1442
4753	GUAAUGUUCGAGGUGUUG	1119	4753	GUAAUGUUCGAGGUGUUG	1119	4775	CAACACUUGCAACACUUAAC	1443
4771	GAAGAUGGGAGGAUUUUGC	1120	4771	GAAGAUGGGAGGAUUUUGC	1120	4793	GCAAAUCCUUCCCAUUCUUC	1444
4789	CAGGCGUGAGUCUAUCCAA	1121	4789	CAGGCGUGAGUCUAUCCAA	1121	4811	UUGGAUAGACUCAGCCUUG	1445
4807	AGAGGUUUUUGUUJAGGACG	1122	4807	AGAGGUUUUUGUUJAGGACG	1122	4829	CGUCCUAAACAAAAGCCUUC	1446
4825	GUGGGUCCCAAGCCAAAGCC	1123	4825	GUGGGUCCCAAGCCAAAGCC	1123	4847	GGCUUGGCUUGGGACCCAC	1447
4843	CUJAAUGUGGAAUUCGGA	1124	4843	CUJAAUGUGGAAUUCGGA	1124	4865	UCCGAUUUCCACACUUAAG	1448
4861	AUUGAUAGAAAGGAAGACU	1125	4861	AUUGAUAGAAAGGAAGACU	1125	4883	AGUCUCCUUUCUAUCAAU	1449
4879	UAACGUUACCUUGCUUUGG	1126	4879	UAACGUUACCUUGCUUUGG	1126	4901	CCAAAGCAAGGUAAACGUUA	1450
4897	GAGAGUACUUGGAGCCUGCA	1127	4897	GAGAGUACUUGGAGCCUGCA	1127	4919	UGCAGGCUCCAGUACUCUC	1451
4915	AAUUGCAUUGUGUJUGCUC	1128	4915	AAUUGCAUUGUGUJUGCUC	1128	4937	GAGCAAAACAAUUGCAUUC	1452
4933	CUGGUGGAGGUGGGCAUGG	1129	4933	CUGGUGGAGGUGGGCAUGG	1129	4955	CCAUGCCCCACCUCCACCAG	1453
4951	GGGUCUGUUCUGAAAUGUA	1130	4951	GGGUCUGUUCUGAAAUGUA	1130	4973	UACAUIUCAGAAACAGACCC	1454
4969	AAAGGGUUCAGACGGGGUUC	1131	4969	AAAGGGUUCAGACGGGGUUC	1131	4991	AACCCCGUUCUGAACCCUUCU	1455
4987	UUCUGGUUUUJAGAAAGGUUG	1132	4987	UUCUGGUUUUJAGAAAGGUUG	1132	5009	CAACCUUCUAAAACCCAGAA	1456
5005	GCGUUGUUCUUCGAGUUGGG	1133	5005	GCGUUGUUCUUCGAGUUGGG	1133	5027	CCCAACUUCGAAAGAACACGC	1457
5023	GCJAAAGUAGAGUUCGUUG	1134	5023	GCJAAAGUAGAGUUCGUUG	1134	5045	CAACGAACUCUACUUAAGC	1458
5041	GUGCUGUUUCUGACUCCUA	1135	5041	GUGCUGUUUCUGACUCCUA	1135	5063	UAGGAGUCAGAAACAGCAC	1459
5059	AAUGAGAGUUCUCCUCCAGA	1136	5059	AAUGAGAGUUCUCCUCCAGA	1136	5081	UCUGGAAGGAACUCUCUUAU	1460

5077	ACCGUAGCUGUCUCCUUG	1137	5077	ACCGUAGCUGUCUCCUUG	1137	5099	CAAGGAGACGCUAACGGU	1461
5095	GCCAAAGCCCAGGAAGAAA	1138	5095	GCCAAAGCCCAGGAAGAAA	1138	5117	UUUUCUCCUGGGCUUGGC	1462
5113	AAUGAUGCAGCUCUGGCUC	1139	5113	AAUGAUGCAGCUCUGGCUC	1139	5135	GAGCCAGAGCUGCAUCAUU	1463
5131	CCUUGUCUCCAGGCUGAU	1140	5131	CCUUGUCUCCAGGCUGAU	1140	5153	AUCAGCCUGGGAGACAAGG	1464
5149	UCCUUUUUUCAGAAUACCA	1141	5149	UCCUUUUUUCAGAAUACCA	1141	5171	UGGUUUUCUGAAUAAAGGA	1465
5167	ACAAAGAAAGGACAUUCAG	1142	5167	ACAAAGAAAGGACAUUCAG	1142	5189	CUGAAUGUCCUUUCUUUGU	1466
5185	GCUCAAGGCUCCUGCCGU	1143	5185	GCUCAAGGCUCCUGCCGU	1143	5207	ACGGCAGGAGCCUUGAGC	1467
5203	UGUUGAAGAGUUCUGACUG	1144	5203	UGUUGAAGAGUUCUGACUG	1144	5225	CAGUCAGAACUCUUCACAAC	1468
5221	GCACAAACCAGCUUCUGGU	1145	5221	GCACAAACCAGCUUCUGGU	1145	5243	ACCAGAAAGCUGGUUUGUC	1469
5239	UUUCUUCUGGAUUAUAC	1146	5239	UUUCUUCUGGAUUAUAC	1146	5261	GUUUCAUUCAGAAAGAAA	1470
5257	CCCUCAUAUCUGUCCUGAU	1147	5257	CCCUCAUAUCUGUCCUGAU	1147	5279	AUCAGGACAGAUUAUGAGG	1471
5275	UGUGAUUAUGUCUGACUG	1148	5275	UGUGAUUAUGUCUGACUG	1148	5297	CAGUCUCAGACAUUAUCACA	1472
5293	GAUUGCGGGAGGUUCAAUG	1149	5293	GAUUGCGGGAGGUUCAAUG	1149	5315	CAUUGAACCUCCCGCAUUC	1473
5311	GUGAAGCUGUGUGUGUGU	1150	5311	GUGAAGCUGUGUGUGUGU	1150	5333	ACACCACACACAGCUUCAC	1474
5329	UCAAAGUUUCAGGAAGGAU	1151	5329	UCAAAGUUUCAGGAAGGAU	1151	5351	AUCCUUCUUGAAACUUGA	1475
5347	UUUUACCCUUUUGUUCUUC	1152	5347	UUUUACCCUUUUGUUCUUC	1152	5369	GAAGAACAACAAAGGGUAAAA	1476
5365	CCCCUGUCCCCAACCCAC	1153	5365	CCCCUGUCCCCAACCCAC	1153	5387	GUGGGUUGGGGACAGGGGG	1477
5383	CUCUCACCCCGCAACCCAU	1154	5383	CUCUCACCCCGCAACCCAU	1154	5405	AUGGGUUGCGGGGUGAGAG	1478
5401	UCAGUAUUUUAAGUUAUUUG	1155	5401	UCAGUAUUUUAAGUUAUUUG	1155	5423	CAAAUAACUAAAAUACUGA	1479
5419	GGCCUCUACUCCAGUAAAC	1156	5419	GGCCUCUACUCCAGUAAAC	1156	5441	GUUUACUGGAGUAGAGGCC	1480
5437	CCUGAUUGGGUUUGUUCAC	1157	5437	CCUGAUUGGGUUUGUUCAC	1157	5459	GUGAACAAACCCCAUUCAGG	1481
5455	CUCUCUGAAUGAUUAUUAG	1158	5455	CUCUCUGAAUGAUUAUUAG	1158	5477	CUAAUAUUAUUAUUCAGAGAG	1482
5473	GCCAGACUUCAAAAUUUAU	1159	5473	GCCAGACUUCAAAAUUUAU	1159	5495	AAUAAUUUUUGAAGUCUGGC	1483
5491	UUUAUAGCCCAAAUUUAUA	1160	5491	UUUAUAGCCCAAAUUUAUA	1160	5513	UUUAUUUUUGGGCUAUAAA	1484
5509	ACAUCUUAUUGAUUAUUUA	1161	5509	ACAUCUUAUUGAUUAUUUA	1161	5531	UAAUAUAUUAUUAUAGUUG	1485
5527	AGACUUUUUAACAUUAUAG	1162	5527	AGACUUUUUAACAUUAUAG	1162	5549	CUCUAUAUGUUAAAAAGUCU	1486
5545	GCUAUUUUCUACUGAUUUUU	1163	5545	GCUAUUUUCUACUGAUUUUU	1163	5567	AAAAUUCAGUAGAAAUAGC	1487
5563	UGCCCUUUGUUCUGUCCUUU	1164	5563	UGCCCUUUGUUCUGUCCUUU	1164	5585	AAAGGACAGAACAAAGGGCA	1488
5581	UUUUUCAAAAAAGAAAUUG	1165	5581	UUUUUCAAAAAAGAAAUUG	1165	5603	CAUUUCUUUUUUUGAAAAAA	1489
5599	GUGUUUUUUUGUUUGGUACC	1166	5599	GUGUUUUUUUGUUUGGUACC	1166	5621	GGUACCAAAACAAAAACAC	1490
5617	CAUAGUGUGAAUUGCUGGG	1167	5617	CAUAGUGUGAAUUGCUGGG	1167	5639	CCCAGCAUUUCACACUAG	1491
5635	GAACAAUGACUAUAAGACA	1168	5635	GAACAAUGACUAUAAGACA	1168	5657	UGUCUUUAUAGUCAUUGUUC	1492
5653	AUGCAUUGGCACAUUAUU	1169	5653	AUGCAUUGGCACAUUAUU	1169	5675	AAUAUAUUGGCCAUJAGCAU	1493
5671	UUUAUGUCUUUUUAUGUAG	1170	5671	UUUAUGUCUUUUUAUGUAG	1170	5693	CUACAUAACACAGACUAA	1494
5689	GAACAAAUUGUAAUAUUU	1171	5689	GAACAAAUUGUAAUAUUU	1171	5711	AAUAUAUUAUUAUUGUUC	1495
5707	UAAAGCCUUUAUAUAUUAUG	1172	5707	UAAAGCCUUUAUAUAUUAUG	1172	5729	CAUUAUAUAUAAGGCUUUA	1496

5725	GAACUUUGUACUAUUCACA	1173	5725	GAACUUUGUACUAUUCACA	1173	5747	UGUGAAUAGUACAAAGUUC	1497
5743	AUUUUGUAUCAGUAUUUUG	1174	5743	AUUUUGUAUCAGUAUUUUG	1174	5765	CAUAAUACUGAUACAAAU	1498
5761	GUAGCAUAACAAAGGUCAU	1175	5761	GUAGCAUAACAAAGGUCAU	1175	5783	AUGACCUUUUGUUAUGCUAC	1499
5779	UAAUGCUUUCAGCAAUUGA	1176	5779	UAAUGCUUUCAGCAAUUGA	1176	5801	UCAAUUGCUGAGAAAGCAUUA	1500
5797	AUGUCAUUUUUAUUAAGAA	1177	5797	AUGUCAUUUUUAUUAAGAA	1177	5819	UUCUUUAUUAUUAUUAAGAA	1501
5812	AGAACAUUGAAAAACUUGA	1178	5812	AGAACAUUGAAAAACUUGA	1178	5834	UCAAGUUUUUCAAUGUUCU	1502

gi|4503752|ref|NM_002020.1

Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPos	Lower seq	Seq ID
1	ACCCACGGCAGCGGCCGG	1503	1	ACCCACGGCAGCGGCCGG	1503	23	CCGGCCGCUGCCGUGGGU	1750
19	GAGAUAGCAGGGGCCGG	1504	19	GAGAUAGCAGGGGCCGG	1504	41	CGGCGCCCGCUGCAUCUC	1751
37	GCGCUGUGCCUGCGACUGU	1505	37	GCGCUGUGCCUGCGACUGU	1505	59	ACAGUCGACGACAGCGC	1752
55	UGGCUUGGCGGACUCC	1506	55	UGGCUUGGCGGACUCC	1506	77	GGAGUCCAGGACAGGCCA	1753
73	CUGGACGGCCUGGUGAGUG	1507	73	CUGGACGGCCUGGUGAGUG	1507	95	CACUCACAGGCGGUCCAG	1754
91	GACUACUCCAUAGACCCCG	1508	91	GACUACUCCAUAGACCCCG	1508	113	GGGGGUGCAUGGAGUAGUC	1755
109	CCGACCUUGAACAUACACGG	1509	109	CCGACCUUGAACAUACACGG	1509	131	CCGUGAUGUUCAAAGGUCGG	1756
127	GAGGAGUCACAGCUAUCG	1510	127	GAGGAGUCACAGCUAUCG	1510	149	CGAUGACGUGUGACUCCUC	1757
145	GACACCGGUGACAGCCUGU	1511	145	GACACCGGUGACAGCCUGU	1511	167	ACAGGCGUGACACCGGUGUC	1758
163	UCCAUCUCCUGCAGGGGAC	1512	163	UCCAUCUCCUGCAGGGGAC	1512	185	GUCCCCUGCAGGAGUUGA	1759
181	CAGCACCCUCCAGUGGG	1513	181	CAGCACCCUCCAGUGGG	1513	203	CCACUCCAGGGGGUGCUG	1760
199	GCUUGGCCAGGAGCUCAGG	1514	199	GCUUGGCCAGGAGCUCAGG	1514	221	CCUGAGCUCCUGGCCAAGC	1761
217	GAGGCGCCAGCCACCGGAG	1515	217	GAGGCGCCAGCCACCGGAG	1515	239	CUCCGGUGGCGUGGCCCUJC	1762
235	GACAAGGACAGCGAGGACA	1516	235	GACAAGGACAGCGAGGACA	1516	257	UGUCCUGCUGUCCUUGUC	1763
253	ACGGGGUGGUGCGAGACU	1517	253	ACGGGGUGGUGCGAGACU	1517	275	AGUCUCGCACACCCCGU	1764
271	UGCGAGGGACAGACGCCA	1518	271	UGCGAGGGACAGACGCCA	1518	293	UGGCGUCUGUGCCCUCCGCA	1765
289	AGCCCCUACUGCAAGGUGU	1519	289	AGGCCUACUGCAAGGUGU	1519	311	ACACCUUGCAGUAGGGCCU	1766
307	UUGCUGCUGCAGGAGUAC	1520	307	UUGCUGCUGCAGGAGUAC	1520	329	GUACCUCUGCAGCAGCAA	1767
325	CAUGCCAACGACAGGCA	1521	325	CAUGCCAACGACAGGCA	1521	347	UGCCUGUGUCGUUGGCAUG	1768
343	AGCUACGUCUGCUACUACA	1522	343	AGCUACGUCUGCUACUACA	1522	365	UGUAGUAGCAGACGUAGCU	1769
361	AAGUACAUAAGGACGCA	1523	361	AAGUACAUAAGGACGCA	1523	383	UGCGUGCCUUGAGUACUU	1770
379	AUCGAGGGCACACGGCCG	1524	379	AUCGAGGGCACACGGCCG	1524	401	CGGCGGUGGUGCCCUCCAU	1771
397	GCCAGCUCCUACGUGUUCG	1525	397	GCCAGCUCCUACGUGUUCG	1525	419	CGAACACGUAGGAGCUGGC	1772

415	GUGAGAGACUUUGAGCAGC	1526	415	GUGAGAGACUUUGAGCAGC	1526	437	GCUGCUCAAAAGUCUCAC	1773
433	CCAUUCAUCAACAAGCCUG	1527	433	CCAUUCAUCAACAAGCCUG	1527	455	CAGGCUUUGUUGAUGAUGG	1774
451	GACACGCUCUUGGUCAACA	1528	451	GACACGCUCUUGGUCAACA	1528	473	UGUUGACCAAGAGCGUGUC	1775
469	AGGAAGGACGCCAUGUGGG	1529	469	AGGAAGGACGCCAUGUGGG	1529	491	CCCACAUUGGCGUCCUCCU	1776
487	GUGCCCUUGUCUGGUCCA	1530	487	GUGCCCUUGUCUGGUCCA	1530	509	UGACACACAGACAGGGCAC	1777
505	AUCCCCGGCCUCAAUGUCA	1531	505	AUCCCCGGCCUCAAUGUCA	1531	527	UGACAUUGAGGCGCGGGGAU	1778
523	ACGUGGCGUCGCAAGCU	1532	523	ACGUGGCGUCGCAAGCU	1532	545	AGCUUUGCGAGCGCAGCGU	1779
541	UCGGUGCUGUGGCCAGAGC	1533	541	UCGGUGCUGUGGCCAGAGC	1533	563	CGUCUGGCCACAGCACCGA	1780
559	GGCAGGAGUGGUGUGGG	1534	559	GGCAGGAGUGGUGUGGG	1534	581	CCACACACCCUCCUGCCC	1781
577	GAUGACCGCGGGGCAUGC	1535	577	GAUGACCGCGGGGCAUGC	1535	599	GCAUGCCCCCGCGGUAUC	1782
595	CUCGUGUCCACGCCACUGC	1536	595	CUCGUGUCCACGCCACUGC	1536	617	GCAGUGCGUGGACACGAG	1783
613	CUGCACGAUGCCCUGUACC	1537	613	CUGCACGAUGCCCUGUACC	1537	635	GGUACAGGGCAUCGUGCAG	1784
631	CUGCAGUGCGAGACCACCU	1538	631	CUGCAGUGCGAGACCACCU	1538	653	AGGUGGUCUCGCACUGCAG	1785
649	UGGGAGACCAAGGACUCC	1539	649	UGGGAGACCAAGGACUCC	1539	671	GGAAGUCCUGGUCUCCCA	1786
667	CUUCCAAACCCUCCUGG	1540	667	CUUCCAAACCCUCCUGG	1540	689	CCAGGAAGGGGUUGGAAAG	1787
685	GUGCACAUCACAGGCAACG	1541	685	GUGCACAUCACAGGCAACG	1541	707	CGUUGCCUGUGAUGUGCAG	1788
703	GAGCUCUAUGACAUCACG	1542	703	GAGCUCUAUGACAUCACG	1542	725	GCUGGAUGUCAUAGAGCUC	1789
721	CUGUUGCCCAGGAAGUCGC	1543	721	CUGUUGCCCAGGAAGUCGC	1543	743	GCAGCUUCCUGGGCAACAG	1790
739	CUGGAGCUGCUGGUAGGGG	1544	739	CUGGAGCUGCUGGUAGGGG	1544	761	CCCCUACCAAGCAGCUCAG	1791
757	GAGAAGCUGGUCUCAACU	1545	757	GAGAAGCUGGUCUCAACU	1545	779	AGUUGAGGACCAAGCUUCUC	1792
775	UGCACCGUGUGGGCUGAGU	1546	775	UGCACCGUGUGGGCUGAGU	1546	797	ACUCAGCCCCACACGGUGCA	1793
793	UUUAACUCAGGUGUCACCU	1547	793	UUUAACUCAGGUGUCACCU	1547	815	AGGUGACACCCUGAGUUAAA	1794
811	UUUGACUGGGACUACCCAG	1548	811	UUUGACUGGGACUACCCAG	1548	833	CUGGGUAGUCCCAAGUAAA	1795
829	GGGAAGCAGGCAGAGCGGG	1549	829	GGGAAGCAGGCAGAGCGGG	1549	851	CCCGCUCUGCCUGCUUCCC	1796
847	GGUAAGUGGGUGCCCGAGC	1550	847	GGUAAGUGGGUGCCCGAGC	1550	869	GCUCGGGACCCACUUAAC	1797
865	CGACGCUCCCAACAGACCC	1551	865	CGACGCUCCCAACAGACCC	1551	887	GGGUCUGUUGGGAGCGUCG	1798
883	CACACAGAACUCUCCAGCA	1552	883	CACACAGAACUCUCCAGCA	1552	905	UGCUGGAGAGUUCUGUGUG	1799
901	AUCCUGACCAUCCACAACG	1553	901	AUCCUGACCAUCCACAACG	1553	923	CGUUGUGGAUGGUCAGGAU	1800
919	GUCAGCCAGCAGCAGCCUGG	1554	919	GUCAGCCAGCAGCAGCCUGG	1554	941	CCAGGUCGUGGUGGCGUGAC	1801
937	GGCUCGUUUGUGUGCAAGG	1555	937	GGCUCGUUUGUGUGCAAGG	1555	959	CCUUGCACACAUACGAGCC	1802
955	GCCAAACAACGGCAUCCAGC	1556	955	GCCAAACAACGGCAUCCAGC	1556	977	GCUGGAUGCCGUGUUGGCG	1803
973	CGAUUUCGGGAGAGACCCG	1557	973	CGAUUUCGGGAGAGACCCG	1557	995	CGGUGCUCUCCCGAAUCCG	1804
991	GAGGUCAUUGUGCAUGAAA	1558	991	GAGGUCAUUGUGCAUGAAA	1558	1013	UUUCAUGCACAAUAGACCUC	1805

1009	AAUCCCUUUAUCAGCGUG	1559	1009	AAUCCCUUUAUCAGCGUG	1559	1031	CGACGCUGAUGAAGGGAUU	1806
1027	GAGUGGCUCUAAAGGACCCA	1560	1027	GAGUGGCUCUAAAGGACCCA	1560	1049	UGGGUCCUUUAGAGCCACUC	1807
1045	AUCCUGGAGGCCACGGCAG	1561	1045	AUCCUGGAGGCCACGGCAG	1561	1067	CUGCCGUGGCCUCCAGGAU	1808
1063	GGAGACGAGCUGGUGAAGC	1562	1063	GGAGACGAGCUGGUGAAGC	1562	1085	GCUUACACGAGCUGGUCUC	1809
1081	CUGCCCGUGAAGCUGGCAG	1563	1081	CUGCCCGUGAAGCUGGCAG	1563	1103	CUGCCAGCUUACAGGGCAG	1810
1099	GCGUACCCCCCGCCCGAGU	1564	1099	GCGUACCCCCCGCCCGAGU	1564	1121	ACUGGGCGGGGGGUAACGC	1811
1117	UUCACGUGGUACAAGGAUG	1565	1117	UUCACGUGGUACAAGGAUG	1565	1139	CAUCCUUGUACCAACUGGAA	1812
1135	GGAAAGGCACUUGCCGGGC	1566	1135	GGAAAGGCACUUGCCGGGC	1566	1157	GCCCGGACAGUGCCUUUCC	1813
1153	CGCCACAGUCCACAUGCC	1567	1153	CGCCACAGUCCACAUGCC	1567	1175	GGCAUGUGGACUGUGGCG	1814
1171	CUGGUGCUCAAGGAGGUGA	1568	1171	CUGGUGCUCAAGGAGGUGA	1568	1193	UCACCUCCUUGAGCACCCAG	1815
1189	ACAGAGGCCAGCACAGGCA	1569	1189	ACAGAGGCCAGCACAGGCA	1569	1211	UGCCUGUGCUGGCCUCUGU	1816
1207	ACCUACACCCUCGCCUCUGU	1570	1207	ACCUACACCCUCGCCUCUGU	1570	1229	ACAGGGCGAGGGUGUAGGU	1817
1225	UGGAACUCCGCGUCUGGCC	1571	1225	UGGAACUCCGCGUCUGGCC	1571	1247	GGCCAGCAGCGGAGUCCA	1818
1243	CUGAGGCGCAACAUCAGCC	1572	1243	CUGAGGCGCAACAUCAGCC	1572	1265	GGCUGAUGUUGCGCCUCAG	1819
1261	CUGGAGCUGGUGUGAAUG	1573	1261	CUGGAGCUGGUGUGAAUG	1573	1283	CAUUCACCAACAGCUCAC	1820
1279	GUGCCCCCCCAGAUACAUG	1574	1279	GUGCCCCCCCAGAUACAUG	1574	1301	CAUGUAUCUGGGGGGCAC	1821
1297	GAGAGGAGGCCUCCUCC	1575	1297	GAGAGGAGGCCUCCUCC	1575	1319	GGGAGGAGGCCUCCUCCUC	1822
1315	CCCAGCAUCUACUCGCGUC	1576	1315	CCCAGCAUCUACUCGCGUC	1576	1337	GACGCGAGUAGUAGCUGGG	1823
1333	CACAGCCGCCAGGCCUUA	1577	1333	CACAGCCGCCAGGCCUUA	1577	1355	UGAGGGCCUGGGCGGUGUG	1824
1351	ACCUGCACGGCCUACGGGG	1578	1351	ACCUGCACGGCCUACGGGG	1578	1373	CCCCGUAGGCCGUGCAGGU	1825
1369	GUGCCCCUGCCUUCACAGA	1579	1369	GUGCCCCUGCCUUCACAGA	1579	1391	UGCUGAGAGGCGAGGGGCAC	1826
1387	AUCCAGUGGCACUGGCGGC	1580	1387	AUCCAGUGGCACUGGCGGC	1580	1409	GCCGCCAGUGCCACUGGAU	1827
1405	CCCUGGACACCCUGCAAGA	1581	1405	CCCUGGACACCCUGCAAGA	1581	1427	UCUUGCAGGGUGUCCAGGG	1828
1423	AUGUUUGCCCCAGCGUAGUC	1582	1423	AUGUUUGCCCCAGCGUAGUC	1582	1445	GACUACGCUGGGCAACAACU	1829
1441	CUCCGGCGGCGGCAGCAGC	1583	1441	CUCCGGCGGCGGCAGCAGC	1583	1463	GCUGCUGCCCGCCCGGAG	1830
1459	CAAGACCUCAUGCCACAGU	1584	1459	CAAGACCUCAUGCCACAGU	1584	1481	ACUGUGGCAUGAGGUCUUG	1831
1477	UGCCGUGACUGGAGGCGG	1585	1477	UGCCGUGACUGGAGGCGG	1585	1499	CCGCCUCCAGUACACGGCA	1832
1495	GUGACCACGCAGGAUGCCG	1586	1495	GUGACCACGCAGGAUGCCG	1586	1517	CGGCAUCCUGCGUGGUCAC	1833
1513	GUGAACCCCAUCGAGAGCC	1587	1513	GUGAACCCCAUCGAGAGCC	1587	1535	GGCUCUCGAUGGGGUACAC	1834
1531	CUGGACACCUUGACCCGAGU	1588	1531	CUGGACACCUUGACCCGAGU	1588	1553	ACUCGGUCCAGGUGUCCAG	1835
1549	UUUGUGGAGGGAAGAAUA	1589	1549	UUUGUGGAGGGAAGAAUA	1589	1571	UAUUCUUUCCUCCACAAA	1836
1567	AAGACUGUGAGCAAGCUGG	1590	1567	AAGACUGUGAGCAAGCUGG	1590	1589	CCAGCUUGCUCACAGUCUU	1837
1585	GUGAUCCAGAAUGCCCAACG	1591	1585	GUGAUCCAGAAUGCCCAACG	1591	1607	CGUUGGCAUUCUGGAUCAC	1838

1603	GUGUCUGCCAUUGUACAAGU	1592	1603	GUGUCUGCCAUUGUACAAGU	1592	1625	ACUUGUACAUUGGACACAC	1839
1621	UGUGUGGUCUCCAACAAGG	1593	1621	UGUGUGGUCUCCAACAAGG	1593	1643	CCUUGUUGGAGACACACA	1840
1639	GUGGGCCAGGAUGAGCGGC	1594	1639	GUGGGCCAGGAUGAGCGGC	1594	1661	GCCGCUCAUCCUGGCCAC	1841
1657	CUCAUCUACUUAUGUGA	1595	1657	CUCAUCUACUUAUGUGA	1595	1679	UCACUAAGAAGUAGUAG	1842
1675	ACCACCAUCCCGACGGCU	1596	1675	ACCACCAUCCCGACGGCU	1596	1697	AGCGUCGGGGAUGGUGU	1843
1693	UUCACCAUCGAUCCAAGC	1597	1693	UUCACCAUCGAUCCAAGC	1597	1715	GCUUGGAUUCGAUGGUGAA	1844
1711	CAUCCGAGGAGCUACUAG	1598	1711	CAUCCGAGGAGCUACUAG	1598	1733	CUAGUAGCUCCUCGGGUGG	1845
1729	GAGGGCCAGCCGGUGCUCC	1599	1729	GAGGGCCAGCCGGUGCUCC	1599	1751	GGAGCACCGGCGUGGCCUC	1846
1747	CUGAGCUGCCAAGCCGACA	1600	1747	CUGAGCUGCCAAGCCGACA	1600	1769	UGUCGGCUUGGCAGCUCAG	1847
1765	AGCUACAAGUACGAGCAUC	1601	1765	AGCUACAAGUACGAGCAUC	1601	1787	GAUCUCGUACUUGUAGCU	1848
1783	CUGCGCUGGUACCGCCUCA	1602	1783	CUGCGCUGGUACCGCCUCA	1602	1805	UGAGCGGUACCGCGCAG	1849
1801	AACUUGUCCAGCUGCAGC	1603	1801	AACUUGUCCAGCUGCAGC	1603	1823	CGUGAGCGUGGACAGGUU	1850
1819	GAUGCGACGGGAACCCGC	1604	1819	GAUGCGACGGGAACCCGC	1604	1841	GCGGUUCCCGUGCGCAUC	1851
1837	CUUCUGCUCGACUGCAAGA	1605	1837	CUUCUGCUCGACUGCAAGA	1605	1859	UCUUGCAGUCGAGCAGAAG	1852
1855	AACGUGCAUCUGUUCGCCA	1606	1855	AACGUGCAUCUGUUCGCCA	1606	1877	UGGCGAACAGAUCCACGUU	1853
1873	ACCCUCUGGCCGCCAGCC	1607	1873	ACCCUCUGGCCGCCAGCC	1607	1895	GGCUGCGGCCAGAGGGGU	1854
1891	CUGGAGGAGGUGGCACCCUG	1608	1891	CUGGAGGAGGUGGCACCCUG	1608	1913	CAGGUGCCACCUCCUCCAG	1855
1909	GGGGCGGCCACGCCACGC	1609	1909	GGGGCGGCCACGCCACGC	1609	1931	GCUGGGCUGGGCGGCCCC	1856
1927	CUCAGCCUGAGUAUCCCCC	1610	1927	CUCAGCCUGAGUAUCCCCC	1610	1949	GGGGUAUACUACAGGUGAG	1857
1945	CGCGUCGCGCCGAGCAGC	1611	1945	CGCGUCGCGCCGAGCAGC	1611	1967	CGUGCUCGGCGCGACGCG	1858
1963	GAGGGCCACUAUGUGCGG	1612	1963	GAGGGCCACUAUGUGCGG	1612	1985	CGCACAUAGUGGCCUCC	1859
1981	GAAGUGCAAGACCGCGCA	1613	1981	GAAGUGCAAGACCGCGCA	1613	2003	UGCGCCGUCUUGCACUUC	1860
1999	AGCAUGACAAGCACUGCC	1614	1999	AGCAUGACAAGCACUGCC	1614	2021	GGCAGUGCUUGUACUAGGCU	1861
2017	CACAAGAAGUACCUUGUCGG	1615	2017	CACAAGAAGUACCUUGUCGG	1615	2039	CCGACAGGUACUUCUUGUG	1862
2035	GUGCAGGCCCUUGGAAGCCC	1616	2035	GUGCAGGCCCUUGGAAGCCC	1616	2057	GGGCUCCAGGGGCGGAC	1863
2053	CCUCGGCUCACGCAGAACU	1617	2053	CCUCGGCUCACGCAGAACU	1617	2075	AGUUCUGCGUGAGCCGAGG	1864
2071	UUGACCGACCUCCUGGUGA	1618	2071	UUGACCGACCUCCUGGUGA	1618	2093	UCACCAGGAGGUGCGUCAA	1865
2089	AACGUGAGCGACUCGUGG	1619	2089	AACGUGAGCGACUCGUGG	1619	2111	CCAGCGAGUCGUCACGUU	1866
2107	GAGUAGCAGUCUUGUGG	1620	2107	GAGUAGCAGUCUUGUGG	1620	2129	CCACCAAGCACUGCAUCUC	1867
2125	GCCGGAGCGCACGCGCCCA	1621	2125	GCCGGAGCGCACGCGCCCA	1621	2147	UGGGCGGUGGCGUCCCGGC	1868
2143	AGCAUCGUGUGGUACAAAG	1622	2143	AGCAUCGUGUGGUACAAAG	1622	2165	CUUUGUACACACGAGUUCU	1869
2161	GACGAGAGGCGUGUGGAGG	1623	2161	GACGAGAGGCGUGUGGAGG	1623	2183	CCUCCAGCAGCCUCCGUC	1870
2179	GAAGAAGUCUGGAGUCGACU	1624	2179	GAAGAAGUCUGGAGUCGACU	1624	2201	AGUCGACUCCAGACUUCUUC	1871

2197	UUGCGGACUCCAAACCAGA	1625	2197	UUGCGGACUCCAAACCAGA	1625	2219	UCUGGUUGGAGUCCGCCAA	1872
2215	AAGCUGAGCAUCCAGCGCG	1626	2215	AAGCUGAGCAUCCAGCGCG	1626	2237	CGCGCUGGAUGCUAGCUU	1873
2233	GUGCGGAGGAGGAUGCGG	1627	2233	GUGCGGAGGAGGAUGCGG	1627	2255	CCGCAUCCUCCUCCGCGCAC	1874
2251	GGACCGUAUCUGCAGCG	1628	2251	GGACCGUAUCUGCAGCG	1628	2273	CGCUGCACAGAUACGGUCC	1875
2269	GUGUGCAGACCCAAAGGCU	1629	2269	GUGUGCAGACCCAAAGGCU	1629	2291	AGCCCUUGGGUCUGCACAC	1876
2287	UGCGUCAAUCUCCGCCA	1630	2287	UGCGUCAAUCUCCGCCA	1630	2309	UGGCGGAGGAGUUGACGCA	1877
2305	AGCGUGCCGUGGAAGGCU	1631	2305	AGCGUGCCGUGGAAGGCU	1631	2327	AGCCUUCACAGGCCACGCU	1878
2323	UCCGAGGAUAGGCGAGCA	1632	2323	UCCGAGGAUAGGCGAGCA	1632	2345	UGCUGCCCUUUAUCCUCGGA	1879
2341	AUGGAGAUCCGUAUCCUUG	1633	2341	AUGGAGAUCCGUAUCCUUG	1633	2363	CAAGGAUCACGAUCCCAU	1880
2359	GUCGGUACCGGCGUCAUCG	1634	2359	GUCGGUACCGGCGUCAUCG	1634	2381	CGAUGACGCCGGUACCGAC	1881
2377	GCUGUCUUCUUGGGUCC	1635	2377	GCUGUCUUCUUGGGUCC	1635	2399	GGACCCAGAAAGACAGC	1882
2395	CUCUCCUCCUCAUCUUCU	1636	2395	CUCUCCUCCUCAUCUUCU	1636	2417	AGAAGAUGAGGAGGAGGAG	1883
2413	UGUAACAUGAGGAGGCCGG	1637	2413	UGUAACAUGAGGAGGCCGG	1637	2435	CCGGCCUCCUCAUGUUAACA	1884
2431	GCCACGCGAGACAUAAGA	1638	2431	GCCACGCGAGACAUAAGA	1638	2453	UCUUGAUGUCUGCGUGGGC	1885
2449	ACGGGCUACCUGUCCAUA	1639	2449	ACGGGCUACCUGUCCAUA	1639	2471	UGAUGGACAGGUAGCCCGU	1886
2467	AUCAUGACCCCGGGAGG	1640	2467	AUCAUGACCCCGGGAGG	1640	2489	CCUCCCGGGGUGCCAUGAU	1887
2485	GUGCCUUGGAGGAGCAAU	1641	2485	GUGCCUUGGAGGAGCAAU	1641	2507	AUUGCUCCUCCAGAGGCAC	1888
2503	UGCGAAUACCUGUCCUACG	1642	2503	UGCGAAUACCUGUCCUACG	1642	2525	CGUAGGACAGGUUAUCCGA	1889
2521	GAUGCCAGCCAGUGGGAAU	1643	2521	GAUGCCAGCCAGUGGGAAU	1643	2543	AUUCACACUGGCGGCAUC	1890
2539	UUCGCGGAGAGCGGCUUC	1644	2539	UUCGCGGAGAGCGGCUUC	1644	2561	GCAGCCGCUUCGGGGGAA	1891
2557	CACCUGGGAGAGUGCUUC	1645	2557	CACCUGGGAGAGUGCUUC	1645	2579	CGAGCACUCUCCCCAGGUG	1892
2575	GGCUACGGGCGCCUUCGGGA	1646	2575	GGCUACGGGCGCCUUCGGGA	1646	2597	UCCCGAAGGCGCGCGUAGCC	1893
2593	AAGGUGGUGGAAGCCUCCG	1647	2593	AAGGUGGUGGAAGCCUCCG	1647	2615	CGGAGGCUUCCACCAACUU	1894
2611	GCUUUCGGCAUCCACAAGG	1648	2611	GCUUUCGGCAUCCACAAGG	1648	2633	CCUUGUGGAUCCCGAAAGC	1895
2629	GGCAGCAGCUUGACACCG	1649	2629	GGCAGCAGCUUGACACCG	1649	2651	CGGUGUCACAGCUUGGCC	1896
2647	GUGCCCGUAAAAUUCUGA	1650	2647	GUGCCCGUAAAAUUCUGA	1650	2669	UCAGCAUUUUCACGGCCAC	1897
2665	AAAGAGGGCGCCACGGCCA	1651	2665	AAAGAGGGCGCCACGGCCA	1651	2687	UGGCCGUGGGCGCCCUUUU	1898
2683	AGCAGCAGCGCGGCUUGA	1652	2683	AGCAGCAGCGCGGCUUGA	1652	2705	UCAGCGCGCGCUUGCGCU	1899
2701	AUGUCGGAGCUCAAGAUCC	1653	2701	AUGUCGGAGCUCAAGAUCC	1653	2723	GGAUUUUGAGCUCCGACAU	1900
2719	CUCAUUCACAUCCGCAACC	1654	2719	CUCAUUCACAUCCGCAACC	1654	2741	GGUUGCCGAUGUGAAUGAG	1901
2737	CACCUCAACGUGGUCAACC	1655	2737	CACCUCAACGUGGUCAACC	1655	2759	GGUUGACCACGUUGAGGUG	1902
2755	CUCUUGGGGCGUGCACCA	1656	2755	CUCUUGGGGCGUGCACCA	1656	2777	UGGUGCACGCCCGGAGGAG	1903
2773	AAGCGCAGGGGCCCCCUCA	1657	2773	AAGCGCAGGGGCCCCCUCA	1657	2795	UGAGGGGGCCCCUGCGGCUU	1904

2791	AUGGUGAUCGUGGAGUUCU	1658	2791	AUGGUGAUCGUGGAGUUCU	1658	2813	1905
2809	UGCAAGUACGGCAACCUUCU	1659	2809	UGCAAGUACGGCAACCUUCU	1659	2831	1906
2827	UCCAACUUCUGCGGCCA	1660	2827	UCCAACUUCUGCGGCCA	1660	2849	1907
2845	AAGCGGACGCCUUCAGCC	1661	2845	AAGCGGACGCCUUCAGCC	1661	2867	1908
2863	CCUUGCGGAGAAAGUUC	1662	2863	CCUUGCGGAGAAAGUUC	1662	2885	1909
2881	CCGAGCAGCGCGGACGCU	1663	2881	CCGAGCAGCGCGGACGCU	1663	2903	1910
2899	UUCGCGCCAUUGGAGC	1664	2899	UUCGCGCCAUUGGAGC	1664	2921	1911
2917	CUCGCCAGGCUUGGAUCGGA	1665	2917	CUCGCCAGGCUUGGAUCGGA	1665	2939	1912
2935	AGCGGCCGCGGAGCAGCG	1666	2935	AGCGGCCGCGGAGCAGCG	1666	2957	1913
2953	GACAGGGUCCUUCGCGC	1667	2953	GACAGGGUCCUUCGCGC	1667	2975	1914
2971	CGGUUCUGAAGACCGAGG	1668	2971	CGGUUCUGAAGACCGAGG	1668	2993	1915
2989	GGCGGAGCGAGCGGGCUU	1669	2989	GGCGGAGCGAGCGGGCUU	1669	3011	1916
3007	UCUCCAGACCAAGAAAGCUG	1670	3007	UCUCCAGACCAAGAAAGCUG	1670	3029	1917
3025	GAGGACCUUGGCUAGGCC	1671	3025	GAGGACCUUGGCUAGGCC	1671	3047	1918
3043	CCGUGACCAUGGAAGAU	1672	3043	CCGUGACCAUGGAAGAU	1672	3065	1919
3061	CUUGUCUACAGCUUCC	1673	3061	CUUGUCUACAGCUUCC	1673	3083	1920
3079	CAGGUGGCCAGAGGAUGG	1674	3079	CAGGUGGCCAGAGGAUGG	1674	3101	1921
3097	GAGUUCUGGCUUCCCGAA	1675	3097	GAGUUCUGGCUUCCCGAA	1675	3119	1922
3115	AAGUGCAUCCACAGAGACC	1676	3115	AAGUGCAUCCACAGAGACC	1676	3137	1923
3133	CUGGCUUCGGAACAUUC	1677	3133	CUGGCUUCGGAACAUUC	1677	3155	1924
3151	CUGCUGUCGGAAGCGACG	1678	3151	CUGCUGUCGGAAGCGACG	1678	3173	1925
3169	GUGGUGAAGAUUCUGACU	1679	3169	GUGGUGAAGAUUCUGACU	1679	3191	1926
3187	UUUGGCCUUGCCCGGACA	1680	3187	UUUGGCCUUGCCCGGACA	1680	3209	1927
3205	AUCUACAAAGACCCCGACU	1681	3205	AUCUACAAAGACCCCGACU	1681	3227	1928
3223	UACGUCGCAAGGGCAGUG	1682	3223	UACGUCGCAAGGGCAGUG	1682	3245	1929
3241	GCCCGGCUGCCCCUGAAGU	1683	3241	GCCCGGCUGCCCCUGAAGU	1683	3263	1930
3259	UGGAUGGCCCCUGAAAGCA	1684	3259	UGGAUGGCCCCUGAAAGCA	1684	3281	1931
3277	AUCUUCGACAAGGUGUACA	1685	3277	AUCUUCGACAAGGUGUACA	1685	3299	1932
3295	ACCAGCAGAGUGACGUGU	1686	3295	ACCAGCAGAGUGACGUGU	1686	3317	1933
3313	UGGUCCUUUGGGGUGCUUC	1687	3313	UGGUCCUUUGGGGUGCUUC	1687	3335	1934
3331	CUCUGGGAGAUUCUUCUC	1688	3331	CUCUGGGAGAUUCUUCUC	1688	3353	1935
3349	CUGGGGCCUCCCCGUACC	1689	3349	CUGGGGCCUCCCCGUACC	1689	3371	1936
3367	CCUGGGGUGCAGAUCAAUG	1690	3367	CCUGGGGUGCAGAUCAAUG	1690	3389	1937

3385	GAGGAGUUCUGCCAGCGCG	1691	3385	GAGGAGUUCUGCCAGCGCG	1691	3407	CGGCGUGGCGAAGACUCCUC	1938
3403	GUGAGAGACGGCACAAAGGA	1692	3403	GUGAGAGACGGCACAAAGGA	1692	3425	UCCUUGUGCCGUCUCUCAC	1939
3421	AUGAGGGCCCCGGAGCUGG	1693	3421	AUGAGGGCCCCGGAGCUGG	1693	3443	CCAGCUCGCGGGGCCUCAU	1940
3439	GCCACUCCCGCCAUACGCC	1694	3439	GCCACUCCCGCCAUACGCC	1694	3461	GGCGUAUGGCGGGAGUGGC	1941
3457	CACAUCAUGCUGAACUCUCU	1695	3457	CACAUCAUGCUGAACUCUCU	1695	3479	AGCAGUUCAGCAUGAUGUG	1942
3475	UGGUCCGGAGACCCCAAGG	1696	3475	UGGUCCGGAGACCCCAAGG	1696	3497	CCUUGGGGUCUCCGGACCA	1943
3493	GCGAGACCUCAUUCUCGG	1697	3493	GCGAGACCUCAUUCUCGG	1697	3515	CCGAGAAUGCAGGUCUCGC	1944
3511	GACCUUGGAGAUCCUGG	1698	3511	GACCUUGGAGAUCCUGG	1698	3533	CCAGGAUCCACCAGGUC	1945
3529	GGGACCUUGCAGGCGCA	1699	3529	GGGACCUUGCAGGCGCA	1699	3551	UGCCUGGAGCAGGUUCCCC	1946
3547	AGGGCCUUGCAAGAGGAAG	1700	3547	AGGGCCUUGCAAGAGGAAG	1700	3569	CUUCCUCUUGCAGGCCCCU	1947
3565	GAGGAGGUCUCAGAGCC	1701	3565	GAGGAGGUCUCAGAGCC	1701	3587	GGCCCAUGCAGACCUCCUC	1948
3583	CCGCGCAGCUCUCAGAGCU	1702	3583	CCGCGCAGCUCUCAGAGCU	1702	3605	AGCUCUGAGAGCUCGCGG	1949
3601	UCAGAAGAGGGCAGCUUCU	1703	3601	UCAGAAGAGGGCAGCUUCU	1703	3623	AGAAGCUGCCCUUCUGA	1950
3619	UCGCAGGUGUCCACCAUGG	1704	3619	UCGCAGGUGUCCACCAUGG	1704	3641	CCAUGGUGGACACCUCCGA	1951
3637	GCCCUACACAUCCGCCAGG	1705	3637	GCCCUACACAUCCGCCAGG	1705	3659	CCUGGCGAUGUGUAGGCG	1952
3655	GCUGACGUCAGGACAGCC	1706	3655	GCUGACGUCAGGACAGCC	1706	3677	GGCUGUCCUACGCGUACG	1953
3673	CCGCCAAGCCUGCAGCGCC	1707	3673	CCGCCAAGCCUGCAGCGCC	1707	3695	GGCGCUGCAGGCUUGGCGG	1954
3691	CACAGCCUGGCCGCCAGGU	1708	3691	CACAGCCUGGCCGCCAGGU	1708	3713	ACCUUGCGGCGCAGGCGUG	1955
3709	UAUUACAACUGGGUGUCCU	1709	3709	UAUUACAACUGGGUGUCCU	1709	3731	AGGACACCCAGUUGUAUA	1956
3727	UUUCCCGGUGCCUGGCCA	1710	3727	UUUCCCGGUGCCUGGCCA	1710	3749	UGGCCAGGCACCCCGGAAA	1957
3745	AGAGGGGUGAGACCCGUG	1711	3745	AGAGGGGUGAGACCCGUG	1711	3767	CACGGGUCUCAGCCCCUCU	1958
3763	GGUUCUCCAGGAUGAAGA	1712	3763	GGUUCUCCAGGAUGAAGA	1712	3785	UCUUAUCCUGGAGGAACC	1959
3781	ACAUUUGAGGAUUUCCCCA	1713	3781	ACAUUUGAGGAUUUCCCCA	1713	3803	UGGGGAUUUCCUCAAUUGU	1960
3799	AUGACCCCAACGACCUACA	1714	3799	AUGACCCCAACGACCUACA	1714	3821	UGUAGGUCUUGGGGUCAU	1961
3817	AAAGGCUCUGUGGACAACC	1715	3817	AAAGGCUCUGUGGACAACC	1715	3839	GGUUGUCCACAGAGCCUUU	1962
3835	CAGACAGACAGUGGGAUGG	1716	3835	CAGACAGACAGUGGGAUGG	1716	3857	CCAUCCACUCUGUCUGUG	1963
3853	GUGCUGGCCUCGGAGGAGU	1717	3853	GUGCUGGCCUCGGAGGAGU	1717	3875	ACUCCUCCGAGGCCAGCAC	1964
3871	UUUGAGCAGAUAGAGCA	1718	3871	UUUGAGCAGAUAGAGCA	1718	3893	UGCUCUCUAUCUGCUCAAA	1965
3889	AGGCAUAGACAAGAAAGCG	1719	3889	AGGCAUAGACAAGAAAGCG	1719	3911	CGCUUUCUUGUCUAUGCCU	1966
3907	GGCUUCAGGUAGCUGAAGC	1720	3907	GGCUUCAGGUAGCUGAAGC	1720	3929	GCUUCAGCUACCUAGAGCC	1967
3925	CAGAGAGAGAGAGGCAGC	1721	3925	CAGAGAGAGAGAGGCAGC	1721	3947	GCUGCCUUCUCUCUCUCUG	1968
3943	CAUACGUCAGCAUUUUCUU	1722	3943	CAUACGUCAGCAUUUUCUU	1722	3965	AAGAAAUGCUGACGUUUG	1969
3961	UCUCUGCACUUUAUAAAGAA	1723	3961	UCUCUGCACUUUAUAAAGAA	1723	3983	UUUCUUUAUUAUGCAGAGA	1970

3979	AGAUCAAAAGACUUUAAGAC	1724	3979	AGAUCAAAAGACUUUAAGAC	1724	4001	GUCUUAAAGUCUUUGAUCU	1971
3997	CUUUCGCUAUUUCUUCUAC	1725	3997	CUUUCGCUAUUUCUUCUAC	1725	4019	GUAGAAAGAAUAGCGAAAG	1972
4015	CUGCUAUCUACUACAAACU	1726	4015	CUGCUAUCUACUACAAACU	1726	4037	AGUUUGUAGUAGAUAGCAG	1973
4033	UUCAAAAGAGGAACCAAGAG	1727	4033	UUCAAAAGAGGAACCAAGAG	1727	4055	CUCCUGGUUCCUUCUUGAA	1974
4051	GGACAAGAGGAGCAUGAAA	1728	4051	GGACAAGAGGAGCAUGAAA	1728	4073	UUUCAUGCUCCUUCUUGUCC	1975
4069	AGUGGACAAGGAGUGUGAC	1729	4069	AGUGGACAAGGAGUGUGAC	1729	4091	GUACACUCCUUGUCCACU	1976
4087	CCACUGAAGACACACAGGG	1730	4087	CCACUGAAGACACACAGGG	1730	4109	CCCUGUGGUGCUUCAGUGG	1977
4105	GAGGGUUUAGGCCUCCGGA	1731	4105	GAGGGUUUAGGCCUCCGGA	1731	4127	UCCGGAGGCCUAAACCCUC	1978
4123	AUGACUGCGGCGAGGCCUG	1732	4123	AUGACUGCGGCGAGGCCUG	1732	4145	CAGGCCUGCCCGCAGUCAU	1979
4141	GGAUAAUAUCCAGCCUCCC	1733	4141	GGAUAAUAUCCAGCCUCCC	1733	4163	GGAGGCGUGGAUAUUAUCC	1980
4159	CACAAGAGCUGGUGGAGC	1734	4159	CACAAGAGCUGGUGGAGC	1734	4181	GCUCCACCAGCUUCUUGUG	1981
4177	CAGAGUGUCCUUGACUCC	1735	4177	CAGAGUGUCCUUGACUCC	1735	4199	GGAGUCAGGGAACACUCUG	1982
4195	CUCCAAGGAAAGGGAGACG	1736	4195	CUCCAAGGAAAGGGAGACG	1736	4217	CGUCUCCUUCUUCUUGGAG	1983
4213	GCCUUUUAUGGUCUCUG	1737	4213	GCCUUUUAUGGUCUCUG	1737	4235	CAGCAGACCAUGAAAGGGC	1984
4231	GAGUAAACAGGUGCCUCCCC	1738	4231	GAGUAAACAGGUGCCUCCCC	1738	4253	GGGAAGGCACCUUGUACUC	1985
4249	CAGACACUGGCGUUAUCUG	1739	4249	CAGACACUGGCGUUAUCUG	1739	4271	GCAGUAAACGCCAGUGUCUG	1986
4267	CUUGACCAAAGAGCCCUCA	1740	4267	CUUGACCAAAGAGCCCUCA	1740	4289	UGAGGGCUUUGGUCUAAAG	1987
4285	AAGCGGCCCUUAUGCCAGC	1741	4285	AAGCGGCCCUUAUGCCAGC	1741	4307	GCUGSCAUAAAGGGCCGCUU	1988
4303	CGUGACAGAGGGCUCACCU	1742	4303	CGUGACAGAGGGCUCACCU	1742	4325	AGGUGAGCCCCUCUGACCG	1989
4321	UCUUGCCUUCUAGGUCACU	1743	4321	UCUUGCCUUCUAGGUCACU	1743	4343	AGUGACCUAAGAAAGGCAAGA	1990
4339	UUCUCACAAUGUCCCUUCA	1744	4339	UUCUCACAAUGUCCCUUCA	1744	4361	UGAAAGGACAUUGUGAGAA	1991
4357	AGCACCUAGCCUUGUGCCC	1745	4357	AGCACCUAGCCUUGUGCCC	1745	4379	GGGCACAGGGUCAGGUGCU	1992
4375	CGCCGAUUAUCCUUGGUA	1746	4375	CGCCGAUUAUCCUUGGUA	1746	4397	UACCAAGGAAUAAUCCGCG	1993
4393	AAUAUGAGUAAUACAUCAA	1747	4393	AAUAUGAGUAAUACAUCAA	1747	4415	UUGAUGUAUUACUCAUAUU	1994
4411	AAGAGUAGUAAUAAAAGCU	1748	4411	AAGAGUAGUAAUAAAAGCU	1748	4433	AGCUUUUAAUAAUACUACU	1995
4429	UAAUUAUUAUGUUAUUA	1749	4429	UAAUUAUUAUGUUAUUA	1749	4451	UUUAUAAACAUGAUUAUUA	1996

The 3'-ends of the Upper sequence and the Lower sequence of the siNA construct can include an overhang sequence, for example about 1, 2, 3, or 4 nucleotides in length, preferably 2 nucleotides in length, wherein the overhanging sequence of the lower sequence is optionally complementary to a portion of the target sequence. The upper sequence is also referred to as the sense strand, whereas the lower sequence is also referred to as the antisense strand. The upper and lower sequences in the Table can further comprise a chemical modification having Formulae I-VII or any combination thereof.

Table III: VEGF and VEGFR Synthetic Modified siNA constructs

VEGFR1

Target Pos	Target	Seq ID	Aliases	Sequence	Seq ID
296	GCUGUCUGCUUUCACACAGGAUCU	1997	FLT1:298U21 siRNA sense	UGUCUGCUUUCACACAGGAUTT	2020
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	FLT1:1956U21 siRNA sense	AGGAGAGGACCUUGAAACUGTT	2021
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	FLT1:1957U21 siRNA sense	GGAGAGGACCUUGAAACUGUTT	2022
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	FLT1:2787U21 siRNA sense	AUUUGGCAUUAAGAAAUACATT	2023
296	GCUGUCUGCUUUCACACAGGAUCU	1997	FLT1:316L21 siRNA (298C) antisense FLT1:1974L21 siRNA (1956C)	AUCCUGUGAGAAAGACGACATT	2024
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	antisense FLT1:1975L21 siRNA (1957C)	CAGUUUCAGGUCCUCUCUCCUTT	2025
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	antisense FLT1:2805L21 siRNA (2787C)	ACAGUUUCAGGUCCUCUCUCCCTT	2026
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	antisense	UGAUUUUCUUAUUGCCAAAUTT	2027
296	GCUGUCUGCUUUCACACAGGAUCU	1997	FLT1:298U21 siRNA stab04 sense	B uGucuGcuuucAcAGGAuTT B	2028
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	FLT1:1956U21 siRNA stab04 sense	B AGGAGAGGAGGACcuGAAAcuGTT B	2029
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	FLT1:1957U21 siRNA stab04 sense	B GGAGAGGAGGACcuGAAAcuGuTT B	2030
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	FLT1:2787U21 siRNA stab04 sense FLT1:316L21 siRNA (298C) stab05 antisense	B AuuuGGcAuuuAAGAAAUcATT B AuccuGuGAGAGGcAGAcATsT	2031 2032
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	antisense FLT1:1975L21 siRNA (1957C) stab05	cAGUuuAcAGGuccucuccuTsT	2033
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	antisense FLT1:1975L21 siRNA (1957C) stab05	AcAGuuuAcAGGuccucuccTsT	2034
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	antisense FLT1:2805L21 siRNA (2787C) stab05	uGAuuuucuuAAuGccAAAuTsT	2035
296	GCUGUCUGCUUUCACACAGGAUCU	1997	FLT1:298U21 siRNA stab07 sense	B uGucuGcuuucAcAGGAuTT B	2036
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	FLT1:1956U21 siRNA stab07 sense	B AGGAGAGGAGGACcuGAAAcuGTT B	2037
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	FLT1:1957U21 siRNA stab07 sense	B GGAGAGGAGGACcuGAAAcuGuTT B	2038
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	FLT1:2787U21 siRNA stab07 sense FLT1:316L21 siRNA (298C) stab11 antisense	B AuuuGGcAuuuAAGAAAUcATT B AuccuGuGAGAGGcAGAcATsT	2039 2040
1954	GAAGGAGAGGACCUUGAAACUGUC	1998	antisense FLT1:1975L21 siRNA (1957C) stab11	cAGUuuAcAGGuccucuccuTsT	2041
1955	AAGGAGAGGACCUUGAAACUGUCU	1999	antisense FLT1:2805L21 siRNA (2787C) stab11	AcAGUuuAcAGGuccucuccTsT	2042
2785	GCAUUUGGCAUUAAGAAAUACACC	2000	antisense	uGAuuuucuuAAuGccAAAuTsT	2043

VEGFR1

Target	SeqID	RPI#	Alias	Sequence	SeqID
AACUGAGUUUAAAAGGCACCCAG	2009	29694	FLT1:349U21 siRNA stab01 sense	CsUsGsAsGsUUUAAAAGGCACCCCTsT	2092
AACAACCCACAAAUAACAACAAGA	2010	29695	FLT1:2340U21 siRNA stab01 sense	CsAsAsCsCACAAAUAACAACAATsT	2093
AGCCUGGAAAGAAUCAAACCCUU	2011	29696	FLT1:3912U21 siRNA stab01 sense	CsCsUsGsGSAAGAAUCAAACCCTsT	2094
AAGCAAGGAGGGCCUCUGAUGGU	2012	29697	FLT1:2949U21 siRNA stab01 sense	GsCsAsAsGsGAGGGCCUCUGAUGTsT	2095
AACUGAGUUUAAAAGGCACCCAG	2009	29698	FLT1:369L21 siRNA (349C) stab01 sense	GsGsGsUsGsCCUUUUAAACUCAGTsT	2096
AACAACCCACAAAUAACAACAAGA	2010	29699	FLT1:2358L21 siRNA (2340C) stab01 sense	UsUsGsUsUsGUAUUUUUGUGGUUGTsT	2097
AGCCUGGAAAGAAUCAAACCCUU	2011	29700	FLT1:3932L21 siRNA (3912C) stab01 sense	GsGsUsUsUsUGAUUCUUUCCAGGTsT	2098
AAGCAAGGAGGGCCUCUGAUGGU	2012	29701	FLT1:2969L21 siRNA (2949C) stab01 sense	CsAsUsCsAsGAGGCCCUCCUUGCTsT	2099
AACUGAGUUUAAAAGGCACCCAG	2009	29702	FLT1:349U21 siRNA stab03 sense	csUsGsAsGuuuAAAAAGGcAcscscsTsT	2100
AACAACCCACAAAUAACAACAAGA	2010	29703	FLT1:2340U21 siRNA stab03 sense	csAsAscscAcAAAAuAcAAcsAsAsTsT	2101
AGCCUGGAAAGAAUCAAACCCUU	2011	29704	FLT1:3912U21 siRNA stab03 sense	cscsUsGsGAAAGAAuAAAAscscsTsT	2102
AAGCAAGGAGGGCCUCUGAUGGU	2012	29705	FLT1:2949U21 siRNA stab03 sense	GscsAsAsGGAGGGGccucUGAsusGsTsT	2103
AACUGAGUUUAAAAGGCACCCAG	2009	29706	FLT1:369L21 siRNA (349C) stab02 antisense	GsGsGsUsGsCsCsUsUsUsAsAsCsUsCsAsGsTsT	2104
AACAACCCACAAAUAACAACAAGA	2010	29707	FLT1:2358L21 siRNA (2340C) stab02 antisense	UsUsGsUsUsGsUsAsUsUsUsGsUsGsUsUsGsTsT	2105
AGCCUGGAAAGAAUCAAACCCUU	2011	29708	FLT1:3932L21 siRNA (3912C) stab02 antisense	GsGsUsUsUsGsAsUsUsCsUsUsCsCsAsGsGsTsT	2106
AAGCAAGGAGGGCCUCUGAUGGU	2012	29709	FLT1:2969L21 siRNA (2949C) stab02 antisense	CsAsUsCsAsGsAsGsCsCsCsUsCsCsUsUsGsCsTsT	2107
AACAACCCACAAAUAACAACAAGA	2010	29981	FLT1:2340U21 siRNA Native sense	CAACCACAAAUAACAACAAGA	2108
AACAACCCACAAAUAACAACAAGA	2010	29982	FLT1:2358L21 siRNA (2340C) Native antisense	UUGUUUAUUUUUGUGGUUGUU	2109
AACAACCCACAAAUAACAACAAGA	2010	29983	FLT1:2342U21 siRNA stab01 inv	AsAsCsAsAsCAUAAAAACACCAACTsT	2110
AACAACCCACAAAUAACAACAAGA	2010	29984	FLT1:2358L21 siRNA (2340C) stab01 inv	GsUsUsGsUGUUUUUAUGUUUUUTsT	2111
AACAACCCACAAAUAACAACAAGA	2010	29985	FLT1:2342U21 siRNA stab03 inv	AsAscAsAcAuAAAAAcAccAsAscTsT	2112
AACAACCCACAAAUAACAACAAGA	2010	29986	FLT1:2358L21 siRNA (2340C) stab02 inv	GsUsUsGsCsUsGsUsUsAsUsGsUsGsUsTsT	2113

AACAACCCACAAAAUACAACAAGA	2010	29987	FLT1:2340U21 siRNA inv Native sense	AGAACCAACAUAACCAACAAC	2114
AACAACCCACAAAAUACAACAAGA	2010	29988	FLT1:2358L21 siRNA (2340C) Inv Native	UUUUUGGUGUUUUUAUGUUGUU	2115
AACAACCCACAAAAUACAACAAGA	2010	30075	FLT1:2340U21 siRNA sense	CAACCAACAAAAUACAACAATT	2116
AACAACCCACAAAAUACAACAAGA	2010	30076	FLT1:2358L21 siRNA (2340C) antisense	UUUUUGUAUUUUUGUGGUUGTT	2117
AACAACCCACAAAAUACAACAAGA	2010	30077	FLT1:2342U21 siRNA inv	AGAACCAACAUAACCAACAATT	2118
AACAACCCACAAAAUACAACAAGA	2010	30078	FLT1:2358L21 siRNA (2340C) inv	UUUUUGGUGUUUUUAUGUUGTT	2119
AACAACCCACAAAAUACAACAAGA	2010	30187	FLT1:2358L21 siRNA (2340C) 2'-F U.C antisense	uuGuuGuAuuuuuGuGGuuGTT	2120
AACAACCCACAAAAUACAACAAGA	2010	30190	FLT1:2358L21 siRNA (2340C) X = nitroindole antisense	uuGuuGuAuuuuuGuGGuuGXX	2121
AACAACCCACAAAAUACAACAAGA	2010	30193	FLT1:2358L21 siRNA (2340C) Z = nitroindole antisense	uuGuuGuAuuuuuGuGGuuGZZ	2122
AACAACCCACAAAAUACAACAAGA	2010	30196	FLT1:2340U21 siRNA sense IB caps w/2'FY's sense	B cAAccAcAAAAAuAcAAcAAATT B	2123
AACAACCCACAAAAUACAACAAGA	2010	30199	FLT1:2340U21 siRNA sense IB caps sense	cAAccAcAAAAAuAcAAcAAATT	2124
AACAACCCACAAAAUACAACAAGA	2010	30340	FLT1:2358L21 siRNA (2340C) X = 3'dT antisense	uuGuuGuAuuuuuGuGGuuGTX	2125
AACAACCCACAAAAUACAACAAGA	2010	30341	FLT1:2358L21 siRNA (2340C) X = glycyl antisense	uuGuuGuAuuuuuGuGGuuGTX	2126
AACAACCCACAAAAUACAACAAGA	2010	30342	FLT1:2358L21 siRNA (2340C) U = 3'OMeU antisense	uuGuuGuAuuuuuGuGGuuGTU	2127
AACAACCCACAAAAUACAACAAGA	2010	30343	FLT1:2358L21 siRNA (2340C) t = L- dT antisense	uuGuuGuAuuuuuGuGGuuGtT	2128
AACAACCCACAAAAUACAACAAGA	2010	30344	FLT1:2358L21 siRNA (2340C) u = L-rU antisense	uuGuuGuAuuuuuGuGGuuGTu	2129
AACAACCCACAAAAUACAACAAGA	2010	30345	FLT1:2358L21 siRNA (2340C) D = IdT antisense	uuGuuGuAuuuuuGuGGuuGTD	2130
AACAACCCACAAAAUACAACAAGA	2010	30346	FLT1:2358L21 siRNA (2340C) X = 3'dT antisense	uuGuuGuAuuuuuGuGGuuGXT	2131
AACAACCCACAAAAUACAACAAGA	2010	30416	FLT1:2358L21 siRNA (2340C) TsT antisense	uuGuuGuAuuuuuGuGGuuGTsT	2132
UCGUGUAAGGAGUGGACCAUCAU	2013	30777	FLT1:1184U21 siRNA stab04 sense	B GuGuAAGGAGUGGAccAucTT B	2133
UUACGGAGUAUUGCUGUGGGAAA	2014	30778	FLT1:3503U21 siRNA stab04 sense	B AcGGAGuAuGcuUGGGATT B	2134
UAGCAGGCCUUAAGACAUUGAGG	2015	30779	FLT1:4715U21 siRNA stab04 sense	B GcAGGccuAAGAcAuGuGATT B	2135
AGCAAAAAGCAAGGGAGAAAAAGA	2016	30780	FLT1:4753U21 siRNA stab04 sense	B cAAAAAGcAAGGGAGAAAAATT B	2136

UCGUGUAGGAGGAGGACCAUCAU	2013	30781	FLT1:1202L21 siRNA (1184C) stab05 antisense	GauGGuccAcuccuuAcAcTsT	2137
UUACGGAGUAUUGCUGUGGAAA	2014	30782	FLT1:3521L21 siRNA (3503C) stab05 antisense	ucccAcAGcAAuAcuccGuTsT	2138
UAGCAGGCCCUAAGACAUGUGAGG	2015	30783	FLT1:4733L21 siRNA (4715C) stab05 antisense	ucAcAuGuccuuAGGccuGcTsT	2139
AGCAAAAAGCAAGGGAGAAAAGA	2016	30784	FLT1:4771L21 siRNA (4753C) stab05 antisense	uuuuuccuccuuGcuuuuuGTsT	2140
AACAACCACAAAAUACAACAAGA	2010	30955	FLT1:2340U21 siRNA stab07 sense	B cAAccAcAAAAuAcAAcAATT B	2141
AACAACCACAAAAUACAACAAGA	2010	30956	FLT1:2358L21 siRNA (2340C) stab08 antisense	uuGuuGuAuuuuuGuGGuuGTsT	2142
AACAACCACAAAAUACAACAAGA	2010	30963	FLT1:2340U21 siRNA inv	AACAACAUA AAAACACCAACIT	2143
AACAACCACAAAAUACAACAAGA	2010	30964	FLT1:2358L21 siRNA (2340C) inv	GUUGGUGUUUUUAUGUUUUT	2144
AACAACCACAAAAUACAACAAGA	2010	30965	FLT1:2340U21 siRNA stab04 inv	B AAcAAcAuAAAAcAccAAcTT B	2145
AACAACCACAAAAUACAACAAGA	2010	30966	FLT1:2358L21 siRNA (2340C) stab05 inv	GuuGGUGuuuuAuGuuGuuTsT	2146
AACAACCACAAAAUACAACAAGA	2010	30967	FLT1:2340U21 siRNA stab07 inv	B AAcAAcAuAAAAcAccAAcTT B	2147
AACAACCACAAAAUACAACAAGA	2010	30968	FLT1:2358L21 siRNA (2340C) stab08 inv	GuuGGUGuuuuAuGuuGuuTsT	2148
AACUGAGUUUAAAAGGCACCCAG	2009	31182	FLT1:349U21 siRNA TT sense	CUGAGUUUAAAAGGCACCCCT	2149
AAGCAAGGAGGGCCUCUGAUGGU	2012	31183	FLT1:2949U21 siRNA TT antisense	GCAAGGAGGGCCUCUGAUGTT	2150
AGCCUGGAAAGAAUCAAACCUU	2011	31184	FLT1:3912U21 siRNA TT sense	CCUGGAAAGAAUCAAACCCCT	2151
AACUGAGUUUAAAAGGCACCCAG	2009	31185	FLT1:367L21 siRNA (349C) TT antisense	GGGUGCCUUUUAAAACUCAGTT	2152
AAGCAAGGAGGGCCUCUGAUGGU	2012	31186	FLT1:2967L21 siRNA (2949C) TT sense	CAUCAGAGGGCCUCUUGCTT	2153
AGCCUGGAAAGAAUCAAACCUU	2011	31187	FLT1:3930L21 siRNA (3912C) TT antisense	GGUUUUGAUUUCUUUCCAGGTT	2154
AACUGAGUUUAAAAGGCACCCAG	2009	31188	FLT1:349U21 siRNA stab04 sense	B cuGAGuuuuAAAAAGcAccTT B	2155
AAGCAAGGAGGGCCUCUGAUGGU	2012	31189	FLT1:2949U21 siRNA stab04 sense	B GcAAAGGAGGGccuccuGAuGTT B	2156
AGCCUGGAAAGAAUCAAACCUU	2011	31190	FLT1:3912U21 siRNA stab04 sense	B ccuGGAAGAAuAAAAaccTT B	2157
AACUGAGUUUAAAAGGCACCCAG	2009	31191	FLT1:367L21 siRNA (349C) stab05 antisense	GGGuGccuuuuAAAAcucAGTsT	2158
AAGCAAGGAGGGCCUCUGAUGGU	2012	31192	FLT1:2967L21 siRNA (2949C) stab05 antisense	cAucAGAGGGccuccuccuGcTsT	2159
AGCCUGGAAAGAAUCAAACCUU	2011	31193	FLT1:3930L21 siRNA (3912C) stab05 antisense	GGuuuuGAuuuuuccAGGTsT	2160

AACUGAGUUUAAAAGGCACCCAG	2009	31194	FLT1:349U21 siRNA stab07 sense	B cuGAGUUUAAAAGGcAcccTT B	2161
AAGCAAGGAGGGCCUCUGAUGGU	2012	31195	FLT1:2949U21 siRNA stab07 sense	B GcAAGGAGGGccuccuGauGTT B	2162
AGCCUGGAAAGAAUCAAACCCUU	2011	31196	FLT1:3912U21 siRNA stab07 sense	B ccuGGAAGAAuacAAAAccTT B	2163
AACUGAGUUUAAAAGGCACCCAG	2009	31197	FLT1:367L21 siRNA (349C) stab08 antisense	GGGUGccuuuuAAAacucAGTsT	2164
AAGCAAGGAGGGCCUCUGAUGGU	2012	31198	FLT1:3930L21 siRNA (3912C) stab08 antisense	cAucAGAGGccuccuuGcTsT	2165
AGCCUGGAAAGAAUCAAACCCUU	2011	31199	FLT1:349U21 siRNA inv TT	GGUUuUGAuucuuuuccAGGTsT	2166
AACUGAGUUUAAAAGGCACCCAG	2009	31200	FLT1:2949U21 siRNA inv TT	CCCACGGAAAUUUUGAGUCTT	2167
AAGCAAGGAGGGCCUCUGAUGGU	2012	31201	FLT1:3912U21 siRNA inv TT	GUAGUCUCCGGGAGGAACGTT	2168
AGCCUGGAAAGAAUCAAACCCUU	2011	31202	FLT1:367L21 siRNA (349C) inv TT	CcAAAAcUAAGAAAGGUCCTT	2169
AACUGAGUUUAAAAGGCACCCAG	2009	31203	FLT1:2967L21 siRNA (2949C) inv TT	GACUCAAAUUUUCGUGGGTT	2170
AAGCAAGGAGGGCCUCUGAUGGU	2012	31204	FLT1:3930L21 siRNA (3912C) inv TT	CGUUCCUCCCGGAGACUACTT	2171
AGCCUGGAAAGAAUCAAACCCUU	2011	31205	FLT1:349U21 siRNA stab04 inv	GGACCUUUUCUUAUUUUUGGTT	2172
AACUGAGUUUAAAAGGCACCCAG	2009	31206	FLT1:2949U21 siRNA stab04 inv	B cccAcGGAAAUuuuGAGucTT B	2173
AAGCAAGGAGGGCCUCUGAUGGU	2012	31207	FLT1:3912U21 siRNA stab04 inv	B GuAGucuccGGGAGGAACGTT B	2174
AGCCUGGAAAGAAUCAAACCCUU	2011	31208	FLT1:367L21 siRNA (349C) stab05 inv	B ccAAAAcUAAGAAAGGucTT B	2175
AACUGAGUUUAAAAGGCACCCAG	2009	31209	FLT1:2967L21 siRNA (2949C) stab05 inv	GAcucAAAUuuuccGuGGGTsT	2176
AAGCAAGGAGGGCCUCUGAUGGU	2012	31210	FLT1:3930L21 siRNA (3912C) stab05 inv	cGuuccuccGGAGAcuAcTsT	2177
AGCCUGGAAAGAAUCAAACCCUU	2011	31211	FLT1:349U21 siRNA stab07 inv	GGAccuuuuuAGUUuuGGTsT	2178
AACUGAGUUUAAAAGGCACCCAG	2009	31212	FLT1:2949U21 siRNA stab07 inv	B cccAcGGAAAUuuuGAGucTT B	2179
AAGCAAGGAGGGCCUCUGAUGGU	2012	31213	FLT1:3912U21 siRNA stab07 inv	B GuAGucuccGGGAGGAACGTT B	2180
AGCCUGGAAAGAAUCAAACCCUU	2011	31214	FLT1:367L21 siRNA (349C) stab08 inv	B ccAAAAcUAAGAAAGGucTT B	2181
AACUGAGUUUAAAAGGCACCCAG	2009	31215	FLT1:2967L21 siRNA (2949C) stab08 inv	GAcucAAAUuuuccGuGGGTsT	2182
AAGCAAGGAGGGCCUCUGAUGGU	2012	31216	FLT1:3930L21 siRNA (3912C) stab08 inv	cGuuccuccGGAGAcuAcTsT	2183
AGCCUGGAAAGAAUCAAACCCUU	2011	31217	FLT1:349U21 siRNA stab09 sense	GGAccuuuuuAGUUuuGGTsT	2184
AACUGAGUUUAAAAGGCACCCAG	2009	31270	FLT1:2949U21 siRNA stab09 sense	B CUAGUUUAAAAGGCACCCCTT B	2185
AAGCAAGGAGGGCCUCUGAUGGU	2012	31271	FLT1:2949U21 siRNA stab09 sense	B GCAAGGAGGGCCUCUGAUGTT B	2186

AGCCUGGAAAGAAUCAAACCUU	2011	31272	sense FLT1:3912U21 siRNA stab09 sense	B CCUGGAAAGAAUCAAACCTT B	2187
AACUGAGUUUAAAAGGCACCCAG	2009	31273	stab10 antisense FLT1:367L21 siRNA (349C)	GGGUGCCUUUUAAACUCAGTsT	2188
AAGCAAGGAGGGCCUCUGAUGGU	2012	31274	stab10 antisense FLT1:2967L21 siRNA (2949C)	CAUCAGAGGCCCUCCUUGCTsT	2189
AGCCUGGAAAGAAUCAAACCUU	2011	31275	stab10 antisense FLT1:3930L21 siRNA (3912C)	GGUUUUGAUUCUUUCCAGGTsT	2190
AACUGAGUUUAAAAGGCACCCAG	2009	31276	FLT1:349U21 siRNA stab09 inv	B CCCACGGAAAAUUUGAGUCTT B	2191
AAGCAAGGAGGGCCUCUGAUGGU	2012	31277	FLT1:2949U21 siRNA stab09 inv	B GUAGUCUCCGGGAGGAACGTT B	2192
AGCCUGGAAAGAAUCAAACCUU	2011	31278	FLT1:3912U21 siRNA stab09 inv	B CCAAAACUAAGAAAGGUCCTT B	2193
AACUGAGUUUAAAAGGCACCCAG	2009	31279	stab10 inv FLT1:367L21 siRNA (349C)	GACUCAAAUUUUCCGUGGGTsT	2194
AAGCAAGGAGGGCCUCUGAUGGU	2012	31280	stab10 inv FLT1:2967L21 siRNA (2949C)	CGUUCUCCCGGAGACUACTsT	2195
AGCCUGGAAAGAAUCAAACCUU	2011	31281	stab10 inv FLT1:3930L21 siRNA (3912C)	GGACCUUUUCUUAGUUUUUGGTsT	2196
AACAACCCACAAAUAACAACAAGA	2010	31424	stab11 X = 3'-BrdU antisense FLT1:2358L21 siRNA (2340C)	uuGuuGuAuuuuuGuGGuuGXsX	2197
AAGCAAGGAGGGCCUCUGAUGGU	2012	31425	stab11 X = 3'-BrdU sense FLT1:2967L21 siRNA (2949C)	cAucAGAGGGccuccuuGcXsX	2198
AACAACCCACAAAUAACAACAAGA	2010	31442	stab11 X = 3'-BrdU antisense FLT1:2358L21 siRNA (2340C)	uuGuuGuAuuuuuGuGGuuGXsT	2199
AAGCAAGGAGGGCCUCUGAUGGU	2012	31443	stab11 X = 3'-BrdU sense FLT1:2967L21 siRNA (2949C)	cAucAGAGGGccuccuuGcXsT	2200
AACAACCCACAAAUAACAACAAGA	2010	31449	sense FLT1:2340U21 siRNA stab09	B CAACCCACAAAUAACAACAATT B	2201
AACAACCCACAAAUAACAACAAGA	2010	31450	sense FLT1:2340U21 siRNA inv stab09	B AACAAACAUAUAAAACCAACTT B	2202
AACAACCCACAAAUAACAACAAGA	2010	31451	stab10 antisense FLT1:2358L21 siRNA (2340C)	UUGUUGUAUUUUUGUGGUUGTsT	2203
AACAACCCACAAAUAACAACAAGA	2010	31452	inv stab10 antisense FLT1:2358L21 siRNA (2340C)	GUUGGUGUUUUUAUGUUGUUTsT	2204

VEGFR2

Target Pos	Target	Seq ID	Allases	Sequence	Seq ID
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA sense	ACCUUGGAGCAUCUCAUCUTT	2044
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3854U21 siRNA sense	UGAGCAUGGAAGAGGAUUCTT	2045
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA sense	ACCUUUUCCUGUAUGGAGTT	2046
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA sense	CAACACAGCAGGAUUCAGUTT	2047
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) antisense	AGAUGAGAUGCUCUCCAAAGGUTT	2048
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUCUCCAUUGCUCATT	2049
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) antisense	CUCCAUACAGGAAACAGGUTT	2050
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) antisense	ACUGAUUCCUUCUGUUGUTT	2051
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA stab04 sense	B AccuuGGAGcAucucAucUTT B	2052
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3854U21 siRNA stab04 sense	B uGAGcAuGGAAAGAGGAucTT B	2053
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA stab04 sense	B AccuGuuuuccuGuAuGGAGTT B	2054
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA stab04 sense	B cAAcAcAGcAGGAAucAGuTT B	2055
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) stab05 antisense	AGAUGAGAuGcuccAAGGuTsT	2056
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3872L21 siRNA (3854C) stab05 antisense	GAAuccuuccuAuGcuATsT	2057
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) stab05 antisense	cuccAuAcAGGAAAcAGGuTsT	2058
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) stab05 antisense	AcuGAuuuccuGcuGuGuuGTsT	2059
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA stab07 sense	B AccuuGGAGcAucucAucUTT B	2060
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3854U21 siRNA stab07 sense	B uGAGcAuGGAAAGAGGAucTT B	2061
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA stab07 sense	B AccuGuuuuccuGuAuGGAGTT B	2062
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA stab07 sense	B cAAcAcAGcAGGAAucAGuTT B	2063
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) stab11 antisense	AGAUGAGAuGcuccAAGGuTsT	2064
3852	UUUGAGCAUGGAAGAGGAUUCUG	2002	KDR:3872L21 siRNA (3854C) stab11 antisense	GAAuccuuccuAuGcuATsT	2065
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) stab11 antisense	cuccAuAcAGGAAAcAGGuTsT	2066
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) stab11 antisense	AcuGAuuuccuGcuGuGuuGTsT	2067

VEGFR2

Target	SeqID	RP#	Alias	Sequence	SeqID
UGUCCACUUAUCCUGAGGAGCAAG	2017	30785	KDR:3076U21 siRNA stab04 sense	B uccAuuAccuGAGGAGcATT B	2205
UUUGAGCAUGGAAGAGGAUUCUG	2002	30786	KDR:3854U21 siRNA stab04 sense	B uGAGcAuGGAAGAGGAuuCTT B	2053
AUGGUUCUUGCCUCAGAAAGAGCU	2018	30787	KDR:4089U21 siRNA stab04 sense	B GGuuuuGccuAcAGAAAGAGTT B	2206
UCUGAAGGCUCUAAACAGACAAG	2019	30788	KDR:4191U21 siRNA stab04 sense	B uGAAGGcucAAAaccAGAcATT B	2207
UGUCCACUUAUCCUGAGGAGCAAG	2017	30789	KDR:3094L21 siRNA (3076C) stab05 antisense	uGcuccuAcAGGuAAGuGGATsT	2208
UUUGAGCAUGGAAGAGGAUUCUG	2002	30790	KDR:3872L21 siRNA (3854C) stab05 antisense	GAAuccuucAuGcucATsT	2057
AUGGUUCUUGCCUCAGAAAGAGCU	2018	30791	KDR:4107L21 siRNA (4089C) stab05 antisense	cucuucGAGGcAAGAAccTsT	2209
UCUGAAGGCUCUAAACAGACAAG	2019	30792	KDR:4209L21 siRNA (4191C) stab05 antisense	uGucuGGuuuGAGccuucATsT	2210
UGUCCACUUAUCCUGAGGAGCAAG	2017	31426	KDR:3076U21 siRNA sense	UCCACUUACCUUGAGGAGCATT	2211
UUUGAGCAUGGAAGAGGAUUCUG	2002	31427	KDR:3854U21 siRNA sense	UGAGCAUUGGAAGAGGGAUUCCT	2045
AUGGUUCUUGCCUCAGAAAGAGCU	2018	31428	KDR:4089U21 siRNA sense	GGUUCUUGCCUCAGAAAGAGTT	2212
UCUGAAGGCUCUAAACAGACAAG	2019	31429	KDR:4191U21 siRNA sense	UGAAGGCUCUAAACCAGACATT	2213
UGUCCACUUAUCCUGAGGAGCAAG	2017	31430	KDR:3094L21 siRNA (3076C) antisense	UGCUCUCAGGUAAAGUGGATT	2214
UUUGAGCAUGGAAGAGGAUUCUG	2002	31431	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUCUUCUCCAUUCUCATT	2049
AUGGUUCUUGCCUCAGAAAGAGCU	2018	31432	KDR:4107L21 siRNA (4089C) antisense	CUCUUCUGAGGCAAGAACCTT	2215
UCUGAAGGCUCUAAACAGACAAG	2019	31433	KDR:4209L21 siRNA (4191C) antisense	UGUCUUGGUUUUGAGCCUUCATT	2216
UGACCUUGGAGCAUCUCAUCUGU	2001	31434	KDR:3304U21 siRNA sense	ACCUUGGAGCAUCUCAUCUTT	2044
UUUGAGCAUGGAAGAGGAUUCUG	2002	31435	KDR:3854U21 siRNA sense	UGAGCAUGGAAGAGGGAUUCCT	2045
UCACCUUUUCCUGUAUGGAGGA	2003	31436	KDR:3894U21 siRNA sense	ACCUUUUCCUGUAUGGAGTT	2046
GACAACACAGCAGGAAUCAGUCA	2004	31437	KDR:3948U21 siRNA sense	CAACACAGCAGGAAUCAGUTT	2047
UGACCUUGGAGCAUCUCAUCUGU	2001	31438	KDR:3322L21 siRNA (3304C) antisense	AGAUGAGAUUCUCCAAGGUTT	2048
UUUGAGCAUGGAAGAGGAUUCUG	2002	31439	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUUCUCCAUUCUCATT	2049
UCACCUUUUCCUGUAUGGAGGA	2003	31440	KDR:3912L21 siRNA (3894C) antisense	CUCCAUAACAGGAAACAGGUTT	2050
GACAACACAGCAGGAAUCAGUCA	2004	31441	KDR:3966L21 siRNA (3948C) antisense	ACUGAUUCCUGCUGUGUUGTT	2051

(400/102)

VEGFR3

Target Pos	Target	Seq ID	Aliases	Sequence	Seq ID
2009	AGCACUGCCACAAGAAAGUACCUG	2005	FLT4:2011U21 siRNA sense	CACUGCCACAAGAAAGUACCIT	2068
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	FLT4:3921U21 siRNA sense	GAAGCAGAGAGAGAGAGGTT	2069
4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4038U21 siRNA sense	AGAGGAACCAAGGAGGACAATT	2070
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA sense	CAAGAGGAGCAUGAAAGUGTT	2071
2009	AGCACUGCCACAAGAAAGUACCUG	2005	antisense	GGUACUUCUUGUGGCAGUGTT	2072
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	FLT4:3939L21 siRNA (3921C) antisense	CCUUCUCUCUCUCUGCUUCTT	2073
4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4056L21 siRNA (4038C) antisense	UUGUCCUCCUGGUUCCUCUTT	2074
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4072L21 siRNA (4054C) antisense	CACUUUCAUGCUCUCCUUGTT	2075
2009	AGCACUGCCACAAGAAAGUACCUG	2005	FLT4:2011U21 siRNA stab04 sense	B cAcuGccAcAAGAAAGuAccTT B	2076
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	FLT4:3921U21 siRNA stab04 sense	B GAAGcAGAGAGAGAGAGGTT B	2077
4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4038U21 siRNA stab04 sense	B AGAGGAaccAGGAGGAcAATT B	2078
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA stab04 sense	B cAAGAGGAGcAuGAAAGuGTT B	2079
2009	AGCACUGCCACAAGAAAGUACCUG	2005	antisense	GGuAcuucuuGuGcAGuGTsT	2080
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	FLT4:3939L21 siRNA (3921C) stab05 antisense	ccuucucucucucucGcuucTsT	2081
4036	AAAGAGGAACCAAGGAGGACAAGA	2007	antisense	uuGuccucuuGGuuccucuuTsT	2082
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	antisense	cAcuuucAuGcuuccucuuGTsT	2083
2009	AGCACUGCCACAAGAAAGUACCUG	2005	FLT4:2011U21 siRNA stab07 sense	B cAcuGccAcAAGAAAGuAccTT B	2084
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	FLT4:3921U21 siRNA stab07 sense	B GAAGcAGAGAGAGAGAGGTT B	2085
4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4038U21 siRNA stab07 sense	B AGAGGAaccAGGAGGAcAATT B	2086
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA stab07 sense	B cAAGAGGAGcAuGAAAGuGTT B	2087
2009	AGCACUGCCACAAGAAAGUACCUG	2005	antisense	GGuAcuucuuGuGcAGuGTsT	2088
3919	CUGAAGCAGAGAGAGAGAGGCA	2006	antisense	ccuucucucucucucGcuucTsT	2089

4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4056L21 siRNA (4038C) stab11 antisense	uuGuccuccuGGuuccucuTsT	2090
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4072L21 siRNA (4054C) stab11 antisense	cAuuuucAuGcuccuccuuGTsT	2091

Uppercase = ribonucleotide
 u,c = 2'-deoxy-2'-fluoro U,C
 T = thymidine
 B = inverted deoxy abasic
 s = phosphorothioate linkage
 A = deoxy Adenosine
 G = deoxy Guanosine

Table IV

Non-limiting examples of Stabilization Chemistries for chemically modified siNA constructs

Chemistry	pyrimidine	Purine	cap	p=S	Strand
"Stab 1"	Ribo	Ribo	-	5 at 5'-end 1 at 3'-end	S/AS
"Stab 2"	Ribo	Ribo	-	All linkages	Usually AS
"Stab 3"	2'-fluoro	Ribo	-	4 at 5'-end 4 at 3'-end	Usually S
"Stab 4"	2'-fluoro	Ribo	5' and 3'-ends	-	Usually S
"Stab 5"	2'-fluoro	Ribo	-	1 at 3'-end	Usually AS
"Stab 6"	2'-O-Methyl	Ribo	5' and 3'-ends	-	Usually S
"Stab 7"	2'-fluoro	2'-deoxy	5' and 3'-ends	-	Usually S
"Stab 8"	2'-fluoro	2'-O-Methyl	-	1 at 3'-end	Usually AS
"Stab 9"	Ribo	Ribo	5' and 3'-ends	-	Usually S
"Stab 10"	Ribo	Ribo	-	1 at 3'-end	Usually AS
"Stab 11"	2'-fluoro	2'-deoxy	-	1 at 3'-end	Usually AS

CAP = any terminal cap, see for example Figure 10.

All Stab 1-11 chemistries can comprise 3'-terminal thymidine (TT) residues

All Stab 1-11 chemistries typically comprise 21 nucleotides, but can vary as described herein.

S = sense strand

AS = antisense strand

Table V

A. 2.5 μ mol Synthesis Cycle ABI 394 Instrument

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	6.5	163 μ L	45 sec	2.5 min	7.5 min
S-Ethyl Tetrazole	23.8	238 μ L	45 sec	2.5 min	7.5 min
Acetic Anhydride	100	233 μ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	186	233 μ L	5 sec	5 sec	5 sec
TCA	176	2.3 mL	21 sec	21 sec	21 sec
Iodine	11.2	1.7 mL	45 sec	45 sec	45 sec
Beaucage	12.9	645 μ L	100 sec	300 sec	300 sec
Acetonitrile	NA	6.67 mL	NA	NA	NA

B. 0.2 μ mol Synthesis Cycle ABI 394 Instrument

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	15	31 μ L	45 sec	233 sec	465 sec
S-Ethyl Tetrazole	38.7	31 μ L	45 sec	233 min	465 sec
Acetic Anhydride	655	124 μ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	1245	124 μ L	5 sec	5 sec	5 sec
TCA	700	732 μ L	10 sec	10 sec	10 sec
Iodine	20.6	244 μ L	15 sec	15 sec	15 sec
Beaucage	7.7	232 μ L	100 sec	300 sec	300 sec
Acetonitrile	NA	2.64 mL	NA	NA	NA

C. 0.2 μ mol Synthesis Cycle 96 well Instrument

Reagent	Equivalents:DNA/ 2'-O-methyl/Ribo	Amount: DNA/2'-O- methyl/Ribo	Wait Time* DNA	Wait Time* 2'-O- methyl	Wait Time* Ribo
Phosphoramidites	22/33/66	40/60/120 μ L	60 sec	180 sec	360sec
S-Ethyl Tetrazole	70/105/210	40/60/120 μ L	60 sec	180 min	360 sec
Acetic Anhydride	265/265/265	50/50/50 μ L	10 sec	10 sec	10 sec
N-Methyl Imidazole	502/502/502	50/50/50 μ L	10 sec	10 sec	10 sec
TCA	238/475/475	250/500/500 μ L	15 sec	15 sec	15 sec
Iodine	6.8/6.8/6.8	80/80/80 μ L	30 sec	30 sec	30 sec
Beaucage	34/51/51	80/120/120	100 sec	200 sec	200 sec
Acetonitrile	NA	1150/1150/1150 μ L	NA	NA	NA

- 5
- Wait time does not include contact time during delivery.
 - Tandem synthesis utilizes double coupling of linker molecule

CLAIMS

What we claim is:

1. A double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a vascular endothelial growth factor receptor (VEGFr) gene, wherein
5 said siNA molecule comprises about 21 nucleotides.
2. The siNA molecule of claim 1, wherein said siNA molecule comprises no ribonucleotides.
3. The siNA molecule of claim 1, wherein said siNA molecule comprises ribonucleotides.
- 10 4. The siNA molecule of claim 1, wherein one of the strands of said double-stranded siNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of a VEGFr gene, and wherein the second strand of said double-stranded siNA molecule comprises a nucleotide sequence
15 substantially similar to the nucleotide sequence or a portion thereof of said VEGFr gene.
5. The siNA molecule of claim 4, wherein each said strand of the siNA molecule comprises about 19 to about 23 nucleotides, and wherein each said strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.
- 20 6. The siNA molecule of claim 1, wherein said siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of a VEGFr gene, and wherein said siNA further comprises a sense region, wherein said sense region comprises a nucleotide sequence
25 substantially similar to the nucleotide sequence or a portion thereof of said VEGFr gene.
7. The siNA molecule of claim 6, wherein said antisense region and said sense region each comprise about 19 to about 23 nucleotides, and wherein said antisense region comprises at least about 19 nucleotides that are complementary to nucleotides of the sense region.
- 30 8. The siNA molecule of claim 1, wherein said siNA molecule comprises a sense region and an antisense region and wherein said antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA


encoded by a VEGFr gene and said sense region comprises a nucleotide sequence that is complementary to said antisense region.

9. The siNA molecule of claim 6, wherein said siNA molecule is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and the second fragment comprises the antisense region of said siNA molecule.
10. The siNA molecule of claim claim 6, wherein said sense region is connected to the antisense region via a linker molecule.
11. The siNA molecule of claim 10, wherein said linker molecule is a polynucleotide linker.
12. The siNA molecule of claim 10, wherein said linker molecule is a non-nucleotide linker.
13. The siNA molecule of claim 6, wherein pyrimidine nucleotides in the sense region are 2'-O-methyl pyrimidine nucleotides.
14. The siNA molecule of claim 6, wherein purine nucleotides in the sense region are 2'-deoxy purine nucleotides.
15. The siNA molecule of claim 6, wherein the pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides.
16. The siNA molecule of claim 9, wherein the fragment comprising said sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the fragment comprising said sense region.
17. The siNA molecule of claim 16, wherein said terminal cap moiety is an inverted deoxy abasic moiety.
18. The siNA molecule of claim 6, wherein the pyrimidine nucleotides of said antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides.
19. The siNA molecule of claim 6, wherein the the purine nucleotides of said antisense region are 2'-O-methyl purine nucleotides.
20. The siNA molecule of claim 6, wherein the purine nucleotides present in said antisense region comprise 2'-deoxy- purine nucleotides.
21. The siNA molecule of claim 18, wherein said antisense region comprises a phosphorothioate internucleotide linkage at the 3' end of said antisense region.

22. The siNA molecule of claim 6, wherein said antisense region comprises a glyceryl modification at the 3' end of said antisense region.
23. The siNA molecule of claim 9, wherein each of the two fragments of said siNA molecule comprise 21 nucleotides.
- 5 24. The siNA molecule of claim 23, wherein about 19 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule and wherein at least two 3' terminal nucleotides of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule.
- 10 25. The siNA molecule of claim 24, wherein each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines.
26. The siNA molecule of claim 25, wherein said 2'-deoxy-pyrimidine is 2'-deoxy-thymidine.
- 15 27. The siNA molecule of claim 23, wherein all 21 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule.
28. The siNA molecule of claim 23, wherein about 19 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by a VEGFr gene.
- 20 29. The siNA molecule of claim 23, wherein 21 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by a VEGFr gene.
30. The siNA molecule of claim 9, wherein the 5'-end of the fragment comprising said antisense region optionally includes a phosphate group.
- 25 31. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr1.
32. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr2.
33. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr3.
34. A double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGFr gene, wherein said siNA molecule comprises no

ribonucleotides and wherein each strand of said double-stranded siNA molecule comprises about 21 nucleotides.

35. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr1.
36. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr2.
- 5 37. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr3.
38. A double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGFr gene, wherein said siNA molecule does not require the presence of a ribonucleotide within the siNA molecule for said inhibition of expression of the VEGFr gene and wherein each strand of said double-stranded siNA molecule comprises about 21 nucleotides.
- 10 39. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr1.
40. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr2.
41. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr3.
42. A pharmaceutical composition comprising the siNA molecule of claim 1 in an acceptable carrier or diluent.
- 15 43. Medicament comprising the siNA molecule of claim 1.
44. Active ingredient comprising the siNA molecule of claim 1.
45. Use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a VEGFr gene, wherein said siNA molecule comprises one or more chemical modifications and each strand of said double-stranded siNA comprises about 21 nucleotides.
- 20


Figure 1

= SOLID SUPPORT

R = TERMINAL PROTECTING GROUP


FOR EXAMPLE:

DIMETHOXYTRITYL (DMT)





(1)

= CLEAVABLE LINKER

(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR
INVERTED DEOXYABASIC SUCCINATE)

(2)

= CLEAVABLE LINKER

(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR
INVERTED DEOXYABASIC SUCCINATE)INVERTED DEOXYABASIC SUCCINATE
LINKAGE

GLYCERYL SUCCINATE LINKAGE


Figure 2




Figure 3

Figure 4



POSITIONS (NN) CAN COMPRISE ANY NUCLEOTIDE, SUCH AS DEOXYNUCLEOTIDES
(eg. THYMIDINE) OR UNIVERSAL BASES
B = ABASIC, INVERTED ABASIC, INVERTED NUCLEOTIDE OR OTHER TERMINAL CAP
THAT IS OPTIONALLY PRESENT
L = GLYCERYL MOIETY THAT IS OPTIONALLY PRESENT
S = PHOSPHOROTHIOATE OR PHOSPHORODITHIOATE

Figure 5

lower case = 2'-O-Methyl or 2'-deoxy-2'-fluoro

italic lower case = 2'-deoxy-2'-fluorounderline = 2'-O-methyl*ITALIC UPPER CASE* = DEOXY

B = INVERTED DEOXYABASIC

L = GLYCERYL MOIETY OPTIONALLY PRESENT

S = PHOSPHOROTHIOATE OR

PHOSPHORODITHIOATE



Figure 6

Figure 7



[illegible]

3'-EXTENSION

[illegible]


↓
CLEAVAGE WITH RESTRICTION
ENZYMES 1 AND 2

C

5'--

3'--

CLONE



R1 = RESTRICTION SITE #1
R2 = RESTRICTION SITE #2
N = A, G, C, or T
X = A, G, C, or T

Figure 9: Target site Selection using siRNA





Figure 10

R = O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl
 B = Independently any nucleotide base, either naturally occurring or chemically modified, or optionally H (abasic).

Figure 11: Modification Strategy





Figure 12: Inhibition of VEGF-Induced Angiogenesis by siRNAs



Condition	Mean Fit-1 Value (approx.)
Untreated	22
31190/31193	8
31208/31211	18
31183/31186	12
31201/31204	20
31184/31187	8
31202/31205	20
30075/30076	8
30077/30078	18
Scram1	20
Scram2	18
Transfection Control	35

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)